

START

0009976

195

**NATIONAL PRIORITY LIST
U.S. DOE HANFORD 100 AREA**

AUGUST 17, 1987

0
5
6
0
6
0
1
1
1
6



NPL CANDIDATE

Update #

Facility name: U.S. DOE Hanford 100 Area

Received: SEP 14 1987

Location: Hanford Site

EPA Region: X

Person(s) in charge of the facility: J. J. Keating, Asst. Mgr.

Safety, Environment and Security

509 - 376 - 7334

Name of Reviewer: D. M. Bennett, EPA Region X

Date: 8-17-87

General description of the facility:

(For example: landfill, surface impoundment, pile, container; types of hazardous substances; location of the facility; contamination route of major concern; types of information needed for rating; agency action, etc.)

The 100 Area Site contains the 116-B-4, 116-B-6, 116-C-2, 116-D-1A,
116-D-1B, 116-F-5, 116-F-10, 116-H-3, 116-K-2, 116-B-1, 116-B-3, 116-C-1,
116-D-2, 116-DR-1, 116-DR-2, 116-DR-4, 116-F-1, 116-F-2, 116-F-3, 116-F-4,
116-H-1, 116-H-4, 116-K-1, 116-K-2, 116-B-5, 116-B-10, 116-C-2, 116-D-3,
116-D-4, 116-F-9, 116-F-13, 116-B-2, 116-D-1A, 116-D-1B, 116-DR-3, 116-F-3,
116-KE-3, 116-KW-2, 116-DR-6, 116-F-1, 116-F-6, 116-F-12, 116-H-2, 116-K-2,
116-DR-8, 116-F-7, 117-B, 117-C, 117-D, 117-H, 100KE*1, 100KE*2, 100KE*3, 100KW*1
100KW*2, 116-B-9, 116-D-6, 116-F-11, 116-DR-7, 116-KE-1, 116-KW-1, 116-KE-2,
107-B, 107-C, 107-D, 107-DR, 107-F, 107-H, 107-KE, 107-KW, White Bluffs
Pickling Acid Cr1b, 118-B-1, 118-B-4, 118-B-5, 118-C-2, 118-D-1, 118-D-3,
118-D-5, 118-DR-1, 118-F-1, 118-F-3, 118-F-7, 118-H-1, 118-H-2, 118-H-3,

Scores: $S_M = 46.38$ ($S_{GW} = 6.12$ $S_{SW} = 80.00$ $S_a = 0.00$)

$S_{FE} = 0.00$

$S_{DC} = 0.00$

QA
1/11/88
Kath E. Hall
Revised 1/14/88
K

FIGURE 1
HRS COVER SHEET

PLEASE RETURN TO:
ENVIRONMENTAL DIVISION
RESOURCE CENTER

91118900960

118-H-4, 118-H-5, 188-K, 118-B-2, 118-B-3, 118-U-4, 118-F-1, 118-H-3,
118-B-6, 118-F-5, 118-F-6, 118-B-1, 118-B-7, 118-C-1, 118-D-1, 118-D-2,
118-F-2, 118-F-4, 118-H-1, 118-K, 100B/C Burn Pit, 100DR Burn Pit, 100F
Burn Pit, 100N Burn Pit and 100K Burn Pit sites.

FIGURE 1 (Continued)
HRS COVER SHEET

11/1/88
KCA

91118970961

 HRS Ground Water Route Work Sheet

Site: U.S. DOE Hanford 100 Area

8/17/87

Rating Factor	Assigned Value	Multiplier	Score	Max. Score	Ref. (Section)
1. Observed Release	45	1	45	45	3.1

If observed release is given a score of 45, proceed to line 4. If observed release is given a score of 0, proceed to line 2.					

2. Route Characteristics					3.2-
Depth to Aquifer of Concern	0	2	0	6	
Net Precipitation	0	1	0	3	
Permeability of the Unsaturated Zone	0	1	0	3	
Physical State	0	1	0	3	

Total Route Characteristics Score			0	15	

3. Containment	0	1	0	3	3.3

4. Waste Characteristics					3.4
Chemical					
a. Toxicity/Persistence	18	1	18	18	
Hazardous Waste Quantity	8	1	8	8	

Total Waste Characteristics Score			26	26	

5. Targets					3.5
Ground Water Use	1	3	3	9	
Distance to Nearest Well/ Population Served	0	1	0	40	

Total Targets Score			3	59	

6. If line 1=45 (1x4x5) If line 1=0 (2x3x4x5)			3510	57330	

7. Line 6/57330 * 100	Sc(gw)=	6.12			

*Revised 1/14/88
 1/11/88
 KJ*

9 1 1 1 8 9 7 0 9 5 9

 HRS Surface Water Route Work Sheet
 =====

Site: U.S. DOE Hanford 100 Area

8/17/87

Rating Factor	Assigned Value	Multiplier	Score	Max. Score	Ref. (Section)
1. Observed Release	45	1	45	45	4.1
If observed release is given a score of 45, proceed to line 4. If observed release is given a score of 0, proceed to line 2.					
2. Route Characteristics					4.2
Facility Slope & Intervening Terrain	0	1	0	3	
1-yr. 24-hr. Rainfall	0	1	0	3	
Distance to Nearest Surface Water	0	2	0	6	
Physical State	0	1	0	3	
Total Route Characteristics Score			0	15	
3. Containment	0	1	0	3	4.3
4. Waste Characteristics					4.4
a. Chemical					
Toxicity/Persistence	18	1	18	18	
Hazardous Waste Quantity	8	1	8	8	
Total Waste Characteristics Score			26	26	
5. Targets					4.5
Surface Water Use	3	3	9	9	
Distance to a Sensitive Environment	0	2	0	6	
Population Served/Distance to Water Intake Downstream	35	1	35	40	
Total Targets Score			44	55	
6. If line 1=45 (1x4x5) If line 1=0 (2x3x4x5)			51480	64350	
7. Line 6/64350 * 100	Sc(sw)=		80.00		

*1/11/88
XCD*

9111097093

	S	S ²
Groundwater Route Score (S _{gw})	6.12	37.45
Surface Water Route Score (S _{sw})	80.00	6400.00
Air Route Score (S _a)	0.00	0.00
$S_{gw}^2 + S_{sw}^2 + S_a^2$	-	6437.45
$\sqrt{S_{gw}^2 + S_{sw}^2 + S_a^2}$	-	80.23
$\sqrt{S_{gw}^2 + S_{sw}^2 + S_a^2} / 1.73 = S_M =$	-	46.38

FIGURE 10
WORKSHEET FOR COMPUTING S_M

Revised 1/14/88
KEA

1/11/88
KEA

91118900965

DOCUMENTATION RECORDS
FOR
HAZARD RANKING SYSTEM

INSTRUMENTATIONS: As briefly as possible summarize the information you used to assign the score for each factor (e.g., "Waste quantity = 4,230 drums plus 800 cubic yards of sludge"). The source of information should be provided for each entry and should be a bibliographic-type reference, include the location of the document.

FACILITY NAME: U.S. DOE Hanford 100 Area

LOCATION: Hanford Site, Benton County, Washington

DATE SCORED: 8-17-87

PERSON SCORING: R. D. Stenner, Pacific Northwest Laboratory for DOE

PRIMARY SOURCE(S) OF INFORMATION (e.g., EPA region, state, FIT, etc.):

The information was taken from Department of Energy documents and databases associated with the Hanford Site, as well as from other publicly available documents addressing conditions at or in the vicinity of the Hanford Site. Information was also gathered through telephone and personal communications with responsible individuals (such information is referenced accordingly in the package).

FACTORS NOT SCORED DUE TO INSUFFICIENT INFORMATION:

Even though air concentrations of some of the constituents of interest can be detected above background offsite, no air monitoring data were found sufficient for HRS scoring of the Hanford CERCLA sites. These constituents of interest detected above background offsite are present in the routine gaseous effluents from operating facilities at Hanford. Therefore, the air route rating factors were not scored.

COMMENTS OR QUALIFICATIONS:

The Department of Energy has completed a preliminary assessment of the hazardous waste sites located on the Hanford Site. This work served as the primary basis for developing the scores for the aggregate 100 Area Site. These preliminary assessment efforts are documented in the "Draft Phase I Installation Assessment of Inactive Waste-Disposal Sites at Hanford" volumes and respective addenda which are referenced throughout the package.

QA
11/11/84
Kathleen D. Young

911390966

GROUND WATER ROUTE

1 OBSERVED RELEASE

Contaminants detected (5 maximum):

An observed release to the ground water beneath the 100 Areas can be verified using several different substances. For the purpose of documenting observed release to the ground water it was determined that the strontium-90 (Sr-90) and chromium (Cr) would provide sufficient evidence. Evidence of observed release can be given in terms of a comparison of background concentrations and down gradient concentrations for both Sr-90 and Cr. Wells used to verify observed release of Sr-90 are listed in the following table.

100 Area Wells for Sr-90

BACKGROUND			DOWN GRADIENT		
Date	Well	Concentration (pCi/L)	Date	Well	Concentration (pCi/L)
4/10/87	1-H4-5	<0.753 pCi/L	4/10/87	1-H4-4	8.05 pCi/L
			5/29/87	1-N-4	12.5 pCi/L

The wells used to verify observed release of Cr to the ground water beneath the 100 Area are listed in the following table.

100 Area Wells for Cr

BACKGROUND			DOWN GRADIENT		
Date	Well	Concentration (µg/L)	Date	Well	Concentration (µg/L)
6/12/87	6-77-54	<10 µg/L	5/21/87	1-B3-1	62 µg/L
			3/25/87	1-D5-12	1,560 µg/L
			3/18/87	1-K-20	137 µg/L

Reference 5, page 3.14; Reference 2; Reference 3, pages 19, 23, 274, 13,16, and 21; Reference 1; Reference 10; Reference 18

Rationale for attributing the contaminants to the facility:

The nine 100 Areas (B, C, D, DR, KE, KW, F, H, and N) border the Columbia River in the northernmost part of the Hanford Site. Each of the nine areas has one production reactor. Eight of these reactors have been shut down; only the N Reactor, used for both plutonium and electricity production is still in operation. When the reactors were operational, cooling water was drawn from the river and treated with alum, sulfuric acid and chlorine. Excess sulfuric acid was used to maintain the pH of the water within a desired range. To control oxidation of aluminum parts in the reactor, sodium dichromate was used to maintain an oxidation coating on aluminum parts. The chlorine was added for algae control in the settling basins; at times copper

11/11/89
XCA

9 1 1 0 9 9 6 8

sulfate was added for additional algae control. As the cooling water passed through the reactor various activation products (e.g, Co-60, Ni-63, etc.) were dissolved in the cooling water. When a fuel element ruptured or developed a pin hole in its' cladding some fission products such as Sr-90 would dissolve in the cooling water. Chromic acid, oxalic acid and nitric acid were used for dummy fuel-element decontamination. In addition to vertical safety rods for emergency reactor shutdown, the reactors were equipped with hoppers of nickel-plated boron steel balls, nickel-plated carbon steel balls, and stainless steel balls that would drop into the vertical safety rod channels for emergency shutdown. This system required no supplementary power source. Although it was never used, a third safety system, one involving the use of potassium borate solution, was in place at the reactors. A supplementary control system, in addition to the normal horizontal control rods, was incorporated into the reactors. This supplementary control system consisted of a Poison(a) Column Control Facility that could charge selected process tubes with a lead-cadmium poison to absorb neutrons. Boron-carbide aluminum poison splines were also used for supplementary control. These activities resulted in waste containing the contaminants discussed above being disposed of in the 100 Area. Down gradient wells show contaminant concentrations significantly above those found in the background wells.

(a) The term poison refers to a material's ability to absorb neutrons and thus control the rate of fission.

Reference 4, pages 2.11-2.13

2 ROUTE CHARACTERISTICS

Depth to Aquifer of Concern

Name/description of aquifer(s) of concern:

The aquifer of concern is the unconfined aquifer which is comprised of the glacio fluvial sediments of the Hanford formation and the late deposits of the Ringold formation. It generally slopes downward from west to east; depth to ground water is from 10 to 15 meters (34 to 48 feet). It is bounded below by either the basalt surface or, in places, the relatively impervious clays and silts of the lower unit of the Ringold formation. Laterally, the unconfined aquifer is bounded by the anticlinal basalt ridges that ring the basin. The Yakima and Columbia Rivers, however, do not entirely transect these sediments and therefore do not constitute a discontinuity for HRS Scoring purposes. The basalt ridges above the water table have a low permeability and act as a barrier to lateral flow of the ground water. The saturated thickness of the unconfined aquifer is greater than 61 m in some areas of the Hanford Site and pinches out along the flanks of the basalt anticlines.

Recharge to the unconfined aquifer originates from several sources. Natural recharge occurs from precipitation at higher elevations and runoff from

ephemeral streams to the west, such as Cold Creek and Dry Creek. The Yakima River recharges the unconfined aquifer as it flows along the southwest boundary of the Hanford Site. The Columbia River recharges the unconfined aquifer during high stages when river water is transferred to the aquifer along the river bank. The unconfined aquifer receives little recharge from precipitation directly on the Hanford Site because of a high rate of evapotranspiration under native soil and vegetation conditions. Large scale artificial recharge occurs from offsite agricultural irrigation and liquid-waste disposal in the operating areas at Hanford.

Underlying the surface sands is a mixture of sand and gravel extending to a depth of about 200 feet (60 meters). Basaltic rock starts at that depth and extends downward over 1.9 miles (3000 meters).

Reference 5, page 2.5; Reference 4, pages 2.11-13; Reference 9

Depth(s) from the ground surface to the highest seasonal level of the saturated zone [water table(s)] of the aquifer of concern:

Not applicable.

Depth from the ground surface to the lowest point of waste disposal/storage:

Not applicable.

1/11/88
KCA

9
5
6
6
6
1
1
1
6

Net Precipitation

Mean annual or seasonal precipitation (list months for seasonal):

Not applicable.

Mean annual lake or seasonal evaporation (list months for seasonal):

Not applicable.

Net precipitation (subtract the above figures):

Not applicable.

Permeability of Unsaturated Zone

Soil type in unsaturated zone:

Not applicable.

Permeability associated with soil type:

Not applicable

Physical State

Physical state of substances at time of disposal (or at present time for generated gases):

Not applicable.

1/11/88
K2A

0
2
6
0
0
0
0
0
1
1
1
9

3 CONTAINMENT

Containment

Method(s) of waste or leachate containment evaluated:

Not applicable.

Method with highest score:

Not applicable.

4 WASTE CHARACTERISTICS

Toxicity and Persistence

Compound(s) evaluated:

These substances are associated with production reactor operation (See Tables 1 and 2):

Substances in the 100 Area

uranium	mercury	cesium-137
lead	plutonium-238	cesium-134
cadmium	plutonium-239	europium-152
sodium dichromate	plutonium-240	europium-154
sodium oxalate	tritium	europium-155
sulfuric acid	cobalt-60	sodium sulfamate
sulfamic acid	strontium-90	nickel-63
potassium borate	copper sulfate	sodium hydroxide

Reference 4, pages 2.11-2.13; Reference 6 (The numerous individual pages are not included in the package, please consult the reference 6 document using the list of individual sites listed on the cover sheet of this package for verification of which individual sites contain which substances); Reference 19

Compound with highest score:

Several of these substances results in a score of 18. Uranium, lead, mercury sodium dichromate and plutonium are among those having scores of 18.

<u>Substance</u>	<u>Toxicity Score</u>	<u>Persistence Score</u>	<u>TOTAL Score</u>
uranium	3	3	18
lead	3	3	18
mercury	3	3	18
dichromate	3	3	18
plutonium	3	3	18

11/11/88
ZCJ

Reference 7, pages 794-797; REference 20.

120000711113

Hazardous Waste Quantity

Total quantity of hazardous substances at the facility, excluding those with a containment score of 0 (Give a reasonable estimate even if quantity is above maximum):

The total quantity of hazardous substances associated with the aggregate 100 Area Site is estimated to be 4.3 billion cubic yards. Table 1 presents the individual sites and associated data for the 100 Area Liquid Waste Sites that were used in creating the aggregate 100 Area Site. Table 2 presents the individual sites and associated data for the 100 Area Solid Waste Sites that were used in creating the aggregate 100 Area Site.

References are listed by waste site in Table 1.

Basis of estimating and/or computing waste quantity:

The volume of wastes disposed of to waste sites at the 100 Area was calculated as the sum of the volumes received by individual waste sites. The basis for this estimate of liquid waste volume is given in Table 1. For purposes of developing the inventories, individual waste sites were aggregated into groups that received similar wastes. These groups, and the individual sites comprising each group, are identified in Table 1. Also included in this table are a description of the general types of waste received by the group of sites, the period of operation covered by the group, the estimated waste volume received by the group, and the references for the waste volumes received by the individual sites. In some cases, an individual site received more than one type of waste. If so, the site was assigned to more than one group and its waste volume was divided into the volumes received by each particular site group. Similar information for 100 Area sites receiving solid wastes is presented in Table 2.

1/11/88
KZ

TABLE 1. 100 AREA LIQUID WASTE SITES

<u>Area No. and Name</u>	<u>Waste Types</u>	<u>Years</u>	<u>Estimated Volume</u>	<u>No. of Cubic Yds</u>	<u>References</u>
100 Area Decontamination Waste Cribs (116-B-4, 116-B-6, 116-C-2, 116-D-1A, 116-D-1B, 116-F-5, 116-F-10, 116-H-3, 116-K-2)	Decontamination solutions consisting of chromic, oxalic, and/or sulfuric acid solutions neutralized with soda ash. Solutions were used to decontaminate dummy fuel elements and reactor hardware.	1947-1970	2.3×10^6 L	3,000	Reference 6, pages 13-14, 17-20, 45-48, 59-60, 61-62, 113-114, 121-122, 151-152, 173-174
100 Area Contaminated Reactor Effluent Cribs and Trenches (116-B-1, 116-B-3, 116-C-1, 116-D-2, 116-DR-1, 116-DR-2, 116-DR-4, 116-F-1, 116-F-2, 116-F-3, 116-F-4, 116-H-1, 116-H-4, 116-K-1, 116-K-2)	Reactor effluent contaminated when fuel element cladding failed. Radiological contamination consisted chiefly of mixed fission products. Chemical contamination consisted of sodium dichromate added to reactor coolant as oxidant.	1946-1971	3.0×10^{11} L	3.9×10^8	Reference 6, pages 7-8, 11-12, 43-44, 63-64, 87-88, 89-90, 93-94, 105-106, 107-108, 109-110, 111-112, 147-148, 153-154, 171-172, 173-174

11/11/88
NCR

<u>Area No. and Name</u>	<u>Waste Types</u>	<u>Years</u>	<u>Estimated Volume</u>	<u>No. of Cubic Yds</u>	<u>References</u>
100 Area Laboratory Waste Cribs and Trenches (116-B-5, 116-B-10, 116-C-2, 116-D-3, 116-D-4, 116-F-9, 116-F-13)	Wastes associated with labs and shops in the 108 Buildings, including 108-B Tritium Lab and Tube Examination Facility, 108-C Metal Examination Facility and 100 F Biology Labs. Contaminants consisted mainly of tritium and low levels of mixed fission and activation products. Negligible chemical contamination except Biology Lab facilities, which received nitrate from animal wastes.	1950-1976	3.2×10^8 L	4.2×10^5	Reference 6, pages 15-16, 23-24, 45-48, 65-66, 67-68, 119-120, 127-128
100 Area Fuel Storage Basin Cribs and Trenches (116-B-2, 116-D-1A, 116-D-1B, 116-DR-3, 116-F-3, 116-KE-3, 116-KW-2)	Storage basin cooling water contaminated with low levels of mixed fission products due to failure of fuel element cladding.	1946-1971	2.0×10^7 L	2.6×10^4	Reference 6, pages 9-10, 59-60, 61-62, 91-92, 109-110, 179-180, 195-196

11/11/98
 881198
 881198

<u>Area No. and Name</u>	<u>Waste Types</u>	<u>Years</u>	<u>Estimated Volume</u>	<u>No. of Cubic Yds</u>	<u>References</u>
100 Area Reactor Decontamination and Maintenance Effluents Cribs and Trenches (116-DR-6, 116-F-1, 116-F-6, 116-F-12, 116-H-2, 116-K-2)	Reactor effluents during reactor maintenance activities. Much of the waste volume was uncontaminated. Radionuclides present were mainly low levels of mixed fission products and activation products. The main chemical contaminant was sulfuric acid, which was used for reactor decontamination. Sodium dichromate was also present in some of the effluents.	1945-1971	6.1 x 10 ⁸ L	8.0 x 10 ⁵	Reference 6, pages 95-96, 105-106, 115- 116, 125-126, 149-150, 173-174
100 Area Reactor Confinement Seal Pit Drainage Cribs (116-DR-8, 116-F-7, 117-B, 117-C, 117-D, 117-H)	Drainage from 117 Building confinement system seal pits. Very low levels of contamination.	1960-1969	2.2 x 10 ⁶ L	2,900	Reference 6, pages 42-44, 49-50, 85-86, 99-100, 117-118, 167-168
100 Area Sulfuric Acid Sludge Disposal Sites (100KE*1, 100KE*2, 100KE*3, 100KW*1, 100KW*2)	Mercury-contaminated sludge from sulfuric acid storage tanks.	1955-1971	3.7 x 10 ⁴ L	48	Reference 6, pages 181-183, 185-187, 189- 191, 197-199, 201-203

11/1/88
RCC

<u>Area No. and Name</u>	<u>Waste Types</u>	<u>Years</u>	<u>Estimated Volume</u>	<u>No. of Cubic Yds</u>	<u>References</u>
100 Area Miscellaneous Floor and Sink Drain Cribs (116-B-9, 116-D-6, 116-F-11)	Drainage from floor and sink drains. Very low levels of radioactive contamination. No chemical contamination.	1952-1967	3.4×10^5 L	440	Reference 6, pages 21-22, 69-70, 123-124
100 Area Potassium Borate Crib (116-DR-7)	Liquid potassium borate solution. No radioactive contamination.	1953	4×10^3 L	5.2	Reference 6, pages 97-98
100 Area Gas Purification Condensate Cribs (116-KE-1, 116-KW-1)	Condensate from reactor gas purification system. Contaminated with tritium and carbon-14 and small amounts of fission products.	1955-1971	1.6×10^6 L	2.1×10^3	Reference 6, pages 175-176, 193-194
100 Area Ion Exchange Regenerant Crib (116-KE-2)	Wastes from regeneration of ion exchange columns. Contained sulfuric acid, caustic, and mixed fission products and activation products.	1955-1971	3.0×10^6 L	3.9×10^3	Reference 6, pages 177-178

11/11/88
NLP

<u>Area No. and Name</u>	<u>Waste Types</u>	<u>Years</u>	<u>Estimated Volume</u>	<u>No. of cubic Yds</u>	<u>References</u>
100 Area Retention Basin Leakage (107-B, 107-C, 107-D, 107-DR, 107-F, 107-H, 107-KE, 107-KW)	Reactor effluent containing very small amounts of fission products and activation products. Chemical contamination limited to small amounts of sodium dichromate added to reactor coolant as oxidant.	1944-1971	3.0×10^{12} L	3.9×10^9	Reference 16
100 Area White Bluffs Pickling Acid Crib	Waste nitric and hydrofluoric acid used to pickle galvanized piping. Acid may or may not have been neutralized.	1943-1945	7×10^5 L	900	Reference 16
		TOTAL	3.3×10^{12} L	4.3×10^9	

11/11/88
 10/19/88

TABLE 2. 100 AREA SOLID WASTE SITES

<u>Area No. and Name</u>	<u>Waste Types</u>	<u>Years</u>	<u>Estimated Volume</u>	<u>No. of Cubic Yds</u>	<u>References</u>
100 Area Metallic Waste Burial Grounds (118-B-1, 118-B-4, 118-B-5, 118-C-2, 118-D-1, 118-D-3, 118-D-5, 118-DR-1, 118-F-1, 118-F-3, 118-F-7, 118-II-1, 118-II-2, 118-II-3, 118-II-4, 118-II-5, 118-K)	Metallic waste consisting mainly of discarded reactor hardware, including such items as aluminum tubes and spacers; lead and cadmium containing slugs, graphite, desiccant, aluminum and boron splines; lead; and mercury. Radioactive contaminants, primarily activation products.	1944-1975	1,000 m ³	1,300	Reference 6, pages 25-27, 33-34, 35-36, 55-56, 71-72, 77-79, 83-84, 101-102, 129-131, 135-136, 143-144, 155-157, 159-160, 161-162, 163-164, 165-166, 205-207
100 Area Construction Waste Burial Grounds (118-B-2, 118-B-3, 118-D-4, 118-F-1, 118-II-3)	Construction waste, demolition debris. Small amounts of radioactive contamination.	1953-1967	48,000 m ³	63,000	Reference 6, pages 29-30, 31-32, 81-82, 129-131, 161-162
100 Area Lab Waste Burial Grounds (118-B-6, 118-F-5, 118-F-6)	Low-level radioactive wastes from 100 Area laboratories, including tritium wastes and animal wastes.	1950-1975	20,000 m ³	26,000	Reference 6, pages 37-38, 139-140, 141-142

11/11/88
KCB

<u>Area No. and Name</u>	<u>Waste Types</u>	<u>Years</u>	<u>Estimated Volume</u>	<u>No. of Cubic Yds</u>	<u>References</u>
100 Area Miscellaneous Solid Waste Burial Grounds (118-B-1, 118-B-7, 118-C-1, 118-D-1, 118-D-2, 118-F-2, 118-F-4, 118-II-1, 118-K)	Radioactivity contaminated trash and miscellaneous solid wastes from 100 Area facilities.	1944-1975	61,000 m ³	80,000	Reference 6, pages 25-27, 39-40, 51-53, 71-72, 73-75, 133-134, 137-138, 155-157, 205-207
100 Area Burn Pits (100 B/C Burn Pit, 100/DR Burn Pit, 100F Burn Pit, 100N Burn Pit, 100K Burn Pit)	Combustible trash from 100 Area buildings and offices.	1943-1971	130,000 m ³	170,000	Reference 6, pages 57-58, 103-104, 145-146, 169-170, 209-210
			<u>TOTAL</u>	<u>260,000 m³</u>	<u>340,000 yd³</u>

11/11/88
KEA

5 TARGETS

Ground Water Use

Use(s) of aquifer(s) of concern within a 3-mile radius of the facility:

There are no wells which draw water for drinking or irrigation purposes within 3 miles of the 100 Area. Therefore, there is no use of the aquifer of concern within 3 miles of the 100 Area.

Reference 8

Distance to Nearest Well

Location of nearest well drawing from aquifer of concern or occupied building not served by a public water supply:

The nearest wells are 699-72-101A, B, and C that are located at the Vernita Bridge Rest Stop.

Reference 8; Reference 9, page II.1-65

Distance to above well or building:

The wells located at the Vernita Bridge Rest Stop are just over 3 miles from the boundary of the 300 Area Site.

Reference 8

Population Served by Ground Water Wells Within a 3-Mile Radius

Identified water-supply well(s) drawing from aquifer(s) of concern within a 3-mile radius and populations served by each:

There are no wells which draw water for drinking or irrigation purposes within 3 miles of the 100 Area. Therefore, there is no use of the aquifer of concern within 3 miles of the 100 Area.

Reference 8

Computation of land areas irrigated by supply well(s) drawing from aquifer(s) of concern within a 3-mile radius, and conversion to population (1.5 people per acre):

There are no documented irrigation wells within 3 miles of the site.

Total population served by ground water within a 3-mile radius:

The total population served by ground water within 3 miles is zero.

1/11/88
ZLN

006006116

SURFACE WATER ROUTE

1 OBSERVED RELEASE

Contaminants detected in surface water at the facility or downhill from it (5 maximum):

The observed release of contaminants to the Columbia River adjacent to the 100 Area can be verified by comparison of upstream and downstream concentrations of selected constituents. Verification of observed release does not require evidence of release for each contaminant disposed at the facility, but relies on identification of at least one constituent in the surface water that can be attributed to the facility. Strontium-90 (Sr-90) was chosen to illustrate observed release from the 100 Area. The following table compares upstream concentrations of Sr-90 with concentrations of Sr-90 at designated surface water use points. The water intakes designated as use points are in the 100N and 100D areas.

BACKGROUND			DOWN GRADIENT(a)		
Date	Sample Point	Concentration (pCi/L)	Date	Sample Point	Concentration (pCi/L)
11/23/82-	Prst Rap	0.34 ± 0.18	12/18/82	100N	28 ± 0.47
12/21/82			12/18/82	100D	1.1 ± 0.05

(a) This data represents an analysis of the composite of river water samples taken every 0.5 mile for the river interval indicated on the date shown.

Reference 11; Reference 5, page 3.14; Reference 12, pages 15 and 8.3; Reference 10

Rationale for attributing the contaminants to the facility:

The nine 100 Areas (B, C, D, DR, KE, KW, F, H, and N) border the Columbia River in the northernmost part of the Hanford Site. Each of the nine areas has one production reactor. Eight of these reactors have been shut down; only the N Reactor, used for both plutonium and electricity production is still in operation. When the reactors were operational, cooling water was drawn from the river and treated with alum, sulfuric acid and chlorine. Excess sulfuric acid was used to maintain the pH of the water within a desired range. To control oxidation of aluminum parts in the reactor, sodium dichromate was used to maintain an oxidation coating on aluminum parts. The chlorine was added for algae control in the settling basins; at times copper sulfate was added for additional algae control. As the cooling water passed through the reactor various activation products (e.g, Co-60, Ni-63, etc.) were dissolved in the cooling water. When a fuel element ruptured or developed a pin hole in its' cladding some fission products such as Sr-90 would dissolve in the cooling water. Chromic acid, oxalic acid and nitric

11/11/88
XCD

acid were used for dummy fuel-element decontamination. In addition to vertical safety rods for emergency reactor shutdown, the reactors were equipped with hoppers of nickel-plated boron steel balls, nickel-plated carbon steel balls, and stainless steel balls that would drop into the vertical safety rod channels for emergency shutdown. This system required no supplementary power source. Although it was never used, a third safety system, one involving the use of potassium borate solution, was in place at the reactors. A supplementary control system, in addition to the normal horizontal control rods, was incorporated into the reactors. This supplementary control system consisted of a Poison^(a) Column Control Facility that could charge selected process tubes with a lead-cadmium poison to absorb neutrons. Boron-carbide aluminum poison splines were also used for supplementary control.

These activities resulted in strontium-90 contaminated waste being disposed of in the 100 Area, and down gradient samples show strontium-90 concentration levels significantly above the strontium-90 concentrations found in background samples.

- (a) The term poison refers to a material's ability to absorb neutrons and thus control the rate of fission.

Reference 4, pages 2.11-2.13

2 ROUTE CHARACTERISTICS

Facility Slope and Intervening Terrain

Average slope of facility in percent:

Not applicable.

Name/description of nearest down slope surface water:

The Columbia River, which originates in the mountains of eastern British Columbia, Canada, flows through the northern edge of the Hanford Site and forms part of the Hanford Site's eastern boundary. The river drains a total area of approximately 70,800 km² enroute to the Pacific Ocean. The flow of the Columbia River is regulated by 11 dams within the United States, 7 upstream and 4 downstream of the Site. Priest Rapids Dam is the nearest impoundment upstream of the Site, and McNary Dam is the nearest dam downstream. (The Hanford reach of the Columbia River extends from Priest Rapids Dam to the head of Lake Wallula, which is created by McNary Dam.) This is the only stretch of the Columbia River within the U.S. that is not impounded by a dam. The width of the river varies from approximately 300 m to about 1000 m. The flow through this stretch of the river is relatively swift, with numerous bends and several islands present throughout the reach. The ground water beneath the site discharges directly into the Columbia River as evidenced by seeps and springs along the river shore.

11/11/88
R27

The flow rate of the Columbia River in this region is regulated primarily by Priest Rapids Dam. Hanford reach flows fluctuate significantly because of the relatively small storage capacity and operational practices of the nearby upstream dams. A minimum flow rate of 1,000 cubic meters per second (36,000 cubic feet per second) has been established at Priest Rapids. Typical daily flows range from 1,000 cubic meters per second (36,000 cubic feet per second) to 7,000 cubic meters per second (250,000 cubic feet per second) with peak spring runoff flows of up to 12,600 cubic meters per second (450,000 cubic feet per second) being recorded. Typical annual average flows at Priest Rapids Dam are 3,100 cubic meters per second (110,000 cubic feet per second) to 3,400 cubic meters per second (120,000 cubic feet per second). Monthly mean flows typically peak from April through June and are at the lowest levels from September through October.

Reference 5, page 2.1-2.3

Average slope of terrain between facility and above-cited surface water body in percent:

Not applicable.

Is the facility located either totally or partially in surface water?

Not applicable.

1/11/88
KIA

Is the facility completely surrounded by areas of higher elevation?

Not applicable.

1-Year 24-Hour Rainfall in Inches

Not applicable.

Distance to Nearest Down slope Surface Water

Not applicable.

Physical State of Waste

Not applicable.

3 CONTAINMENT

Containment

Method(s) of waste or leachate containment evaluated:

Not applicable.

Method with highest score:

Not applicable.

1/11/88
KED

4 WASTE CHARACTERISTICS

Toxicity and Persistence

Compound(s) evaluated

These substances are associated with production reactor operation (See Tables 1 and 2):

Substances in the 100 Area

uranium	mercury	cesium-137
lead	plutonium-238	cesium-134
cadmium	plutonium-239	europium-152
sodium dichromate	plutonium-240	europium-154
sodium oxalate	tritium	europium-155
sulfuric acid	cobalt-60	sodium sulfamate
sulfamic acid	strontium-90	nickel-63
potassium borate	copper sulfate	sodium hydroxide

Reference 4, pages 2.11-2.13; Reference 6 (The numerous individual pages are not included in the package, please consult the reference 6 document using the list of individual sites listed on the cover sheet of this package for verification of which individual sites contain which substances); Reference 19

Compound with highest score:

Several of these substances results in a score of 18. Uranium, lead, mercury sodium dichromate and plutonium are among those having scores of 18.

<u>Substance</u>	<u>Toxicity Score</u>	<u>Persistence Score</u>	<u>TOTAL Score</u>
uranium	3	3	18
lead	3	3	18
mercury	3	3	18
dichromate	3	3	18
plutonium	3	3	18

Reference 7, pages 794-797; Reference 20.

Hazardous Waste Quantity

Total quantity of hazardous substances at the facility, excluding those with a containment score of 0 (Give a reasonable estimate even if quantity is above maximum):

The total quantity of hazardous substances associated with the aggregate 100

1/11/88
RCA

Area Site is estimated to be 4.3 billion cubic yards. Table 1 presents the individual sites and associated data for the 100 Area Liquid Waste Sites that were used in creating the aggregate 100 Area Site. Table 2 presents the individual sites and associated data for the 100 Area Solid Waste Sites that were used in creating the aggregate 100 Area Site.

References are listed by waste site in Table 1.

Basis of estimating and/or computing waste quantity:

The volume of wastes disposed of to waste sites at the 100 Area was calculated as the sum of the volumes received by individual waste sites. The basis for this estimate of liquid waste volume is given in Table 1. For purposes of developing the inventories, individual waste sites were aggregated into groups that received similar wastes. These groups, and the individual sites comprising each group, are identified in Table 1. Also included in this table are a description of the general types of waste received by the group of sites, the period of operation covered by the group, the estimated waste volume received by the group, and the references for the waste volumes received by the individual sites. In some cases, an individual site received more than one type of waste. If so, the site was assigned to more than one group and its waste volume was divided into the volumes received by each particular site group. Similar information for 100 Area sites receiving solid wastes is presented in Table 2.

In compiling the inventory, since groundwater seeps into the Columbia River, it is assumed that the containment associated with the surface water route is considered not to be 0, and all the waste quantities cited were available for migration.

Reference 12; Reference 19.

5 TARGETS

Surface Water Use

Use(s) of surface water within 3 miles downstream of the hazardous substance:

The Columbia River is used as the source of drinking and process water for the 100 and 200 areas. The intake at 100-B Pumphouse supplies water to 100-B/C, 100-D, 100-N, 100-K, and the 200 areas. The intake at 100-D is used as a backup to 100-B intake. The 100-N intake supplies water to 100-N area and Washington Public Power Supply System Generation Station attached to the 100-N reactor. The 100-K Pumphouse supplies water to 100-K area.

The Columbia River both near and down stream of the 100 Area is used for recreation.

Reference 9, pages II.1-64-II.1-65; Reference 13, page 3; Reference 17.

11/11/88
KTA

TABLE 1. 100 AREA LIQUID WASTE SITES

<u>Area No. and Name</u>	<u>Waste Types</u>	<u>Years</u>	<u>Estimated Volume</u>	<u>No. of Cubic Yds</u>	<u>References</u>
100 Area Decontamination Waste Cribs (116-B-4, 116-B-6, 116-C-2, 116-D-1A, 116-D-1B, 116-F-5, 116-F-10, 116-H-3, 116-K-2)	Decontamination solutions consisting of chromic, oxalic, and/or sulfuric acid solutions neutralized with soda ash. Solutions were used to decontaminate dummy fuel elements and reactor hardware.	1947-1970	2.3×10^6 L	3,000	Reference 6, pages 13-14, 17- 20, 45-48, 59- 60, 61-62, 113- 114, 121-122, 151-152, 173-174
100 Area Contaminated Reactor Effluent Cribs and Trenches (116-B-1, 116-B-3, 116-C-1, 116-D-2, 116-DR-1, 116-DR-2, 116-DR-4, 116-F-1, 116-F-2, 116-F-3, 116-F-4, 116-H-1, 116-H-4, 116-K-1, 116-K-2)	Reactor effluent con- taminated when fuel element cladding failed. Radiological contamination consisted chiefly of mixed fission products. Chemical contamination consisted of sodium dichromate added to reactor coolant as oxidant.	1946-1971	3.0×10^{11} L	3.9×10^8	Reference 6, pages 7-8, 11- 12, 43-44, 63- 64, 87-88, 89- 90, 93-94, 105- 106, 107-108, 109-110, 111- 112, 147-148, 153-154, 171- 172, 173-174

11/11/88
KFB

<u>Area No. and Name</u>	<u>Waste Types</u>	<u>Years</u>	<u>Estimated Volume</u>	<u>No. of Cubic Yds</u>	<u>References</u>
100 Area Laboratory Waste Cribs and Trenches (116-B-5, 116-B-10, 116-C-2, 116-D-3, 116-D-4, 116-F-9, 116-F-13)	Wastes associated with labs and shops in the 108 Buildings, including 108-B Tritium Lab and Tube Examination Facility, 108-C Metal Examination Facility and 100 F Biology Labs. Contaminants consisted mainly of tritium and low levels of mixed fission and activation products. Negligible chemical contamination except Biology Lab facilities, which received nitrate from animal wastes.	1950-1976	3.2 x 10 ⁸ L	4.2 x 10 ⁵	Reference 6, pages 15-16, 23-24, 45-48, 65-66, 67-68, 119-120, 127-128
100 Area Fuel Storage Basin Cribs and Trenches (116-B-2, 116-D-1A, 116-D-1B, 116-D-3, 116-F-3, 116-KE-3, 116-KW-2)	Storage basin cooling water contaminated with low levels of mixed fission products due to failure of fuel element cladding.	1946-1971	2.0 x 10 ⁷ l.	2.6 x 10 ⁴	Reference 6, pages 9-10, 59-60, 61-62, 91-92, 109-110, 179-180, 195-196

11/11/88
KFA

<u>Area No. and Name</u>	<u>Waste Types</u>	<u>Years</u>	<u>Estimated Volume</u>	<u>No. of Cubic Yds</u>	<u>References</u>
100 Area Reactor Decontamination and Maintenance Effluents Cribs and Trenches (116-DR-6, 116-F-1, 116-F-6, 116-F-12, 116-H-2, 116-K-2)	Reactor effluents during reactor maintenance activities. Much of the waste volume was uncontaminated. Radionuclides present were mainly low levels of mixed fission products and activation products. The main chemical contaminant was sulfuric acid, which was used for reactor decontamination. Sodium dichromate was also present in some of the effluents.	1945-1971	6.1 x 10 ⁸ L	8.0 x 10 ⁵	Reference 6, pages 95-96, 105-106, 115-116, 125-126, 149-150, 173-174
100 Area Reactor Confinement Seal Pit Drainage Cribs (116-DR-8, 116-F-7, 117-B, 117-C, 117-D, 117-H)	Drainage from 117 Building confinement system seal pits. Very low levels of contamination.	1960-1969	2.2 x 10 ⁶ L	2,900	Reference 6, pages 42-44, 49-50, 85-86, 99-100, 117-118, 167-168
100 Area Sulfuric Acid Sludge Disposal Sites (100KE*1, 100KE*2, 100KE*3, 100KW*1, 100KW*2)	Mercury-contaminated sludge from sulfuric acid storage tanks.	1955-1971	3.7 x 10 ⁴ L	48	Reference 6, pages 181-183, 185-187, 189-191, 197-199, 201-203

11/1/88
KPA

<u>Area No. and Name</u>	<u>Waste Types</u>	<u>Years</u>	<u>Estimated Volume</u>	<u>No. of Cubic Yds</u>	<u>References</u>
100 Area Miscellaneous Floor and Sink Drain Cribs (116-B-9, 116-D-6, 116-F-11)	Drainage from floor and sink drains. Very low levels of radioactive contamination. No chemical contamination.	1952-1967	3.4×10^5 L	440	Reference 6, pages 21-22, 69-70, 123-124
100 Area Potassium Borate Crib (116-DR-7)	Liquid potassium borate solution. No radioactive contamination.	1953	4×10^3 L	5.2	Reference 6, pages 97-98
100 Area Gas Purification Condensate Cribs (116-KE-1, 116-KW-1)	Condensate from reactor gas purification system. Contaminated with tritium and carbon-14 and small amounts of fission products.	1955-1971	1.6×10^6 L	2.1×10^3	Reference 6, pages 175-176, 193-194
100 Area Ion Exchange Regenerant Crib (116-KE-2)	Wastes from regeneration of ion exchange columns. Contained sulfuric acid, caustic, and mixed fission products and activation products.	1955-1971	3.0×10^6 L	3.9×10^3	Reference 6, pages 177-178

11/11/88
KE-2

<u>Area No. and Name</u>	<u>Waste Types</u>	<u>Years</u>	<u>Estimated Volume</u>	<u>No. of Cubic Yds</u>	<u>References</u>
100 Area Retention Basin Leakage (107-B, 107-C, 107-D, 107-DR, 107-F, 107-II, 107-KE, 107-KW)	Reactor effluent containing very small amounts of fission products and activation products. Chemical contamination limited to small amounts of sodium dichromate added to reactor coolant as oxidant.	1944-1971	3.0×10^{12} L	3.9×10^9	Reference 16
100 Area White Bluffs Pickling Acid Crib	Waste nitric and hydrofluoric acid used to pickle galvanized piping. Acid may or may not have been neutralized.	1943-1945	7×10^5 L	900	Reference 16
TOTAL			3.3×10^{12} L	4.3×10^9	

11/11/81
KTP

TABLE 2. 100 AREA SOLID WASTE SITES

<u>Area No. and Name</u>	<u>Waste Types</u>	<u>Years</u>	<u>Estimated Volume</u>	<u>No. of Cubic Yds</u>	<u>References</u>
100 Area Metallic Waste Burial Grounds (118-B-1, 118-B-4, 118-B-5, 118-C-2, 118-D-1, 118-D-3, 118-D-5, 118-DR-1, 118-F-1, 118-F-3, 118-F-7, 118-H-1, 118-H-2, 118-H-3, 118-H-4, 118-H-5, 118-K)	Metallic waste consisting mainly of discarded reactor hardware, including such items as aluminum tubes and spacers; lead and cadmium containing slugs, graphite, desiccant, aluminum and boron splines; lead; and mercury. Radioactive contaminants, primarily activation products.	1944-1975	1,000 m ³	1,300	Reference 6, pages 25-27, 33-34, 35-36, 55-56, 71-72, 77-79, 83-84, 101-102, 129-131, 135-136, 143-144, 155-157, 159-160, 161-162, 163-164, 165-166, 205-207
100 Area Construction Waste Burial Grounds (118-B-2, 118-B-3, 118-D-4, 118-F-1, 118-H-3)	Construction waste, demolition debris. Small amounts of radioactive contamination.	1953-1967	48,000 m ³	63,000	Reference 6, pages 29-30, 31-32, 81-82, 129-131, 161-162
100 Area Lab Waste Burial Grounds (118-B-6, 118-F-5, 118-F-6)	Low-level radioactive wastes from 100 Area laboratories, including tritium wastes and animal wastes.	1950-1975	20,000 m ³	26,000	Reference 6, pages 37-38, 139-140, 141-142

11/11/88
KTP

<u>Area No. and Name</u>	<u>Waste Types</u>	<u>Years</u>	<u>Estimated Volume</u>	<u>No. of Cubic Yds</u>	<u>References</u>
100 Area Miscellaneous Solid Waste Burial Grounds (118-B-1, 118-B-7, 118-C-1, 118-D-1, 118-D-2, 118-F-2, 118-F-4, 118-H-1, 118-K)	Radioactivity contaminated trash and miscellaneous solid wastes from 100 Area facilities.	1944-1975	61,000 m ³	80,000	Reference 6, pages 25-27, 39- 40, 51-53, 71- 72, 73-75, 133- 134, 137-138, 155-157, 205-207
100 Area Burn Pits (100 B/C Burn Pit, 100/DR Burn Pit, 100F Burn Pit, 100N Burn Pit, 100K Burn Pit)	Combustible trash from 100 Area buildings and offices.	1943-1971	130,000 m ³	170,000	Reference 6, pages 57-58, 103-104, 145- 146, 169-170, 209-210
		TOTAL	260,000 m ³	340,000 yd ³	

1/11/88
KLN

Is there tidal influence?

The site is located upstream of four dams, therefore, there is not tidal influence at the site.

Reference 5, page 2.1

Distance to a Sensitive Environment

Distance to 5-acre (minimum) coastal wetland, if 2 miles or less:

Not applicable.

Distance to 5-acre (minimum) fresh-water wetland, if 1 mile or less:

No fresh-water wetlands were found near the Hanford Site.

Reference 8

Distance to critical habitat of an endangered species or national wildlife refuge, if 1 mile or less:

Although there are several sensitive and threatened species that are residents of Hanford Site (for at least part of the year), no endangered species are known to be residents of the site. Two threatened species, the bald eagle and the ferruginous hawk, are residents of the site (for at least part of the year). Because there are no endangered species (federal listing) that reside at the site, there is no critical habitat to be considered in the ranking of the 300 Area Site.

The Draft Phase I Installation Assessment of Inactive Waste-Disposal Sites at Hanford, Volume 1, July 1986 listed the Merriam's Shrew as being on the State Endangered Species List. This information was taken from a list published in a preliminary draft of an environmental impact statement, and, since the referenced February 1987 list shows the Merriam's Shrew as only a proposed sensitive species, it is assumed that the previous list taken from the preliminary draft is in error.

Reference 4; Reference 14

Population Served by Surface Water

Location(s) of water-supply intake(s) within 3 miles (free-flowing bodies) or 1 mile (static water bodies) downstream of the hazardous substance and population served by each intake:

1/11/88
X12

9111000994

The 100-B Pumphouse (at 100-B/C Area) supplies water to 100-B/C, 100-D, 100-N, 100-K and 200 areas. The 100-D Pumphouse (at 100 D/DR Area) is a backup to the 100-B Pumphouse. The water plant at 100-N supplies water to both 100-N and the Washington Public Power Supply System Generating Station which is attached to the 100-N reactor. The 100-K Pumphouse supplies water to the 100-K area. The worker population of the 100 Area is 760 and the population of the 200 Area is 2,355.

Reference 9, pages II.1-64-II.1-65; Reference 13, page 3; Reference 15, page 1; Reference 4; Reference 8

Computation of land area irrigated by above-cited intake(s) and conversion to population (1.5 people per acre):

There are no crops irrigated by the 100 Area intakes, and there are no documented irrigation intakes drawing from the Columbia river within 3 miles of the 100 Area Site.

Total population served:

The total population served is calculated by summing all the respective population estimates:

$$760 + 2,355 = 3115$$

Name/description of nearest of the above water intakes:

The intake at the 100-D Area is the 181-D Pumphouse. The intake at the 100-N Area is the 181-N Pumphouse.

Reference 4; Reference 8

Distance to above-cited intakes, measured in stream miles.

The distance to the above intakes is zero because the intakes have documented observed releases.

Reference 4; Reference 8; Reference 11; Reference 12

1/11/88
KTP

AIR ROUTE

1 OBSERVED RELEASE

Contaminants detected:

Even though air concentrations of some of the constituents of interest can be detected above background offsite, no air monitoring data were found sufficient for HRS scoring of the Hanford CERCLA sites. These constituents of interest detected above background offsite are present in the routine gaseous effluents from operating facilities at Hanford. Therefore, the air route rating factors were not scored.

Date and location of detection of contaminants:

Not Applicable

Methods used to detect the contaminants:

Not Applicable

Rationale for attributing the contaminants to the site:

Not Applicable

2 WASTE CHARACTERISTICS

Reactivity and Incompatibility

Most reactive compound:

Not Applicable

Most incompatible pair of compounds:

Not Applicable

1/11/88
KWA

911899999

Toxicity

Most toxic compound:

Not Applicable

Hazardous Waste Quantity

Total quantity of hazardous waste:

Not Applicable

Basis of estimating and/or computing waste quantity:

Not Applicable

3 TARGETS

Population Within 4-Mile Radius

Circle radius used, give population, and indicate how determined:

0 to 4 mi 0 to 1 mi 0 to 1/2 mi 0 to 1/4 mi

Not applicable

Distance to a Sensitive Environment

Distance to 5-acre (minimum) coastal wetland, if 2 miles or less:

Not applicable

Distance to 5-acre (minimum) fresh-water wetland, if 1 mile or less:

Not Applicable

1/11/89
KCA

91118900997

Distance to critical habitat of an endangered species, if 1 mile or less:

Not Applicable

Land Use

Distance to commercial/industrial area, if 1 mile or less:

Not Applicable

Distance to national or state park, forest, or wildlife reserve, if 2 miles or less:

Not Applicable

Distance to residential area, if 2 miles or less:

Not Applicable

Distance to agricultural land in production within past 5 years, if 1 mile or less:

Not Applicable

Distance to prime agricultural land in production within past 5 years, if 2 miles or less:

Not Applicable

Is a historic or landmark site (National Register or Historic Places and National Natural Landmarks) within the view of the site?

Not Applicable

11/1/88
KCR

91119970999

HRS DOCUMENTATION LOG SHEET		SITE NAME <u>U.S. DOE Hanford - 100 Area</u>	
		CITY <u>Benton County</u> STATE <u>WA</u>	
		IDENTIFICATION NUMBER _____	
REFERENCE NUMBER	DESCRIPTION OF THE REFERENCE		
1	<u>Strontium Determination ALL Matrices, UST-RD-PM-9-80</u>		
2	Computer Printouts from Hanford Ground Water Data Base		
3	<u>Hanford Wells, PNL-5397, February 1985</u>		
4	<u>Draft Phase I Installation Assessment of Inactive Waste-</u>		
	<u>Disposal Sites at Hanford, Volume 1, July 1986</u>		
5	<u>Environmental Monitoring at Hanford for 1986, PNL-6120,</u>		
	May 1987		
6	<u>Draft Phase I Installation Assessment of Inactive Waste-</u>		
	<u>Disposal Site at Hanford, Volume 2, July 1986</u>		
7	<u>Uncontrolled Hazardous Waste Site Ranking System; A Users</u>		
	<u>Manual, 40 CFR 300, Appendix A</u>		
8	U.S.G.S. Maps of the Area Around 100 Area		
9	<u>Waste Management Operations, Volume 1, ERDA-1538</u>		
10	<u>Inductively Coupled Plasma Method, Method 6010</u>		
11	Hanford Stream data base printout showing Priest Rapids Dam		
12	<u>Investigation of Ground-Water Seepage from Hanford Shoreline</u>		
	<u>of the Columbia River, PNL-5289, November 1984</u>		
13	<u>Hanford Reservation Area Worker Census, BNWL-2298, July 1977</u>		
14	Endangered and Threatened Wildlife and Plants, 50 CFR,		
	Part 17, Subpart B, October 86		

11/11/88
X17

9118900999

REFERENCE 1

Strontium Determination ALL Matrices, UST-RD-PM-9-80

91118901001

STRONTIUM DETERMINATION
ALL MATRICES

Principle

Strontium is precipitated sequentially first as the nitrate, and then as the carbonate. The fuming nitric acid separation removes most of the other interfering ions and concentrated nitric precipitations remove calcium. Radioisotopes of barium, radium, and lead are co-precipitated with barium chromate. Iron and final traces of fission products are removed by a hydroxide scavenge using mixed hold back carrier. An additional barium and hydroxide scavenge is done on difficult matrices. Following a final carbonate precipitation, gravimetric yield and Strontium-89 plus Strontium-90 activity is determined or just Sr-85 tracer yield is established. Yttrium-90 is permitted to grow into equilibrium with the strontium-90, then separated from the strontium by successive hydroxide and oxalate precipitations. The yttrium oxalate is transferred to a counting dish, dried under a heat lamp, ignited to Yttrium Oxide, weighed for chemical recovery, and counted on a low background Beta Proportional Counter. Decay counts are made to check the purity of the Yttrium-90.

Scope: Procedures are described for the following matrices: milk, water, urine, feces, filter, soil, vegetation, and ion exchange resin. Food, bone, tissue and grains are treated as vegetation.

The different sample matrices analyzed as described in the scope require slightly different preparations. Refer to the Appropriate Preparation Procedure; 30-19-01, for Bioassay Samples and 30-20-01 for Environmental Matrices.

For Strontium analysis, varying sample types require even preliminary separations to bring them to a common point for the Strontium Determination steps summarized below.

Milk

Strontium is separated from milk and concentrated by equilibration with cation exchange resin using a batch technique. The resin after equilibration is washed with water to remove milk solids. Strontium is desorbed from the resin by equilibrating with 4M NaCl solution. Strontium is separated from the NaCl solution by precipitating as the carbonate.

Soil

Following leaching the sample is dissolved in 9M HCl and the majority of iron is removed using an isopropyl ether extraction. The HCl solution is converted back to the nitric form and a fuming nitric acid precipitation is performed. The nitric acid supernate is reserved for actinide (usually plutonium) chemistry.

Vegetation

In the case of vegetation (especially alfalfa) and produce, the sample residue may contain some radioactivity, depending on the sulfate content of the sample. Hence, following sample preparation and a fuming nitric acid precipitation, the residue is dissolved in water and any insoluble material is boiled with a concentrated solution of Na_2CO_3 to convert the insoluble sulfate residue to the acid soluble carbonate (metathesis). The carbonate is then refluxed with nitric acid and combined with the soluble water fraction.

Water

For large volume sample, the sample is acidified and concentrated by evaporation, neutralized with ammonium hydroxide, and strontium then is precipitated as the carbonate. Strontium is precipitated directly from small volume samples, one liter or less, by neutralizing with ammonium hydroxide to the red end point of phenolphthalein and then adding about 25 mL saturated Na_2CO_3 .

Urine and Feces

Following ashing, the sample salts are dissolved in nitric acid (aliquot of a fecal sample is analyzed), then made basic with hydroxide and carbonate heated and precipitated as the carbonate.

Ion Exchange Resin Samples

The resin is transferred to a beaker and dried and muffled for up to 1 week or more. Sodium carbonate (Na_2CO_3) is added and a fusion is performed. The fusion "cake" is dissolved in water and a small amount of hydrochloric acid and then reprecipitated as the carbonate.

Sequential Analyses

All samples requiring other analysis (usually plutonium analysis) in addition to strontium must first have a fuming nitric separation is performed on the sample residue instead of a carbonate.

Literature References

Sunderman, D. N. and Townley, C. W., The Radiochemistry of Barium, Calcium, and Strontium. Subcommittee on Radiochemistry, National Academy of Science - National Research Council, Office of Technical Services, Department of Commerce, Washington, D.C.

Baratta, E. G. and Knowles, F. E., Jr., Association of Official Analytical Chemists, 56, No. 1 (1973) 213-218.

Krieger, H. L. and E. L. Whittaker, Prescribed Procedures for Measurement of Radioactivity in Drinking Water, EPA-600/4-80-032 August 1980.

EML - Procedure Manual, HASL 300 Edited by Herbert L. Volchok and Gail de Planque, 26th edition, 1983.

Urine and Feces Sample Preparation Procedure, 30-BA-SP.

Sample Preparation Procedure For Environmental Matrices, 30-ENV-SP.

Stevenson, P. C. and Nervik, W. E., The Radiochemistry of the Rare Earths: Scandium, Yttrium, and Actinium. Subcommittee on Radiochemistry, National Academy of Science - National Research Council, Office of Technical Services, Department of Commerce, Washington, D.C.

Harley, J. P., Editor. Manual of Standard Procedures. Health and Safety Laboratory, United States Atomic Energy Commission, New York, 1976.

Reagents: All chemicals are of "reagent grade" and deionized water is used throughout this procedure.

Sr carrier: 39.4 mg Sr/mL, 143.5 g $\text{Sr}(\text{NO}_3)_2$ dissolved and diluted to 1 liter with 2M HNO_3 . Calibrated to 100.0 mg SrCO_3 per mL.

91118921004

Y carrier: 20 mg Y per mL, 25.4 g Y_2O_3 dissolved in 125 mL conc. HNO_3 and diluted to 1 liter with D.I. H_2O . Calibrated to 25.4 mg Y_2O_3 per mL.

Ba carrier: 10 mg Ba^{+2} /mL, 20 g $Ba(NO_3)_2$ dissolved and diluted to 1 liter with D.I. water.

HNO_3 : (nitric acid).

Fuming; 90% (or 24M HNO_3)

Concentrated; 16M HNO_3 .

2M HNO_3 ; 125 mL of concentrated HNO_3 diluted to 1 liter with D.I. water.

HCl: (hydrochloric acid) concentrated or 12M HCl.

9M HCl; 750 mL of concentrated HCl diluted to 1 liter with D.I. water.

6M HCl; 500 mL of concentrated HCl diluted to 1 liter with D.I. water.

CH_3COOH : (glacial acetic), 17.4M.

1.5M acetic acid, 86 mL glacial acetic acid diluted to 1 liter with D.I. water.

$NH_4C_2H_3O_2$: (ammonium acetate) 3M.

231 g of $NH_4C_2H_3O_2$ dissolved and diluted to 1 liter with D.I. water.

Na_2CrO_4 : (sodium chromate) 1.5M.

125 g Na_2CrO_4 dissolved and diluted to 1 liter with D.I. water.

NH_4OH : (ammonium hydroxide) 14.7M or concentrated.

Na_2CO_3 : (sodium carbonate) saturated aqueous - 225 g Na_2CO_3 diluted to 1 liter with D.I. water.

NaOH: (sodium hydroxide) 12M or concentrated. 540 g of sodium hydroxide dissolved and diluted to 1 liter with D.I. water.

6M NaOH; 270 g of sodium hydroxide diluted to 1 liter with D.I. water.

$H_2C_2O_4 \cdot 2H_2O$: (oxalic acid) 15% solution, 150 g of $H_2C_2O_4$ dissolved and diluted to 1 liter with D.I. water.

Collodion: 10% solution, (50 mL of collodion, 300 mL of diethyl ether, 150 mL of ethyl alcohol)

Methyl red indicator, 0.1%: Dissolve 0.1 g methyl red in 100 mL D.I. water.

Phenylphthalein indicator, 0.1%: 1 g phenylphthalein dissolved in 700 mL of 95% ethanol and 300 mL D.I. water.

Mixed Co, Ce, Cs, Mn, Ni, Fe Hold Back Carrier:

This solution when prepared as below will contain 2 mg/mL Co and Ni, 1.0 mg/mL Cs, Mn, Fe, and Ce. Transfer the following into a 250 mL beaker:

9.91 g $\text{Ni}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ (Nickel nitrate), and
7.24 g $\text{Fe}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$ (Ferric nitrate), and
3.60 g $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ (Manganese chloride), and
8.07 g $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ (Cobalt chloride), and
2.34 g $\text{Ce}(\text{NO}_3)_3 \cdot 6\text{H}_2\text{O}$ (Cerous nitrate)
1.27 g CsCl (Cesium chloride)

Add fifty mL of water and 5 mL of 8 M nitric acid. Warm gently until everything has dissolved, then transfer quantitatively to a 1 liter volumetric flask. Bring the volume to 1000.0 mL mark, mix well, then pour into a storage bottle without rinsing the volumetric flask.

Reagents for milk only:

Cation Exchange Resin - Dowex 50W-X8 (Na form), 50-100 mesh size, available from Bio-Rad Laboratories. (The commercially available H^+ form is converted into the Na^+ form by mixing 1.5l of 4M NaCl and 200 mL of resin on a magnetic stirrer for 30 minutes. The resin is filtered in a large Buchner funnel and washed with deionized water until the wash water is Cl-free when tested with 10% AgNO_3 . If a white AgCl precipitated forms, additional water washings are necessary. It is advantageous to prepare enough resin for several samples ahead of time.)

4M NaCl: 234 g NaCl dissolved in a 1 liter graduated mixing cylinder in distilled water, then diluted to 1 liter.

10% AgNO_3 Silver Nitrate: 50 grams of AgNO_3 dissolved in 500 mL of deionized water. Store in dark bottle, away from light.

Equipment and Materials

Beakers, 2 liter, 1 liter, 250 mL
Centrifuge
Centrifuge tubes, 100 mL plastic and/or glass, 50 mL conical bottom
Hot plate
Heat lamp
Meeker burner

Magnetic stirrer and stirring bar
Filter paper 7 cm GFA, Whatman #42 12.5 cm
Buchner funnel, 7 cm
Funnel, 60°
Side arm flask, 1000 mL
125 mL separatory funnel and stopper
Wrist action shaker
Platinum crucible, 150 mL
Muffle furnace 500°C, 1000°C
Stainless steel planchet 1 1/2", 1" and cardboard holders
Plastic jar 250 mL or plastic vial, 25 mL
Balance, 100 g

Calculation of Strontium Activities in Urine, Feces, and Environmental Matrices

The calculations for manual mode or computer mode (CALCTL for ERA samples, PERCTL for BA samples) are basically the same except that in the manual mode, results obtained should be considered as tentative, as only the uncertainty from counting statistics are considered.

If the program is not available, use the following calculation:

Sr-89 or Sr-90 decay corrected value =
(Sr-89 or Sr-90 reference value) x e^{-K t}

where:

t = difference in reference date and current date

"K" = decay constant for Sr-89 and Sr-90, expressed in the same time units as "t" = .693/t_{1/2}.

- Total strontium activity of the sample and the corresponding counting uncertainty are calculated as follows:

Sr (Total) dpm per sample or pCi per unit =

$$\frac{(C_1 - B)((dpc)_{Tot}) \times [V \text{ or } W]}{[2.22] \times Y_s \times (V_A \text{ or } W_A)}$$

Counting Uncertainty dpm or [pCi] per unit or [sample] =

$$\frac{[2] \times \text{SQR} (C_1/T_1 + B/T_B) \times (\text{dpc})_{\text{Tot}} \times [V \text{ or } W]}{[2.22] \times Y_S \times (V_A \text{ or } W_A)}$$

where:

[] = The square brackets indicate parameters exclusively used for environmental or bioassay analyses only.

(dpc)_{Tot} = Disintegrations per count value corresponding to the weight of Sr(NO₃)₂ in mg from the dpc table. Sr-89 and Sr-90 activities are assumed to contribute equally to the total activity.

C₁ = Gross cpm from the first strontium count

T₁ = Sample counting time in minutes

B = Background cpm

T_B = Background counting time in minutes

Y_S = Chemical recovery of strontium expressed as a decimal

[V. or W] = Volume or weight of the sample in the same units as V_A or W_A (applicable for bioassay samples only)

V_A or W_A = Volume or weight of the aliquot in the required unit

2.22 = Factor for conversion of activity from dpm to pCi (only for environmental samples)

[2] = Two sigma uncertainty reported for environmental samples only

2. Sr-90 activity of the sample and the corresponding counting uncertainty from the Y-90 milking is calculated as follows:

Sr-90 dpm per sample or pCi per unit =

$$\frac{(C_Y - B) \times e^{K_2 t_2} \times (\text{dpc})_{Y_2} \times [V \text{ or } W]}{[2.22] \times I \times Y_S \times Y_Y \times V_A \text{ or } W_A}$$

Counting Uncertainty dpm per sample or pCi per unit =

$$\frac{[2] \times \text{SQR} (C_Y/T_Y + B/T_B) \times e^{K_2 t_2} \times (\text{dpc})_{Y_2} \times [V \text{ or } W]}{[2.22] \times I \times Y_s \times Y_y \times V_A \text{ or } W_A}$$

where:

[] = parameters exclusively used for environmental or bioassay analysis

C_Y = Gross Y-90 cpm

T_Y = Time of Y-90 counting in minutes

B = Background cpm

T_B = Length of background counting in minutes

$e^{K_2 t_2}$ = Correction for the decay of Y-90 during the time (t_2 hours) elapsed between Y-90 milking and counting. "K" is the decay constant for Y-90 = 0.693/64.1 hours.

$(\text{dpc})_{Y_2}$ = Disintegrations per count of Y-90 corresponding to the weight of yttrium oxide and the counter used

[V or W] = Volume or weight of the total sample in the same units as V_A or W_A for Bioassay only

V_A or W_A = Volume or weight of the sample aliquot in the units required

[2.22] = Factor, dpm or pCi (Applicable for environmental samples only)

I = Correction for the fractional ingrowth of Y-90 in the strontium sample at the time of milking (from the table)

Y_s = Gravimetric recovery of strontium as strontium carbonate

Y_y = Gravimetric recovery of yttrium oxide (weight of final Y_2O_3 precipitated (mg)/25.4 mg expected).

3. Sr-89 Determination: Sr-89 activity is determined by subtracting Sr-90 activity and Y-90 ingrowth, which are estimated from Y-90 milking and determination, from the total strontium activity observed during the first strontium count with appropriate corrections for self absorption, scattering, and counting efficiency.

Sr-89 dpm per sample or pCi per unit =

$$(C_1 - B) - \frac{(dpm)_{S90}}{(dpc)_{S90}} - \frac{(dpm)_{S90}}{(dpc)_{Y1}} * (1 - e^{-K_1 t_1})$$

$$\frac{x (dpc)_{S89} * e^{K_3 t_3} * V \text{ or } W}{[2.22] * Y_S * V_A \text{ or } W_A}$$

where:

- [] = Parameters exclusive for environmental or bioassay analyses
- $(dpm)_{S90}$ = Disintegrations per minute of Sr-90 nominal value before correcting for factors, such as 2.22, Y_S , volume and weight terms as shown below:

$$\frac{(C_y - B) * e^{K_2 t_2} * (dpc)_{Y_2}}{I * Y_y}$$
- $(dpc)_{S90}, (dpc)_{Y1}, (dpc)_{S89}$ = Disintegration per count of Sr-90, Y-90, and Sr-89 respectively corresponding to the counter used and the weight of strontium carbonate.
- $(1 - e^{-K_1 t_1})$ = Factor expressing Y-90 ingrowth in the strontium sample during the time (t_1 in hours) elapsed between the time " t_0 " of the first Y-90 removal from the strontium sample and the first (strontium) count. " K_1 " is the decay constant for Sr-90 = .693/28.8 years.
- 2.22 = Factor, dpm per pCi - for environmental samples only.
- [V or W] = Volume or weight of the sample in the same units as V_A or W_A - only for Bioassay analysis.

$e^{K_3 t_3}$ = Correction for the decay of Sr-89 from the date of first strontium counting to the sample date. " K_3 " is the decay constant for Sr-89 = 0.693/30.5 days.

Counting Uncertainty dpm per sample or pCi per unit =

$$[2] \times \text{SQR} (C_1/T_1 + B/T_B) + \frac{E_{S90}^2}{(\text{dpc})_{S90}^2} + \frac{E_{S90}^2(1 - e^{-K_1 t_1})^2}{(\text{dpc})_{Y_1}^2}$$

$$\frac{x (\text{dpc})_{Sr89} \times e^{K_3 t_3} \times [V \text{ or } W]}{[2.22] \times Y_3 \times V_A \text{ or } W_A}$$

where:

$$E_{S90} = \frac{\text{SQR}(C_Y/T_Y + B/T_B) \times e^{K_2 t_2} \times (\text{dpc})_{Y_2}}{I \times Y_Y}$$

All the known uncertainties are propagated using a computer program.

PRE-SEPARATION:

Milk

- 1(M). Add 100 cc of Dowex 50W-X8 resin (Na-form) equilibrated as described in the reagent section, to the milk sample aliquots and stir for 3 hours using a magnetic stirrer. Allow the resin to settle for 5-10 minutes. Decant as much of the milk as possible into a separate beaker without losing any resin.
- 2(M). Add about 500 mL of D.I. Water to the resin beaker and stir for about one minute using magnetic stirrer. Allow the resin to settle, decant and discard the supernate. Repeat this process until visible milk solids (cloudiness) have been removed from the resin.
- 3(M). Add 400 mL of 4M NaCl to the resin sample and stir for 15 minutes. Allow the resin to settle and filter through a 7-Cm GFA filter and collect the filtrate in a 1000 mL Erlenmeyer flask. Transfer the NaCl solution from the flask to a clean 1000 mL beaker. Transfer the resin from the filter reservoir back to the resin beaker, add 200 mL of 4M NaCl, stir for 15

min., filter under is before vacuum and rinse the resin with H₂O, collect the filtrate in the 1000 mL Erlynmeyer flask. Add the second NaCl solution and rinse from the flask to the 1000 mL beaker containing the first NaCl solution as wash solution. Discard the resin to waste.

4(M). Add 2-3 drops of phenolphthalein indicator to the NaCl solution, then add conc NH₄OH while stirring until the solution turns pink (end point). Add 70-100 mL of saturated Na₂CO₃, and heat for 10 minutes, then cool to room temperature. Decant most of the supernate to waste. Centrifuge to collect the precipitate remaining.

5(M). Proceed to the Strontium Determination Section.

KEYPOINT: Exercise extreme caution when performing the fuming nitric acid step since a vigorous reaction is possible if any resin is remaining.

Soil

1(S). Evaporate the 8M HNO₃ sample leachate to dryness, then add about 5 mL conc. HCl and carefully evaporate to near dryness at least two times.

2(S). Dissolve the sample in 9M HCl, transfer to a separatory funnel, add about 20 mL of isopropyl ether and shake with frequent venting for 1-2 minutes. Reject the ether (upper phase). Repeat the isopropyl ether extraction two more times or until the yellowish tint indicating iron does not appear in the ether phase.

3(S). Transfer 9M HCl phase back to the original beaker and evaporate to dryness. Convert the sample back to the nitric media using 2-10 mL additions of conc. nitric acid and cooperating to dryness after each addition.

4(S). Proceed to step one of the Strontium Determination Section.

Water

1(W). Add saturated Na₂CO₃ directly to the water sample or to the water sample that has been concentrated by evaporating and allow the precipitate to settle overnight. Decant most of the supernate to waste and collect the precipitate in a centrifuge tube. Dissolve the precipitate in about 5 mL of 2M HNO₃, dilute to 20 mL with D.I. H₂O and proceed to step 4

of the Strontium Determination Section.

Urine and Feces

- 1(U). Dissolve sample salts by adding about 50 mL of D.I. water, warm and add 5 to 10 mL of conc. HNO_3 . Add phenolphthalein indicator, and adjust to the red end point using concentrated NH_4OH . Add 25 mL of saturated Na_2CO_3 , heat for 2-5 min. (metathesis), then cool for 10 minutes.
- 2(U). Transfer, using D.I. water to 1 or 2 100 mL plastic centrifuge tubes and allow the sample to cool for at least 1/2 hour. Centrifuge and decant the supernate to waste. Add about 30-40 mL of D.I. Water. Stir, centrifuge, and decant the supernate to waste.
- 3(U). Proceed to the Strontium Determination Section.

Ion Exchange Resin Samples

- 1(R). A prepared amount (see 30-20-01) of strontium-85 and strontium carrier is added to the sample in its original container. Add a small amount of deionized water, cap the container and shake vigorously. Transfer the sample quantitatively to a 600 mL beaker (labeled with a Tech pen) with DI water followed by a rinse with about 5 mL of concentrated nitric acid. The entire sample is again mixed with a stirrer and dried under a heat lamp.
- 2(R). The sample is ashed to a pinkish white residue at 500-550°C in a muffle furnace (process may take up to a week or more) and then transferred quantitatively to a platinum crucible. Na_2CO_3 is added in an amount equal to 4-5 times that of the sample weight and mixed in thoroughly. The mixture is fused at 900-1000°C for approximately 1-2 hours.
- 3(R). The platinum crucible is placed in a 600 mL beaker and the fusion mixture is dissolved with approximately 200 mL of hot water. The platinum crucible is removed and rinsed with a minimum of 6M HCl .
- 4(R). Approximately 25 mL of sat. Na_2CO_3 is added to the sample with stirring. After allowing the precipitate to settle the majority of the supernate is decanted to waste and the remaining is centrifuged. Proceed with the strontium carbonate precipitate to the Strontium Determination Section.

STRONTIUM DETERMINATION

SPECIAL SAFETY PRECAUTION: Fuming nitric acid is used in this procedure. It can be hazardous; especially when brought in contact with organics, therefore exercise extreme care.

All waste fractions are generally saved to the end of the procedure; after acceptable recoveries are received they may be discarded.

1. Dissolve the sample residue or carbonate precipitate in a minimum amount of HNO_3 while stirring. Add 50-70 mL of fuming nitric acid and cool in an ice bath for 20-30 minutes. Centrifuge and save the supernate for Sequential Analysis if requested; proceed to the plutonium procedure (20-Pu-01) with this fraction.

The fuming nitric acid addition may be done in a beaker first, then transferred to centrifuge tubes, especially in the case of soil and vegetation.

KEY POINT: The fuming nitric precipitation is an excellent separation point for sequential chemistry and is usually included in the sample preparation chemistry (30-ENV-SP).

2. To the precipitate, add 30-40 mL of conc. HNO_3 to remove excess calcium. Stir, centrifuge and decant to waste. Repeat one or more times depending on the calcium content of the particular matrix.
3. Add 20-30 mL of warm D.I. water. Stir to dissolve completely. If any insoluble material is present separate this precipitate by centrifuging and decanting the supernate to a clean centrifuge tube.

For Vegetation Samples (or any other sample that has been muffled) when insoluble residue remains after dry washing: (metathesis) Transfer the insoluble material to a beaker with 2M HNO_3 , add excess Na_2CO_3 and boil for 15 minutes. Transfer back to the tube, centrifuge and decant supernate to waste. Dissolve precipitate in 5-15 mL conc. HNO_3 , warm, stir, centrifuge and combine this supernate with the water soluble fraction.

KEY POINT: Insoluble material not undergoing metathesis should be retained until an acceptable final precipitate weight

recovery or tracer recovery is established, then it can be discarded.

4. Add 1-2 drops of phenolphthalein indicator and adjust dropwise just to Basic (pink endpoint) using conc. NH_4OH , then add 5 drops in excess. Add 1 or 2 drops of mixed (hold back) carrier, allow iron and other metal hydroxides to flocculate. Centrifuge and transfer the supernate to a clean centrifuge tube. Rinse the precipitate with 10 mL of D.I. water centrifuge and combine the supernates.
5. Add 10 mL of freshly prepared buffer solution prepared by mixing equal amounts of 1.5M acetic acid and 3M ammonium acetate. Add 2-3 drops of methyl red indicator and adjust the pH to about 5.0, straw color of methyl red, using glacial acetic acid or conc. NH_4OH . Heat for 5-10 min. then add 1 mL of barium carrier and 1 mL of 1.5M Na_2CrO_4 and continue heating for 5-10 min.
6. Centrifuge and/or filter the BaCrO_4 through a 2" 600 funnel (using Whatman #42 filter paper) into a clean centrifuge tube. Rinse the filter with D.I. water. Fill the centrifuge tube or tubes with saturated Na_2CO_3 , centrifuge, and decant the supernate to waste.

KEY POINT: Centrifuge only if the BaCrO_4 precipitate appears to be heavy, otherwise filter.

For vegetation, soil and filter samples. Dissolve the precipitate in a minimum of 2M HNO_3 and repeat steps 4, 5, and 6.

7. Add fuming HNO_3 dropwise to the precipitate until the effervescence has ceased, then add 50 mL of fuming HNO_3 and cool in the ice water bath for 5-10 minutes. Centrifuge and decant the supernate to waste.

KEY POINT: Add the fuming nitric very slowly at first to avoid sample loss due to the effervescence.

8. Wash the sample one or more times using 20-30 mL of conc. HNO_3 depending on sample matrix. High calcium content matrices may require up to 5 nitric acid washes. Record the Sr-Y separation time on the counting request card.

KEY POINT: 1. If an unusually large precipitate

does not appear to diminish in size after two conc HNO₃ washes discontinue further washes and check with your supervisor.

2. The time of the last nitric acid wash should be considered the separation time.
9. If Sr-85 tracer was added, dissolve the sample precipitate in D.I. H₂O and transfer to 25 mL volume in a styrene jar or 20 mL volume in a plastic (scintillation) vial and submit for a gamma count for Sr-85 recovery.
10. If only Sr (cold) carrier was added, dissolve the precipitate in 10 mL of H₂O, add 25-50 mL of saturated Na₂CO₃, warm for 5-10 minutes, centrifuge, and discard the supernate. Wash the precipitate twice with 20-30 mL D.I. H₂O, centrifuge and discard the wash. Slurry the strontium carbonate precipitate with 1-2 mL of D.I. H₂O and transfer quantitatively to a tared 1" or 1 1/2" stainless steel planchet. Dry under infrared lamp. Cool, weigh, and record on the request card and beta count on either a Low Beta (plated on 1 1/2" planchet) or a Quad Low Background Beta proportional counter (plated on 1" or 1 1/2" planchet).
11. Allow up to 14 days to permit Y-90 to grow into equilibrium with the Sr-90 (the ingrowth time may be shortened if necessary provided a growth correction is made).

STRONTIUM-90 (BY YTTRIUM-90)

12. Using 2M HNO₃, carefully dissolve and rinse the strontium carbonate precipitate from the planchet into a 50-mL centrifuge tube. If Sr-85 yield determination was performed, transfer from the Styrene jar or plastic and to a 100-mL tube. Transfer one Yttrium carrier vial containing exactly 1 mL of Yttrium carrier (20.0 mg Y/mL). Rinse the vial with D.I. water, stir and let stand for 10 minutes.
13. Add 12M NaOH until a permanent clear gelatinous precipitate forms, then add 2 to 3 mL excess and decant supernate to a Styrene jar for possible re-milk. Stir, centrifuge for 5 minutes. Note the Yttrium separation date and time on the analysis request card.
14. Dissolve the precipitate with (1-2 mL) of conc. HNO₃, dilute to approximately 25-30 mL with water, then add conc. NH₄OH (approximately 15 mL) until a permanent precipitate forms. Add 2 to 3 mL excess NH₄OH, stir, centrifuge for 5 minutes and decant the supernate to waste.
15. Repeat Step #14 then wash the precipitate with 25 mL of water, centrifuge for 5 minutes and decant the supernate to waste.
16. Dissolve the precipitate using 1 mL of conc. HCl, then add 40 mL of boiling hot 15% oxalic acid solution and let stand for 5 to 10 minutes. Centrifuge for 5 minutes and decant and discard the supernate. Wash the oxalate precipitate twice with 10 mL of water, centrifuge for 5 minutes and decant and discard the supernate.
17. Transfer the precipitate with water to a previously flamed, labeled, and tared 1" or 1 1/2" stainless steel planchet (depending on the beta counter to be used) with a plastic transfer pipet and dry under a heat lamp. Ignite slowly over a Meker burner to convert the oxalate to the oxide. Cool, weigh and record weight on the request card, fix the sample to the planchet with collodion (2 or 3 drops), and submit for a beta proportional count.

L A S T P A G E

APPROVED FOR USE

RG J... 2/6/86

Mathew M Hardy 2/6/86

Al Robinson 2/4/86

91119901010

REFERENCE 2

Computer Printouts from Hanford Ground Water Data Base

6
1
0
1
0
6
0
1
1
1
1
6

00000000
00000000
00000000

Earth Sciences VAX Computer Center

0000000000
0000000000
0000000000

DATE: 14-AUG-1987 10:41

H	H	EEEE	TTTT	PPPP	RRRR	III	H	H	TTTT		
H	H	E	T	P	P	R	R	I	H	H	T
HH	H	E	T	P	P	R	R	I	HH	H	T
H	H	EEEE	T	PPPP	RRRR	I	H	H	T		
I	HH	E	T	P	R	R	I	N	HH	T	
I	H	E	T	P	R	R	I	H	H	T	
I	H	EEEE	T	P	R	R	III	H	H	T	

11	000000	000000			
11	000000	000000			
1111	00	00	00	00	
1111	00	00	00	00	
11	00	0000	00	0000	
11	00	0000	00	0000	
11	00	00	00	00	00
11	00	00	00	00	00
11	0000	00	0000	00	
11	0000	00	0000	00	
11	00	00	00	00	
11	00	00	00	00	
111111	000000	000000			
111111	000000	000000			

TTTTTTTTTT	HH	HH	PPPPPPP	1111	77777777	
TTTTTTTTTT	HH	HH	PPPPPPP	1111	77777777	
TT	HH	HH	PP	PP	1111	77
TT	HH	HH	PP	PP	1111	77
TT	HH	HH	PP	PP	1111	77
TT	HH	HH	PP	PP	1111	77
TT	HH	HH	PPPPPPP	1111	77	
TT	HH	HH	PPPPPPP	1111	77	
TT	HH	HH	PP	1111	77	
TT	HH	HH	PP	1111	77	
TT	HH	HH	PP	11	77	
TT	HH	HH	PP	11	77	
TT	HH	HH	PP	11	77	
TT	HH	HH	PP	11	77	

File #BKA; (100.TMP)7 (4227,77,0), last revised on 14-AUG-1987 10:41, is a 32 block sequential file owned by UIC (NETPRINT). The records are variable length with implied (CR) carriage control. The longest record is 0 bytes.

Job 100 (148) queued to LPA0 on 14-AUG-1987 10:41 by user NETPRINT, UIC (NETPRINT), under account H8610010 at priority 100, started on printer *LPA0; on 14-AUG-1987 10:41 from queue LPA0.

00000000
00000000
00000000

Digital Equipment Corporation - VAX/VMS Version V4.5

0000000000
0000000000
0000000000

9 1 1 1 0 2 0 1 0 2 1

WELL NAME	COLLECTION DATE	CONSTITUENT NAME	LESS THAN FLAG	ANALYSIS VALUE	ANALYSIS SPONSOR
1-13-1	5/21/87	FCR0011		6.2000E+01	H D6 for Chromium
1-13-12	3/25/87	FCR1011		1.5300E+03	H D6 " "
1-13-21	5/18/87	FCR1011		1.3700E+02	H D6 " "

WELL NAME	COLLECTION DATE	CONSTITUENT NAME	LESS THAN FLAG	ANALYSIS VALUE	ANALYSIS SPONSOR
1-14-1	1/10/87	SR001		0.0500E+00	H D6 for 90Sr
1-14-1	5/29/87	SR001		1.2500E+01	H D6 for 90Sr

WELL NAME	COLLECTION DATE	CONSTITUENT NAME	LESS THAN FLAG	ANALYSIS VALUE	ANALYSIS SPONSOR
1-11-5	4/10/87	1,1,1-T	<	1.0000E+01	H
1-11-5	4/10/87	1,1,2-T	<	1.0000E+01	H
1-11-5	4/10/87	ALUMINUM	<	1.5000E+02	H
1-11-5	4/10/87	ANTIMONY	<	5.0000E+01	H
1-11-5	4/10/87	ARSENIC	<	1.0000E+02	H
1-11-5	4/10/87	BARBIT	<	5.0000E+00	H
1-11-5	4/10/87	BERYLLIUM	<	5.0000E+00	H
1-11-5	4/10/87	BETA	<	8.7100E+00	H
1-11-5	4/10/87	CADMIUM	<	2.0000E+00	H
1-11-5	4/10/87	CALCIUM	<	6.0300E+04	H
1-11-5	4/10/87	CILFORN	<	2.0000E+01	H
1-11-5	4/10/87	CHLORIDE	<	6.5300E+03	H
1-11-5	4/10/87	CHROMIUM	<	1.7700E+02	H
1-11-5	4/10/87	COBALT	<	-1.6000E+00	H
1-11-5	4/10/87	CILIFRIN	<	2.2000E+00	H
1-11-5	4/10/87	CONDUCT	<	4.1400E+02	H
1-11-5	4/10/87	COPPER	<	1.0000E+01	H
1-11-5	4/10/87	CS-137	<	3.6600E+00	H
1-11-5	4/10/87	FALMIN	<	1.5000E+02	H
1-11-5	4/10/87	FANTHIN	<	1.0000E+02	H
1-11-5	4/10/87	FARGE II	<	5.0000E+00	H
1-11-5	4/10/87	FRANCIUM	<	5.5000E+01	H
1-11-5	4/10/87	FIBRYLL	<	5.0000E+00	H
1-11-5	4/10/87	FCADMIUM	<	2.0000E+00	H
1-11-5	4/10/87	FCALCIUM	<	5.0000E+04	H
1-11-5	4/10/87	FCR0011	<	1.6000E+02	H
1-11-5	4/10/87	FCOPPER	<	1.0000E+01	H
1-11-5	4/10/87	FIRIN	<	5.0000E+01	H
1-11-5	4/10/87	FLEAD	<	5.0000E+00	H
1-11-5	4/10/87	FLUORIDE	<	5.0000E+02	H
1-11-5	4/10/87	F IAGNES	<	1.0000E+04	H
1-11-5	4/10/87	F IALGAN	<	5.0000E+00	H
1-11-5	4/10/87	F IERCHUR	<	1.0000E-01	H
1-11-5	4/10/87	F IICKEL	<	1.0000E+01	H
1-11-5	4/10/87	F IUSIUM	<	3.0000E+02	H
1-11-5	4/10/87	F IOTASS	<	4.6000E+03	H
1-11-5	4/10/87	F ISELENI	<	5.0000E+00	H
1-11-5	4/10/87	F IILVER	<	1.0000E+01	H
1-11-5	4/10/87	F IODINE	<	1.0000E+04	H
1-11-5	4/10/87	F ITRIUM	<	3.0000E+02	H

5
5
5

0 9 0 1 0 2 7

1-11-5	1/10/87	F/ALADI		5.0000E+00	H
1-11-5	1/10/87	FZINC		4.2000E+01	H
1-11-5	1/10/87	IRON		5.0000E+01	H
1-11-5	4/10/87	LEADCF	<	5.0000E+00	H
1-11-5	1/10/87	LIALP/IA		2.3600E+00	H
1-11-5	1/10/87	M-XYLE	<	1.0000E+01	H
1-11-5	1/10/87	MAGNES		1.0400E+04	H
1-11-5	1/10/87	MAGNESE	<	5.0000E+00	H
1-11-5	1/10/87	MERCURY	<	1.0000E+01	H
1-11-5	1/10/87	MET-IO IE	<	1.0000E+01	H
1-11-5	1/10/87	MET-IVCH	<	1.0000E+01	H
1-11-5	1/10/87	NICKEL	<	1.0000E+01	H
1-11-5	1/10/87	NITRATE		2.8700E+04	H
1-11-5	1/10/87	OPXYLE	<	1.0000E+01	H
1-11-5	1/10/87	OSMIUM	<	3.0000E+02	H
1-11-5	4/10/87	PERCE IE	<	1.0000E+01	H
1-11-5	1/10/87	PH		7.3000E+00	H
1-11-5	1/10/87	P/IOBP IA	<	1.0000E+03	H
1-11-5	1/10/87	POTASSIUM		4.5500E+03	H
1-11-5	1/10/87	RADIUM	<	3.4700E+02	H
1-11-5	1/10/87	R/IO	<	-1.9300E+01	B
1-11-5	1/10/87	SELENIUM	<	5.0000E+00	H
1-11-5	1/10/87	SILVER	<	1.0000E+01	H
1-11-5	1/10/87	SODIUM		1.1000E+04	H
1-11-5	1/10/87	SR	<	7.5300E+01	H
1-11-5	1/10/87	SULFUR	<	3.0000E+02	H
1-11-5	1/10/87	SULFATE	<	5.7000E+04	H
1-11-5	1/10/87	TETRA IE	<	1.0000E+01	H
1-11-5	1/10/87	TIC	<	0.6500E+02	H
1-11-5	1/10/87	TIX	<	1.0000E+02	H
1-11-5	4/10/87	TRICE IE	<	1.0000E+01	H
1-11-5	1/10/87	TRITIUM		1.0700E+03	B
1-11-5	1/10/87	U		2.0700E+00	B
1-11-5	1/10/87	VAIAD IUM		5.0000E+00	H
1-11-5	1/10/87	ZINC		4.8000E+01	H

BKG for 90Sr

WELL NAME	COLLECTION DATE	CONSTITUENT NAME	LESS FLAG	ANALYSIS VALUE	ANALYSIS SPONSOR
6-77-54	6/12/87	1,1,1-T	<	1.0000E+01	H
6-77-54	6/12/87	1,1,2-T	<	1.0000E+01	H
6-77-54	6/12/87	1,1-DIM	<	3.0000E+03	H
6-77-54	6/12/87	1,2-DIM	<	3.0000E+03	H
6-77-54	6/12/87	12-IBEN	<	1.0000E+01	H
6-77-54	6/12/87	1234TE	<	1.0000E+01	H
6-77-54	6/12/87	1235TE	<	1.0000E+01	H
6-77-54	6/12/87	123THI	<	1.0000E+01	H
6-77-54	6/12/87	13-IBEN	<	1.0000E+01	H
6-77-54	6/12/87	135TRI	<	1.0000E+01	H
6-77-54	6/12/87	14-IBEN	<	1.0000E+01	H
6-77-54	6/12/87	ACETYLE	<	3.0000E+03	H
6-77-54	6/12/87	ACRYLIE	<	3.0000E+03	H
6-77-54	6/12/87	ALKALIN		1.1000E+05	H
6-77-54	6/12/87	ALLYLAL	<	3.0000E+03	H
6-77-54	6/12/87	AMMONIU	<	5.0000E+01	H
6-77-54	6/12/87	BETA		0.5100E+00	H
6-77-54	6/12/87	CILACET	<	3.0000E+03	H
6-77-54	6/12/87	CILFUM	<	1.0000E+01	H
6-77-54	6/12/87	CILURAL	<	3.0000E+03	H

Ref. 2.3

6-77-54	6/12/87	CILTRID		5.7101E+03	H
6-77-54	6/12/87	CILPRIP	<	3.0700E+03	H
6-77-54	6/12/87	COMBCT		2.7700E+02	H
6-77-54	6/12/87	CYALID	<	3.0000E+03	H
6-77-54	6/12/87	CYALIL	<	3.0000E+03	H
6-77-54	6/12/87	CYALJG	<	3.0000E+03	H
6-77-54	6/12/87	DICPRIP	<	3.0000E+03	H
6-77-54	6/12/87	ETHCALD	<	3.0000E+03	H
6-77-54	6/12/87	ETHCYAN	<	3.0000E+03	H
6-77-54	6/12/87	ETHIXID	<	3.0000E+03	H
6-77-54	6/12/87	FALPHID	<	1.5300E+02	H
6-77-54	6/12/87	FANTID	<	1.0000E+02	H
6-77-54	6/12/87	FARSE II	<	5.0000E+00	H
6-77-54	6/12/87	FHAIID	<	2.4000E+01	H
6-77-54	6/12/87	FOERYLL	<	5.0000E+00	H
6-77-54	6/12/87	FCAPIID	<	2.0000E+00	H
6-77-54	6/12/87	FCALEID	<	3.5000E+00	H
6-77-54	6/12/87	FCAID II	<	1.0000E+01	H
6-77-54	6/12/87	FCOMPER	<	1.0000E+01	H
6-77-54	6/12/87	FIRID	<	5.0000E+01	H
6-77-54	6/12/87	FLURID	<	5.0000E+02	H
6-77-54	6/12/87	FLURIDA	<	3.0000E+03	H
6-77-54	6/12/87	FRAGNES		1.1600E+04	H
6-77-54	6/12/87	FRANGAN	<	5.0000E+00	H
6-77-54	6/12/87	FRENCIR	<	1.0000E+01	H
6-77-54	6/12/87	FRICKEL	<	1.0000E+01	H
6-77-54	6/12/87	FUSID II	<	3.0000E+02	H
6-77-54	6/12/87	FROTAS	<	4.0000E+03	H
6-77-54	6/12/87	FRLEII	<	5.0000E+00	H
6-77-54	6/12/87	FRILVER	<	1.0000E+01	H
6-77-54	6/12/87	FRUJIM	<	1.5000E+04	H
6-77-54	6/12/87	FRTRIT	<	3.0000E+02	H
6-77-54	6/12/87	FRVIAII	<	2.7000E+01	H
6-77-54	6/12/87	FZID	<	1.0000E+01	H
6-77-54	6/12/87	GLYCIDY	<	3.0000E+03	H
6-77-54	6/12/87	HEXACIL	<	1.0000E+01	H
6-77-54	6/12/87	HEXCHEN	<	1.0000E+01	H
6-77-54	6/12/87	HYDRAZI	<	3.0000E+03	H
6-77-54	6/12/87	ISOLUTY	<	3.0000E+03	H
6-77-54	6/12/87	KERCHEN	<	1.0000E+04	H
6-77-54	6/12/87	LALPIIA	<	1.0000E+00	H
6-77-54	6/12/87	LXYLE	<	1.0000E+01	H
6-77-54	6/12/87	NETIDIE	<	1.0000E+01	H
6-77-54	6/12/87	NETIYCI	<	1.0000E+01	H
6-77-54	6/12/87	NETZIE	<	3.0000E+03	H
6-77-54	6/12/87	NAPHTIA	<	1.0000E+01	H
6-77-54	6/12/87	NITRATE	<	7.9000E+03	H
6-77-54	6/12/87	OPXYLE	<	1.0000E+01	H
6-77-54	6/12/87	PARALIE	<	3.0000E+03	H
6-77-54	6/12/87	PENTCHJ	<	1.0000E+01	H
6-77-54	6/12/87	PERCEIE	<	1.0000E+01	H
6-77-54	6/12/87	PJ		7.3000E+00	H
6-77-54	6/12/87	PJ		7.6400E+00	H
6-77-54	6/12/87	PJEDL	<	1.0000E+01	H
6-77-54	6/12/87	PHOSP. IA	<	1.0000E+03	H
6-77-54	6/12/87	PHOPYLA	<	3.0000E+03	H
6-77-54	6/12/87	PHOPYLU	<	3.0000E+03	H
6-77-54	6/12/87	SULFATE	<	4.1500E+04	H
6-77-54	6/12/87	TETRAIE	<	1.0000E+01	H
6-77-54	6/12/87	TETRIIB	<	1.0000E+01	H
6-77-54	6/12/87	TIC	<	4.4500E+02	H
6-77-54	6/12/87	TIXIDL	<	1.6500E+00	H
6-77-54	6/12/87	TRECEIE	<	1.0000E+01	H

BKG for Chromium

REFERENCE 3

Hanford Wells, PNL-5397, February 1985

91118901024

WELL DESIGNATION EMA NO.	COORDINATES	CASING ELEV. (FT-ASL)	DRILL DEPTH (FT)	INIT. DEPTH (FT) TO WATER	DIA. (IN)	DEPTH TO BOTTOM (FT)	MIN-MAX PERFORATED DEPTH (FT)	DATE COMP. (M-Y)	FORMER DESIGNATION	COMMENTS
199 H3 1890	1 P N 094994 W 040052	421.90	75	45	8.0	75	29 - 74	8-60		SAMPLE PUMP
199 H4 1875	1 P N 095700 W 038400	417.75	75		6.0	30	25 - 50	3-52	107-H-1	CASING REMOVED
199 H4 1876	2 P N 095200 W 030565	420.43	306	3	6.0	311	25 - 50	5-52	107-H-2	CONFINED AQUIFER
199 H4 1877	3 P N 096373 W 039000	420.35	55	39	6.0	55	34 - 55	5-74		SAMPLE PUMP
199 H4 1878	4 P N 093560 W 038605	413.76	55	36	6.0	50		6-83		SCREEN 33-43 GROUTED SAMPLE PUMP
199 H4 1873	5 P N 096639 W 039065	416.26	60	36	6.0			5-83		SCREEN 32-42 GROUTED SAMPLE PUMP
199 H4 1874	6 P N 096473 W 040245	419.50	55	41	6.0	54		5-83		SCREEN 39-49 GROUTED SAMPLE PUMP

19

WELL DESIGNATION ----- EMA NO.	COORDINATES	CASING ELEV. (FT-MSL)	DRILL DEPTH (FT)	INIT. DEPTH (FT) WATER	TO DIA. (IN)	DEPTH TO BOTTOM (FT)	MIN-MAX PERFORATED DEPTH (FT)	DATE COMP. (M-Y)	FORMER DESIGNATION	COMMENTS
199 N 1988	N 086157 P W 060593	456.10	100	54	8.0	62	34 - 95	5-64		
	O	456.37	60	50	1.50		40 - 60	10-64		
	P	456.38	98	55	1.50		88 - 98	10-64		
	Q	456.36	74	55	1.50		72 - 74	10-64		
199 N 1984	N 086577 P W 060306	459.83	125	57	8.0	95	35-120	6-64		PLUG AT 95
	O	460.10	76	57	1.50		60 - 70	10-64		
	P	460.10	122	57	1.50	132	112-122	10-64		
	Q	460.10	107	61	1.50		105-107	10-64		
	R	460.10	79		1.50		77 - 79	10-64		
199 N 1896	N 086365 P W 060828	459.45	125	63	8.0	70	34 - 95	6-64		SAMPLE PUMP
	O	459.72	68		1.50		60 - 68	10-64		REMOVED
	P	459.71	122		1.50		112-122	10-64		REMOVED
	Q	459.72	98		1.50		96 - 98	10-64		REMOVED
	R	459.71	80		1.50		78 - 80	10-64		REMOVED
199 N 1899	N 085921 P W 060042	458.73	150	52	8.0	73	50 - 60	6-64		CEMENT BRIDGE @73 FT SAMPLE PUMP
	O	457.90	63	54	1.50		50 - 60	4-64		REMOVED 6-74
	P	457.90	138		1.50		128-138	10-64		REMOVED 6-74
	Q	457.90	123		1.50		113-123	10-64		REMOVED 6-74
	R	457.90	105		1.50		103-105	10-64		REMOVED 6-74
	S	457.90	78		1.50		76 - 78	10-64		REMOVED 6-74

23

WELL DESIGNATION ----- EPA NO.	COORDINATES	CASING ELEV. (FT-MSL)	INIT. DRILL DEPTH (FT) WATER	DEPTH TO DIA. (IN)	DEPTH TO BOTTOM (FT)	MIN-MAX PERFORATED DEPTH (FT)	DATE COMP. (M-Y)	FORMER DESIGNATION	COMMENTS
699 75 33A P	N 074590 W 023350	379.07	35	6.0	35		5-43	HR-11	FILLED IN WITH SILT
699 75 31B P	N 074690 W 023370	380.00	36	6.0	36		5-43	HR-12	FILLED IN WITH SILT
699 76 30		375.01	19	24.0	19		0-0	REF.2 NO.70	FILLED IN
699 76 30		414.00	41	36.0			0-0	13/25-3D1 REF.7	DUG WELL
699 77 34 P	N 076925 W 034275	397.24	21	72.0	21		0-0	T14N R27E 32-Q1	FILLED IN
699 77 36 P 4500	N 076700 W 036150	412.20	150 35	0.0	82	32 - 82	4-57		CEMENT PLUG AT 82FT SAMPLE PUMP
699 77 43 P	N 076600 W 042500	441.37	44	72.0	44		0-0	REF.2 NO.74 S1610	FILLED IN
699 77 54 P 4512	N 076700 W 054100	480.59	150 04	0.0	110	70 -120	5-57		CEMENT PLUG AT 120FT SAMPLE PUMP
699 78 36 4489		405.00	30	40.0			0-0	REF.2 NO.20 N.RUN NO.4	FILLED IN FARM WELL
699 78 45 P	N 078050 W 044625	432.00	36	60.0			0-0	REF.2 NO.111	FILLED IN
699 78 62 P 4511	N 077750 W 062300	469.00	150 76	0.0	109	70 -120	5-57		015 SCREEN 67-107 FT SAMPLE PUMP
699 79 104		774.00	699 370	16.0	703	429-678	2-53	PSN 515 14/25-31M1	ARMY CAMP WELL

274

WELL DESIGNATION ----- EMA NO.	COORDINATES	CASING ELEV. (FT-MSL)	DRILL DEPTH (FT)	INIT. DEPTH (FT) TO WATER	DIA. (IN)	DEPTH TO BOTTOM (FT)	MIN-MAX PERFORATED DEPTH (FT)	DATE COMP. (M-Y)	FORMER DESIGNATION	COMMENTS
199 B3 1 1851	N 071000 P W 079030	439.79	63	46	8.0	63	20 - 60	3-53	107-B-1	SAMPLE PUMP
199 B3 2 1852	N 071752 P W 078010	442.59	790	28	8.0	768	635-645	8-53	107-B-2	
1856	P	443.30	778		1.50		758-778	3-70		
1857	Q	442.81	642	42	1.50	429	632-642	3-70		
199 B4 1 1853	N 070000 P W 080650	461.80	90	65	8.0	83	50 - 90	2-49	100-B-1	SAMPLE PUMP
199 B4 2 1854	N 069933 P W 080672	461.99	90	63	6.0	86	62 - 90	2-49	100-B-2	PLUG AT 86
199 B4 3 1855	N 069933 P W 080636	461.75	91	63	8.0	86	60 - 90	2-49	100-B-3	PLUG AT 86
199 B4 4 1891	N 068978 P W 080367	472.14	105	76	8.0	96	49 -102	9-60		SAMPLE PUMP
199 B5 1895	N 069930 P W 082000	455.56	151	50	8.0	100	40 -148	8-62		PLUG AT 100
199 B8 1	N 067375 P W 080490	494.59	75		8.0	76	NONE	3-51	C431-1	NOT ACCESSIBLE
199 B8 2	N 067400 P W 080000	495.59	25		8.0	26	NONE	4-51	C431-2	CASING REMOVED
199 B8 3	N 067455 P W 081110	488.59	25		8.0	26	NONE	4-51	C431-3	CASING REMOVED
199 B8 4	N 067510 P W 081420	488.59	25		8.0	26	NONE	4-51	C431-4	CASING REMOVED

WELL DESIGNATION EMA NO.	COORDINATES	CASING ELEV. (FT-MSL)	INIT. DEPTH (FT) TO WATER	TO DIA. (IN)	DEPTH TO BOTTOM (FT)	MIN-MAX PERFORATED DEPTH (FT)	DATE COMP. (M-Y)	FORMER DESIGNATION	COMMENTS
199 D5 7	N 093520 P W 052140	465.57	35	8.0	36		3-49	107-DR-4	CASING REMOVED
199 D5 8	N 093520 P W 052090	465.57	31	8.0	32		3-49	107-DR-5	CASING REMOVED
199 D5 9	N 093520 P W 051990	462.57	35	8.0	36		3-49	107-DR-6	CASING REMOVED
199 D5 10		464.57	28 DRY	8.0	29	14 - 27	4-57	100-DR-1	
199 D5 11		464.57	27 DRY	8.0	28	14 - 27	4-57	100-DR-2	
199 D5 12 1892	N 092125 P W 052546	469.67	91 85	8.0	91	35 - 90	8-60		SAMPLE PUMP
199 D7 1		420.00	51	48.0	51		0-0	T14N R26E 14-F1	FILLED IN
199 D7 2		420.00	39	84.0	102		0-0	T14N R26E 14-F2	FILLED IN
199 D8 1	N 095230 P W 053140	400.00	37	6.0	37		5-43	HR-9	FILLED IN
199 D8 1861	N 094725 P W 053018	444.61	77 DRY	8.0	44	30 - 75	6-52	107-D-1	
199 D8 1862	N 094720 P W 052205	449.06	81 62	6.0	81	35 - 79	6-52	107-D-2	SAMPLE PUMP

16

WELL DESIGNATION		COORDINATES	CASING ELEV. (FT-MSL)	DRILL DEPTH (FT)	INIT. DEPTH (FT) TO WATER	DIA. (IN)	DEPTH TO BOTTOM (FT)	MIN-MAX PERFORATED DEPTH (FT)	DATE COMP. (M-Y)	FORMER DESIGNATION	COMMENTS
EMA NO.											
199	K 13		464.00	159	72	12.0	138		3-53	100-K	OIL IN WELL
199	K 14		469.85	95		8.0	95		12-52	105-KW-1	
199	K 15	N 077160 P W 069050	408.80	150		6.0	150		4-43	U.S.E.D.TH. NO. 1	
199	K 16	N 076300 P W 067800	404.80	50		8.0	50		2-53	105-KE-1	
199	K 17		406.00	75		8.0	75	50 - 75	9-53	105-KE-2	
199	K 18		409.00	60	21	8.0	40		10-54	107-KE-3	
1879											
199	K 19	N 078000 P W 067000	422.17	51	30	8.0	51	10 - 50	4-55	107-KE-4	SCREEN 26-46 FT. SAMPLE PUMP
1884											
199	K 20	N 079500 P W 066125	422.57	50	31	8.0	48	10 - 50	5-55	107-KE-5	SAMPLE PUMP
1885											
199	K 21	N 080000 P W 066000	421.73	50	DRY	8.0	16	10 - 50	5-55	107-KE-6	
1886											
199	K 22	N 081000 P W 065000	421.60	50	30	8.0	49	10 - 50	5-55	107-KE-7	Ø15 SCREEN 29-49 FT. SAMPLE PUMP
1887											
199	K 23	N 078000 P W 068000	405.80	80	DRY	8.0	25	65 - 80	2-56	1706-KER-1	
199	K 24	N 077000 P W 069000	467.00	50		8.0	50	NONE	12-52	105-KW-2	

21

REFERENCE 4

Draft Phase I Installation Assessment of Inactive Waste-Disposal
Sites at Hanford, Volume 1, July 1986

91113221071

2.0 DESCRIPTION OF HANFORD SITE

This section provides a summary of environmental conditions at the Hanford Site and a brief discussion of the Site's purpose and history. It also describes specific environmental features and the process history of each operational area (i.e., the 100, 200, 300, 400, and 600 Areas).

2.1 ENVIRONMENTAL SUMMARY

The semiarid Hanford Site, operated by the DOE, occupies about 1,476 square kilometers (570 sq mi) of the southeastern part of Washington State north of where the Yakima River flows into the Columbia (see Figure 2.1). The Site lies about 320 kilometers (200 mi) east of Portland, Oregon, 270 kilometers (170 mi) southeast of Seattle, Washington, and 200 kilometers (125 mi) southwest of Spokane, Washington.

Environmental conditions common to all areas at Hanford Site are summarized below. Descriptions of these environmental aspects are based on several recent reports (U.S. DOE 1984; Sommer et al. 1981; Yandon 1977; U.S. ERDA 1975).

2.1.1 Geology and Soils

The Hanford Site lies in the Pasco Basin, a structural and topographic basin of eastern Washington and the Columbia River Basalt Plateau. The region is underlain by three geologic units. In ascending order these are: 1) the sequential beds of basaltic lavas and interbed sediments of the Columbia River Basalt Group at the base; 2) the Pliocene-age Ringold Formation (lacustrine formation), consisting of well-rounded pebbles and cobbles with interstitial spaces filled with medium sand; and 3) the Hanford Formation, consisting of the Pasco (glaciofluvial) gravels and associated sediments of the late Pleistocene age lying at the surface.

The surface geology of the Site is characterized by a surface layer of light brown, fine, slightly silty, wind-deposited sand, sparsely covered by vegetation. Although the surface soil is fertile, it has little agricultural value without irrigation. Underlying the surface sands is a mixture of sand

91118901033




FIGURE 2.1. Features of the Hanford Site.

and gravel extending to a depth of about 60 meters (200 ft). Basaltic rock starts at that depth and extends downward over 3000 meters (1.9 mi).

Elevations range from a low of about 105 meters (345 ft) above mean sea level (MSL) in the southeastern part of the Hanford Site to a maximum of 1,091 meters (3,579 ft) at the crest of Rattlesnake Mountain to the west. (See Section 2.3 for a discussion of geologic features peculiar to each operational area.)

2.1.2 Meteorology

The Site lies east of the Cascade Mountains and, as a result, has a semiarid climate reflecting the rainshadow effect of the mountains. The average annual precipitation for the Site is about 160 millimeters (6.3 in.). Ten percent of this amount falls from July through September, and 42% falls from November through January. The greatest amount of rainfall recorded in a 12-hour period was 47.8 millimeters (1.9 in.).

Because of the limited rainfall, surface runoff from the Hanford Site is minimal. The annual precipitation mostly evaporates, resulting in small amounts of water available for runoff or infiltration.

2.1.3 Hydrology and Hydrogeology

The Columbia River (the fifth largest river by volume in North America) is the dominant aquatic ecosystem on the Hanford Site. Numerous dams have been built on the river. The only free-flowing section in the United States is between Priest Rapids Dam and McNary Reservoir, along the Hanford Site. No significant tributaries enter the stream in this section.

The Columbia has a long-term annual average flow of about 3,600 cubic meters per second (127,000 cfs). [The Yakima River, by comparison, flows an average of about 90 cubic meters per second (3,180 cfs).] The flow rates of the Columbia are influenced by water usage and upstream reservoir projects. The reservoirs provide active storage of more than 4.6×10^{10} cubic meters (37,000,000 acre-feet) of water.

The uppermost aquifer in the Pasco Basin is an unconfined system within the Hanford and Ringold Formations. The elevation of this aquifer ranges from

excluded from public use by the DOE and are used as a wildlife refuge and for DOE environmental research. The land north of the Columbia River is controlled by the Washington State Department of Game and the U.S. Fish and Wildlife Service as controlled hunting areas and a game refuge.

Land use within a 50-kilometer (30-mi) radius of the Site includes residential, suburban, corporate city, agricultural, industrial and commercial, scenic, recreational, and general use areas. The predominant use of lands within the 50-kilometer (30-mi) radius is agricultural, with farms located along or near all the Site boundaries.

2.1.6 Population

Population in the area surrounding the Hanford Site is sparse, consisting primarily of farms and farming communities to the north, east, and west of the Site. The Tri-Cities (Kennewick, Pasco, and Richland), located to the south and southeast of the Hanford Site, represent the major population concentration in the area (Sommer et al. 1981).

In 1980, an estimated 341,000 people were living within an 80-kilometer (50-mi) radius of the Hanford Meteorological Station (HMS) (see Figure 2.4); it is estimated that the number will grow to 417,000 by 1990.

2.1.7 Air Quality

Air quality in the vicinity of the Hanford Site is generally quite good. Wind-eroded dust is a problem in the area, and the dust storms that occur in the region can produce high total-suspended particulate concentrations. However, on both an annual and a short-term basis, the region is in compliance with the National Ambient Air Quality Standards (NAAQS) for particulates. All other pollutant levels also satisfy the federal and State of Washington standards (U.S. DOE 1984).

2.2 PURPOSE AND HISTORY

In 1943, after the Fermi experiment showed that nuclear fission could be controlled in a small reactor, the U.S. Army Corps of Engineers selected Hanford as the location to build larger versions of the Fermi reactor to produce plutonium for possible use in military weapons. Construction started




FIGURE 2.4. U.S. Census Populations for 1980 of Cities Within 80 Kilometers of the Hanford Meteorological Station (U.S. DOE 1986)

in l
fac
afte
4 mc

were
Colu
reac

faci
ficat
heavi

other
pluto
Since
gener

A
older
eight
only o

2.3 Di

Er

below;
is ider
(e.g.,
provide
waste-p

2.3.1

The
River in

91118901035

in March 1943 on three reactor facilities and three chemical processing facilities. The first of the reactors went into operation about 18 months after the start of construction, and the first plutonium was available some 4 months later.

After World War II, five reactors similar to those built during the war were constructed. A total of eight graphite-moderated reactors used the Columbia River for once-through cooling (i.e., water circulated through the reactors only once before being released back to the river).

Early in the 1950s construction began on the research and development facilities known as the Hanford Laboratories. This marked the first diversification of Hanford from a purely defense-materials production facility to one heavily involved in peacetime uses of the atom.

In 1963 the N Reactor was built. The N Reactor is different from the other eight reactors in that it generates steam as a by-product of the plutonium production and does not use river water as a once-through coolant. Since 1966 the Washington Public Power Supply System has used the steam to generate electricity.

A presidential decision was made in early 1964 to begin shutting down the older Hanford reactors. This decision resulted in the closing down of all eight of the older reactors by the end of 1971, leaving the N Reactor as the only operational production reactor.

2.3 DESCRIPTION OF OPERATIONAL AREAS

Environmental features specific to each operational area are described below; the waste-processing history of each area is also discussed. Each area is identified by number (i.e., 100, 200, 300, 400, and 600) and by letter (e.g., the 100-F Area is the location of the 100-Area F Reactor). Appendix B provides further information on waste-disposal site locations and types of waste-processing facilities in Hanford's operational areas.

2.3.1 100 Areas

The nine 100 Areas (B, C, D, DR, KE, KW, F, H, N) border the Columbia River in the northernmost part of the Hanford Site. Each of the nine areas

has one production reactor. Eight of these reactors have been shut down; only the N Reactor, used for both plutonium and electricity production, is still operating. Because some of the areas are contiguous (B/C, D/DR, KE/KW), the Hanford Site map shows only six 100 Areas (Figure 2.1).

The 100 Areas are generally flat with no major surface features. The Hanford Formation lies near the surface of the 100 Areas, covered by a thin layer of wind-deposited silt and fine sand. The water table is found in these sediments at a depth of about 20 meters (66 ft), except in the F and H Areas where the depth to the water table is about 35 meters (115 ft) and 40 meters (131 ft), respectively. The depth to the Ringold Formation is about 25 meters (82 ft); the top of the basalt bedrock is approximately 240 meters (790 ft) below the surface.

Because the water table occurs within the highly permeable sandy gravels of the Hanford Formation, it fluctuates as the river level rises and falls. The ground water generally flows from the 100 Areas and toward the river. When active, each of the 100 Areas included support facilities such as powerhouses. Except for 100-N, these powerhouses produced process steam from coal-fired boilers; 100-N has oil-fired boilers. Adjacent to each area's powerhouse were large storage areas that received railroad carloads of coal and disposal areas for flyash/clinker disposal. Most areas also included water-treatment plants, water-storage tanks, subsurface sewage-disposal systems, raw-water intake structures, and process sewers.

B and C Areas. The B and C reactors are located adjacent to each other on a 2.6-square-kilometer (650-acre) site (the 100 B/C Area) and are the farthest upstream of the 100 Areas. The B Reactor was operated from 1944 to 1968, and the C Reactor was operated from 1952 to 1969. Virtually all the facilities in the area are inactive, with the exception of the B/C export water system, which continues to provide the raw water supply to the 200 Area and some 100 Areas. An electrical substation in the area taps power for the pumps providing the 200-Area water. Fewer than 100 people work in this area (Yandon 1977).

When the reactors were operational, cooling water was drawn from the river and treated with alum, sulfuric acid, and chlorine. Excess sulfuric

acid was used to maintain the pH of the water within a desired range. To control oxidation of aluminum parts in the reactor, sodium dichromate was used to maintain an oxidation coating on aluminum parts. The chlorine was added for algae control in the settling basins; at times copper sulfate was added for additional algae control. Chromic acid, oxalic acid, and nitric acid were used for dummy fuel-element decontamination.

In addition to vertical safety rods for emergency reactor shutdown, the reactors were equipped with hoppers of nickel-plated boron steel balls, nickel-plated carbon steel balls, and stainless steel balls that would drop into the vertical safety rod channels for emergency shutdown. This system required no supplementary power source. Although it was never used, a third safety system, one involving the use of a potassium borate solution, was in place at the reactors.


A supplementary control system, in addition to the normal horizontal control rods, was incorporated into the reactors. This supplementary control system consisted of a Poison^(a) Column Control Facility that could charge selected process tubes with a lead-cadmium poison to absorb neutrons. Boron-carbide aluminum poison splines were also used for supplementary control.

The coolant water system and backup control and shutdown systems at the other 7 once-through-cooled reactors were similar to the those in 100 B/C Area.


D and DR Areas. The 100-D/DR Areas, covering about 3.9 square kilometers (970 acres), are located 11 kilometers (7 mi) downriver of the 100-B/C Area. The D Reactor was operated from 1944 to 1967 and the DR Reactor from 1950 to 1965. These areas are extensively used, and their utilities and services are still in operation. The electrical substation serves as a backup supply for the 100-N Area. The water system is a backup system for the 100-B water system, which supplies water to the 200 Areas. The UNC Nuclear Industries engineering laboratory here is operated in support of the N Reactor. Approximately 20 people are employed in the D and DR Areas (Yandon 1977).

(a) The term poison refers to a material's ability to absorb neutrons and thus control the rate of fission.

Number
scribble
only in
For
area;
of
numbering



9 1 1 1 8 9 0 1 0 4 0



BUILDING LIST

Building Number	Description	Building Number	Description
100-1	...	100-10	...
100-2	...	100-11	...
100-3	...	100-12	...
100-4	...	100-13	...
100-5	...	100-14	...
100-6	...	100-15	...
100-7	...	100-16	...
100-8	...	100-17	...
100-9	...	100-18	...
100-10	...	100-19	...
100-11	...	100-20	...
100-12	...	100-21	...
100-13	...	100-22	...
100-14	...	100-23	...
100-15	...	100-24	...
100-16	...	100-25	...
100-17	...	100-26	...
100-18	...	100-27	...
100-19	...	100-28	...
100-20	...	100-29	...
100-21	...	100-30	...
100-22	...	100-31	...
100-23	...	100-32	...
100-24	...	100-33	...
100-25	...	100-34	...
100-26	...	100-35	...
100-27	...	100-36	...
100-28	...	100-37	...
100-29	...	100-38	...
100-30	...	100-39	...
100-31	...	100-40	...
100-32	...	100-41	...
100-33	...	100-42	...
100-34	...	100-43	...
100-35	...	100-44	...
100-36	...	100-45	...
100-37	...	100-46	...
100-38	...	100-47	...
100-39	...	100-48	...
100-40	...	100-49	...
100-41	...	100-50	...
100-42	...	100-51	...
100-43	...	100-52	...
100-44	...	100-53	...
100-45	...	100-54	...
100-46	...	100-55	...
100-47	...	100-56	...
100-48	...	100-57	...
100-49	...	100-58	...
100-50	...	100-59	...
100-51	...	100-60	...
100-52	...	100-61	...
100-53	...	100-62	...
100-54	...	100-63	...
100-55	...	100-64	...
100-56	...	100-65	...
100-57	...	100-66	...
100-58	...	100-67	...
100-59	...	100-68	...
100-60	...	100-69	...
100-61	...	100-70	...
100-62	...	100-71	...
100-63	...	100-72	...
100-64	...	100-73	...
100-65	...	100-74	...
100-66	...	100-75	...
100-67	...	100-76	...
100-68	...	100-77	...
100-69	...	100-78	...
100-70	...	100-79	...
100-71	...	100-80	...
100-72	...	100-81	...
100-73	...	100-82	...
100-74	...	100-83	...
100-75	...	100-84	...
100-76	...	100-85	...
100-77	...	100-86	...
100-78	...	100-87	...
100-79	...	100-88	...
100-80	...	100-89	...
100-81	...	100-90	...
100-82	...	100-91	...
100-83	...	100-92	...
100-84	...	100-93	...
100-85	...	100-94	...
100-86	...	100-95	...
100-87	...	100-96	...
100-88	...	100-97	...
100-89	...	100-98	...
100-90	...	100-99	...

U.S. Department of Energy
 Pacific Northwest Laboratory
 100-N AREA
 WASTE SITES

H-3-5720 7 10

9111971040

REFERENCE 5

Environmental Monitoring at Hanford for 1986, PNL-6120, May 1987

91118901044

2.0. BACKGROUND INFORMATION

2.1. DESCRIPTION OF THE HANFORD SITE

K. R. Price, P. J. Mitchell, and M. D. Freshley

The U.S. Department of Energy's Hanford Site is located in a rural region of southeastern Washington and occupies an area of 1,500 km². The Site (shown in Figure 2.1) lies about 320 km northeast of Portland, Oregon, 270 km southeast of Seattle, Washington, and 200 km southwest of Spokane, Washington. The Columbia River flows through the northern edge of the Hanford Site and forms part of the eastern boundary. The southern boundary of the Site includes the Rattlesnake Hills, which exceed 1000 m in elevation. Both confined and unconfined aquifers are present beneath the Site. The main geologic units are the Columbia River Basalt Group, the Ringold Formation, and a series of glaciofluvial sediments. The Hanford Project was established in 1943 and was originally designed, built, and operated to produce plutonium for nuclear weapons.

SURFACE CHARACTERISTICS OF THE SITE

The semiarid land on which the Hanford Site is located has a sparse covering of desert shrubs and drought-resistant grasses. The most broadly distributed type of vegetation on the Site is the sagebrush/cheatgrass/bluegrass community. Most abundant of the mammals is the Great Basin pocket mouse. Of the big-game animals, the mule deer is the most abundant, while the cottontail rabbit is the most abundant of the small-game animals. Coyotes are also abundant. The bald eagle is a regular winter visitor to the relatively large areas of uninhabited land comprising the Hanford Site.

The Columbia River, which originates in the mountains of eastern British Columbia, Canada, flows through the northern edge of the Hanford Site and forms part of the Hanford Site's eastern boundary. The river drains a total area of approximately 70,800 km² enroute to the Pacific Ocean. The flow of the Columbia River is regulated by 11 dams within the United States, 7 upstream and 4 downstream of the Site. Priest Rapids Dam is the nearest impoundment upstream of the Site, and McNary Dam is the nearest dam downstream. (The Hanford reach of the Columbia River extends from Priest Rapids Dam to the head of Lake Wallula, which is created by McNary Dam.) This is the only stretch of the Columbia River within the U.S. that is not impounded by a dam. The width of the river

varies from approximately 300 m to about 1000 m. The flow through this stretch of the river is relatively swift, with numerous bends and several islands present throughout the reach.

The flow rate of the Columbia River in this region is regulated primarily by Priest Rapids Dam. Hanford reach flows fluctuate significantly because of the relatively small storage capacity and operational practices of the nearby upstream dams. A minimum flow rate of 1,000 cubic meters per second (m³/s) [36,000 cubic feet per second (cfs)] has been established at Priest Rapids. Typical daily flows range from 1,000 m³/s (36,000 cfs) to 7,000 m³/s (250,000 cfs) with peak spring runoff flows of up to 12,600 m³/s (450,000 cfs) being recorded. Typical annual average flows at Priest Rapids Dam are 3,100 m³/s (110,000 cfs) to 3,400 m³/s (120,000 cfs). Monthly mean flows typically peak from April through June and are at the lowest levels from September through October.

The temperature of the Columbia River varies seasonally. Minimum temperatures are observed during January and February while maximum temperatures typically occur during August and September. Monthly temperatures for the river range from approximately 3°C to about 20°C during the course of a year. Water storage management practices at upstream dams and the flow rate of the river dictate, to a large extent, the thermal characteristics of the Columbia River along the Hanford reach.

91118901045

91118901046




FIGURE 2.1. DOE's Hanford Site

91118901047

The Columbia River system has been developed extensively for hydroelectric power, flood control, navigation, irrigation, and municipal and industrial water supplies. In addition, the Hanford reach is used for a variety of recreational activities including fishing, hunting, boating, water skiing, and swimming. The State of Washington has classified the stretch of the Columbia River from the Washington-Oregon border to Grand Coulee Dam (which includes the Hanford reach) as Class A and established water quality criteria and water use guidelines for this class designation. Because these criteria do not include specific limits for radionuclides, Environmental Protection Agency (EPA) and State of Washington drinking water limits were used for comparison. Other surface water on the Site consists of West Lake (a small, natural pond) and a number of ditches and artificial ponds created for routine disposal of waste water.

Hanford's climate is dry and mild; the area receives approximately 16 cm of precipitation annually. About 40% of the total precipitation occurs during November, December, and January; only 10% falls in July, August, and September. Approximately 45% of all precipitation from December through February is snow. The average minimum and maximum temperatures in July are 16°C and 32°C. For January, the average temperatures are 3°C and -6°C.

Monthly average wind speeds range from about 10 km/h in the summer to 14 km/h in the winter. The prevailing regional winds are from the northwest, with occasional cold-air drainage into valleys and occurrences of strong crosswinds. The region is a typical desert area with frequent strong inversions that occur at night and break during the day, resulting in unstable and turbulent wind conditions.

Land near the Hanford Site is primarily used for agriculture and for livestock grazing. Agricultural lands are found north and east of the Columbia River and south of the Yakima River. These areas contain orchards, vineyards, and fields of alfalfa, wheat, and vegetables. The Hanford Site north of the Columbia River is shared between a state wildlife management area and a federal wildlife refuge. The northeast slope of the Rattlesnake Hills along the southwestern boundary of the Site is designated as the Arid Lands Ecology Reserve (ALE) and is used for ecological research by DOE.

The major population center nearest to the Hanford Site is the Tri-Cities area (Richland, Pasco, and Kennewick), which is situated on the Columbia River downstream from the Site and has a population of approximately 90,000. Approximately 340,000 people live within an 80-km radius of the Hanford Site. This number includes people living in the Tri-Cities, the Yakima area, several small communities, and the surrounding agricultural area. More detail on Site characteristics and activities is available in "The Final Environmental Statement, Waste Management Operations, Hanford Reservation" (ERDA 1975).

SUBSURFACE CHARACTERISTICS OF THE SITE

The DOE operations on the Site have resulted in the production of large volumes of waste water that have historically been discharged to the ground through cribs, ditches, and ponds. These discharges greatly influence the physics and chemistry of the subsurface. Approximately 25 billion liters of liquid effluent in the 200 Areas and 2.6 billion liters of liquid effluent in the 100N Area were disposed to the ground during 1986, including process cooling water and water containing low-level radioactive wastes. The discharge of waste water to the ground at the Hanford Site began in the mid-forties and reached a peak in 1955. After 1955, discharge to cribs declined because of improved treatment of waste streams and the deactivation of various facilities (Graham et al. 1981). Since the restart of the Plutonium and Uranium Extraction (PUREX) Plant and related facilities in late 1983, discharge of PUREX-related effluents has resumed.

Subsurface structures, such as cribs, have primarily been used for the disposal of water containing radioactive wastes, while surface ponds and ditches have primarily been used for the disposal of uncontaminated cooling water (Graham et al. 1981). Sanitary wastes are discharged to the ground via tile fields. The majority of liquid disposal occurred in the Separations Area, which includes the 200-East (200E) and 200-West (200W) Areas (Figure 2.1). Smaller amounts of waste water were disposed in the 100 and 300 Areas. Discharges of waste water to the ground in the 400 Area were minimal.

the Ringold formation has been removed. These sediments were deposited by the ancestral Columbia River when it was swollen by glacial meltwater. The glaciofluvial sediments consist primarily of gravels and sands, with some silts (Newcomb, Strand and Frank 1972).

Hydrology

Both confined and unconfined aquifers are present beneath the Hanford Site. The confined aquifers, in which the ground water is under pressure greater than that of the atmosphere, are found primarily within the Columbia River basalts. In general, the unconfined or water-table aquifer is located in the Ringold Formation and glaciofluvial sediments, as well as some more recent alluvial sediments in areas adjacent to the Columbia River (Gephart et al. 1979). This relatively shallow aquifer has been affected by waste-water disposal at Hanford more than the confined aquifers (Graham et al. 1981). Therefore, the unconfined aquifer is the most thoroughly monitored aquifer beneath the Site.

The unconfined aquifer is bounded below by either the basalt surface or, in places, the relatively impervious clays and silts of the lower unit of the Ringold Formation. Laterally, the unconfined aquifer is bounded by the anticlinal basalt ridges that ring the basin and by the Yakima and Columbia rivers. The basalt ridges above the water table have a low permeability and act as a barrier to lateral flow of the ground water (Gephart et al. 1979). The saturated thickness of the unconfined aquifer is greater than 61 m in some areas of the Hanford Site and pinches out along the flanks of the basalt anticlines. The depth from the ground surface to the water table ranges from less than 0.3 m near the Columbia River to over 106 m in the center of the Site. The elevation of the water table above mean sea level for June of 1986 is shown in Figure 2.3.

Recharge to the unconfined aquifer originates from several sources (Graham et al. 1981). Natural recharge occurs from precipitation at higher elevations and runoff from ephemeral streams to the west, such as Cold Creek and Dry Creek. The Yakima River recharges the unconfined aquifer as it flows along the southwest boundary of the Hanford Site. The Columbia River recharges the unconfined aquifer during high stages when river water is transferred to the aquifer along the river bank. The unconfined aquifer receives little, if any, recharge from pre-

cipitation directly on the Hanford Site because of a high rate of evapotranspiration under native soil and vegetation conditions. However, present studies, such as those described by Heller, Gee, and Meyers (1985), suggest that precipitation may contribute more recharge to the ground water than was originally thought.

Large scale artificial recharge occurs from offsite agricultural irrigation and liquid-waste disposal in the operating areas at Hanford. Recharge from irrigation in the Cold Creek Valley enters the Hanford Site as ground-water flow across the western boundary. Artificial recharge from waste-water disposal at Hanford occurs principally in the Separations Area. It was estimated that recharge to the ground water from facilities in the Separations Area (including B Pond and Gable Mountain Pond, as well as the various cribs and trenches in the 200W and 200E Areas) adds ten times as great an annual volume of water to the unconfined aquifer as is contributed by natural inflow to the area from precipitation and irrigation waters to the west (Graham et al. 1981).

The operational discharge of water has created ground-water mounds near each of the major waste-water disposal facilities in the Separations Area and in the 100 and 300 Areas (Figure 2.3). These mounds have altered the local flow pattern in the aquifer, which is generally from the recharge areas in the west to the discharge areas (primarily the Columbia River) in the east. Water levels in the unconfined aquifer have changed continuously during Site operations because of variations in the volume of waste water discharged. Consequently, the movement of ground water and its associated constituents has also changed with time.

In addition to the Separations Area, ground-water mounding also occurs in the 100 and 300 Areas. Ground-water mounding in these areas is not as significant as in the Separations Area because of differences in discharge volumes and subsurface geology. However, in the 100 and 300 Areas, water levels are also greatly influenced by river stage.

Liquid Effluent Movement

If significant quantities of liquid effluents are discharged to the ground at the Hanford Site waste disposal facilities, then these effluents would percolate downward through the unsaturated zone to the water table. As

100 Area Background and Down Gradient Wells




FIGURE 3.10. Location of Ground-Water Monitoring Wells Sampled in 1986 (first digit of well number has been dropped)

91118901049

REFERENCE 6

Draft Phase I Installation Assessment of Inactive
Waste-Disposal Site at Hanford, Volume 2, July 1986

NOTE: This site contains numerous sheets
from the document; reference is made
to the document instead of attaching
all of the individual sheets.

91118901050

REFERENCE 7

Uncontrolled Hazardous Waste Site Ranking System; A Users Manual,

40 CFR 300, Appendix A

91118901051

Part 300, App. A

TABLE 5—PERSISTENCE (BIODEGRADABILITY) OF SOME ORGANIC COMPOUNDS*—Continued

chloromethyl ethyl ether 3 chloropyridine o 1 butyl p benzoguanone diisobutyl ether dihydrocarvone dimethyl sulfide 2,6 dimethylurea	methylenedio methylene chloride nitrobenzene nitrobenzene 1,1,2 trichloroethylene isomethyl bromo-hydroxy- bisphenol A
Value = 1 Somewhat Persistent Compounds	
acetylene dichloride benzoic acid, methyl ester benzene benzene sulfonic acid butyl benzene butyl bromide o-cresol carbon disulfide o-cresol decane 1,2 dichloroethane 1,2 dimethyl benzene 1,3 dimethyl naphthalene 1,4 dimethyl phenol diethyl adipate n-dodecane ethyl benzene p-ethyl n-hexane p-ethyltoluene hexadecane isopropyl benzene	bronans methyl ester of tyrosine acid methane 2-methyl-6-ethyl pyridine methyl naphthalene methyl paraffins methyl phenyl carbonyl methyl stearate naphthalene nursene octane octyl chloride paraffin phenyl benzene propyl acrylate propylbenzene 1-terphenyl toluene vinyl benzene xylene
Value = 5 Nonpersistent Compounds	
acetaldehyde acetic acid acetone acetylacetone benzoic acid o 1 butyl carbonyl sucrose sucrose sucrose sucrose sucrose sucrose sucrose sucrose sucrose sucrose sucrose	methyl benzoate 3-methyl butanol methyl ethyl ketone 3-methylpentanol pentadecane pentanol propyl propylene tetradecane tridecane n-undecane

* JIS Associates, Inc., Methodology for Assessing the Hazard Potential for Waste Disposal Sites, May 8, 1980

Toxicity of each hazardous substance being evaluated is given a value using the rating scheme of Sax (Table 6) or the National Fire Protection Association (NFPA) (Table 7) and the following guidance:

Toxicity	Assigned value
Sax level 0 or NFPA level 0	0
Sax level 1 or NFPA level 1	1
Sax level 2 or NFPA level 2	2
Sax level 3 or NFPA level 3 or 4	3

40 CFR Ch. I (7-1-86 Edition)

TABLE 6—SAX TOXICITY RATINGS

0—No Toxicity* (None)**
The designation is given to materials which fall into one of the following categories: (a) Materials which cause no harm under any conditions of normal use (b) Materials which produce toxic effects on humans only under the most unusual conditions or by overwhelming dosage
1—Slight Toxicity* (Low)**
(a) Acute local Materials which on single exposures lasting seconds, minutes, or hours cause only slight effects on the skin or mucous membranes regardless of the extent of the exposure (b) Acute systemic Materials which can be absorbed into the body by inhalation, ingestion, or through the skin and which produce only slight effects following single exposures lasting seconds, minutes, or hours, or following ingestion of a single dose regardless of the quantity absorbed or the extent of exposure (c) Chronic local Materials which on continuous or repeated exposures extending over periods of days, months, or years cause only slight and usually reversible harm to the skin or mucous membranes. The extent of exposure may be great or small (d) Chronic systemic Materials which can be absorbed into the body by inhalation, ingestion, or through the skin and which produce only slightly usually reversible effects extending over days, months, or years. The extent of the exposure may be great or small In general, these substances classified as having "slight toxicity" produce changes in the human body which are readily reversible and which will disappear following termination of exposure, either with or without medical treatment
2—Moderate Toxicity* (Med)**
(a) Acute local Materials which on single exposures lasting seconds, minutes, or hours cause moderate effects on the skin or mucous membranes. These effects may be the result of intense exposure for a matter of seconds or moderate exposure for a matter of hours (b) Acute systemic Materials which can be absorbed into the body by inhalation, ingestion, or through the skin and produce moderate effects following single exposures lasting seconds, minutes, or hours, or following ingestion of a single dose

Environmental Protection Agency

TABLE 6—SAX TOXICITY RATINGS—Continued

3—Moderate Toxicity* (Med)**
(a) Chronic local Materials which on continuous or repeated exposures extending over periods of days, months, or years cause moderate harm to the skin or mucous membranes (b) Chronic systemic Materials which can be absorbed into the body by inhalation, ingestion, or through the skin and which produce moderate effects following continuous or repeated exposures extending over periods of days, months, or years These substances classified as having "moderate toxicity" may produce irreversible as well as reversible changes in the human body. These changes are not of such severity as to threaten life or to produce serious physical impairment
4—Severe Toxicity* (High)**
(a) Acute local Materials which on single exposures lasting seconds or minutes cause injury to skin or mucous membranes of sufficient severity to threaten life or to cause permanent physical impairment or disfigurement (b) Acute systemic Materials which can be absorbed into the body by inhalation, ingestion, or through the skin and which can cause injury of sufficient severity to threaten life following a single exposure lasting seconds, minutes, or hours, or following ingestion of a single dose (c) Chronic local Materials which on continuous or repeated exposures extending over periods of days, months, or years can cause injury to skin or mucous membranes of sufficient severity to threaten life or cause permanent impairment, which disfigurement, or irreversible change (d) Chronic systemic Materials which can be absorbed into the body by inhalation, ingestion or through the skin and which can cause death or serious physical impairment following continuous or repeated exposures to small amounts extending over periods of days, months, or years

*Sax, N. I., Dangerous Properties of Industrial Materials, Van Nostrand Reinhold Co., New York, New York, 4th edition, 1974
**Sax, N. I., Dangerous Properties of Industrial Materials, Van Nostrand Reinhold Co., New York, New York, 5th edition, 1978

TABLE 7—NFPA TOXICITY RATINGS*

0	Materials which on exposure under the conditions would offer no health hazard beyond that of ordinary combustible material
1	Materials only slightly hazardous to health. It may be desirable to wear self-contained breathing apparatus.
2	Materials hazardous to health, but areas may be entered freely with self-contained breathing apparatus.
3	Materials extremely hazardous to health, but areas may be entered with extreme care. Full protective clothing, including self-contained breathing apparatus, rubber gloves, boots and bands around legs, arms and waist should be provided. No skin surface should be exposed.
4	A low whiff of the gas or vapor could cause death, or the gas, vapor, or liquid could be fatal on penetrating the fire fighters' normal full protective clothing which is designed for resistance to heat. For most chemicals having a health 4 rating, the normal full protective clothing available to the average fire department will not provide adequate protection against skin contact with these materials. Only special protective clothing designed to protect against the specific hazard should be worn.

Part 300, App. A

*National Fire Protection Association National Fire Codes, Vol. 12, No. 48, 1977.

Table 4 presents values for some common compounds.

Hazardous waste quantity includes all hazardous substances at a facility (as received) except that with a containment value of 0. Do not include amounts of contaminated soil or water; in such cases, the amount of contaminating hazardous substance may be estimated.

On occasion, it may be necessary to convert data to a common unit to combine them. In such cases, 1 ton = 1 cubic yard = 6 drums and for the purposes of converting bulk storage, 1 drum = 55 gallons. Assign a value as follows:

Tons in cubic yards	Number of drums	Assigned value
0	0	0
1-10	1-40	1
11-62	41-250	2
63-126	251-500	3
127-250	501-1,000	4
251-500	1,001-2,500	5
501-1,250	2,501-5,000	6
1,251-2,500	5,001-10,000	7
>2,500	>10,000	8

3.5 Targets. Ground water use indicates the nature of the use made of ground water drawn from the aquifer of concern within 3 miles of the hazardous substance, including the geographical extent of the measurable concentration in the aquifer. Assign a value using the following guidance:

Ground water use	Assigned value
Unusable (e.g., extremely saline aquifer, or extremely low yield, etc.)	0
Commercial, industrial or irrigation and another water source presently available, not used, but usable	1
Drinking water with municipal water from alternate unrestricted sources presently available (e.g., minimal hookup requirements), or commercial, industrial or irrigation with no other water source presently available	2
Drinking water, no municipal water from alternate unrestricted sources presently available	3

Distance to nearest well and population served have been combined in the matrix below to better reflect the important relationship between the distance of a population from hazardous substances and the size of the population served by ground water that might be contaminated by those substances. To determine the overall value for this combined factor, score each individual



Ref. 7.2

REFERENCE 8

U.S.G.S. Maps of the area around the 100 Area:

Coyote Rapids, Wash 15 Minute Map
Smyrna, Wash 15 Minute Map
Corfu, Wash 15 Minute Map
Priest Rapids, Wash 15 Minute Map
Hanford, Wash 15 Minute Map

91118901054



40 FEET
FOOT CONTOURS
LEVEL

ACCURACY STANDARDS
GSD 80225, OR WASHINGTON, D. C. 20242
GSD 80225 IS AVAILABLE ON REQUEST



ROAD CLASSIFICATION

Medium-duty ——— Light-duty - - - - -
Unimproved dirt - - - - -

○ State Route

The NW, SW, and SE quarters also covered by 7.5 minute
1:24 000-scale maps: Priest Rapids 1950, Carrn Hope
Peak 1950, and Emerson Nipple 1950



PRIEST RAPIDS, WASH.
N4630—W11945/15

1965


AMS 2077 III—SERIES V791

CORRAL CANYON
2078

6
9976



ROAD CLASSIFICATION
 Medium-duty ——— Light duty ———
 Unimproved dirt - - - - -
 State Route ○





CORFU, WASH.
 N4645—W11915/15

1965

AMS 2177 IV—SERIES V791

MAP ACCURACY STANDARDS
 COLORADO 80225, OR WASHINGTON, D. C. 20242
 NO SYMBOLS IS AVAILABLE ON REQUEST

7
 9976





ROAD CLASSIFICATION
 Medium duty ——— Light duty ———
 Unimproved dirt - - - - -
 ○ State Route




SMYRNA, WASH.
 N4645—W11930/15
 1965
 AMS 2077 I—SERIES V791

MAP ACCURACY STANDARDS
 COLORADO 80225, OR WASHINGTON, D. C. 20242
 NO SYMBOLS IS AVAILABLE ON REQUEST



ROAD CLASSIFICATION

- Heavy duty —————
- Medium duty - - - - -
- Light duty ————
- Unimproved dirt - - - - -
- State Route ○



RDS
STON, VIRGINIA 22092
N REQUEST


HANFORD, WASH.
N4630—W11915/15

1965


AMS 2177 III—SERIES V791

TEL: 216

9
9976



ORRAL CANYON 2076 I RICHLAND 21 MI. R. 26 E. 35' 302 303 304 305 306 119° 30'



VERTICAL INTERVAL 20 FEET
MEAN SEA LEVEL DATUM OF 1929



QUADRANGLE LOCATION

ROAD CLASSIFICATION

Heavy-duty	—————	Light-duty	—————
Medium-duty	—————	Unimproved dirt
	○	State Route	

COYOTE RAPIDS, WASH.
N4630—W11930/15

1965


DMA 2077 II—SERIES V791

10TH NATIONAL MAP ACCURACY STANDARDS
DENVER, COLORADO 80225. OR RESTON, VIRGINIA 22092
PHOTOCOPY MAPS AND SYMBOLS IS AVAILABLE ON REQUEST


10
9976

UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

PRIEST RAPIDS QUADRANGLE
WASHINGTON
15 MINUTE SERIES (TOPOGRAPHIC)



Mapped, edited, and published by the Geological Survey in cooperation with the Army Map Service
Control by USGS, USC&GS, and U. S. Bureau of Reclamation
Topography by photogrammetric methods from aerial photographs taken 1948 and planetable surveys by USBR 1935-41. Revised from aerial photographs taken 1964. Field checked 1965
Polyconic projection. 1927 North American datum
10,000-foot grid based on Washington coordinate system, south zone
100-meter Universal Transverse Mercator grid ticks, zone 11, shown in blue
Elevations at section corners are for top of marker
Where omitted, land lines have not been established



ROAD CLASSIFICATION
Medium-duty ——— Light-duty ———
Unimproved dirt - - - - -
State Route
The NW, SW, and SE quarters also covered by 7.5 minute, 1:24,000 scale maps: Priest Rapids 1950, Carrn Hope Peak 1950, and Emerson Nipple 1950

PRIEST RAPIDS, WASH.
N4630-W1945/15
1965
AMS 2077 III-SERIES V791


THIS MAP COMPLIES WITH NATIONAL MAP ACCURACY STANDARDS
FOR SALE BY U. S. GEOLOGICAL SURVEY, DENVER, COLORADO 80225, OR WASHINGTON, D. C. 20242
A FOLDER DESCRIBING TOPOGRAPHIC MAPS AND SYMBOLS IS AVAILABLE ON REQUEST

100° 55' 00" W
100° 50' 00" W
100° 45' 00" W
100° 40' 00" W
100° 35' 00" W
100° 30' 00" W
100° 25' 00" W
100° 20' 00" W
100° 15' 00" W
100° 10' 00" W
100° 05' 00" W
100° 00' 00" W




46° 45' 00" N
46° 40' 00" N
46° 35' 00" N
46° 30' 00" N
46° 25' 00" N
46° 20' 00" N
46° 15' 00" N
46° 10' 00" N
46° 05' 00" N
46° 00' 00" N

UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

CORFU QUADRANGLE
WASHINGTON
15 MINUTE SERIES (TOPOGRAPHIC)




Mapped, edited, and published by the Geological Survey in cooperation with the Army Map Service. Control by USGS, USC&GS, and U.S. Bureau of Reclamation. Topography by photogrammetric methods from aerial photographs taken 1948 and planetable surveys by USBR 1935-1941. Field checked 1950. Revised from aerial photographs taken 1964. Field checked 1965. Polyconic projection. 1927 North American datum. 10,000-foot grid based on Washington coordinate system, south zone. 1000-meter Universal Transverse Mercator grid ticks, zone 11, shown in blue. Elevations at section corners are for top of marker.





THIS MAP COMPLIES WITH NATIONAL MAP ACCURACY STANDARDS FOR SALE BY U.S. GEOLOGICAL SURVEY, DENVER, COLORADO 80225, OR WASHINGTON, D.C. 20242. A FOLDER DESCRIBING TOPOGRAPHIC MAPS AND SYMBOLS IS AVAILABLE ON REQUEST.

CORFU, WASH.
N4645-W11915/15
1965

AMS 2177 IV-SERIES V791




Mapped, edited, and published by the Geological Survey in cooperation with the Army Map Service
 Control by USGS, USC&GS, and U. S. Bureau of Reclamation
 Topography by photogrammetric methods from aerial photographs taken 1949 and planimetric surveys by USBR 1935-41. Field checked 1950. Revised from aerial photographs taken 1964. Field checked 1965
 Polyconic projection, 1927 North American datum
 10,000-foot grid based on Washington coordinate system, south zone
 1000-meter Universal Transverse Mercator grid ticks, zone 11, shown in blue
 Elevations at section corners are for top of marker




THIS MAP COMPLIES WITH NATIONAL MAP ACCURACY STANDARDS
 FOR SALE BY U. S. GEOLOGICAL SURVEY, DENVER, COLORADO 80225, OR WASHINGTON, D. C. 20242
 A FOLDER DESCRIBING TOPOGRAPHIC MAPS AND SYMBOLS IS AVAILABLE ON REQUEST

SMYRNA, WASH.
 N4645—W11930/15
 1965
 AMS 2077 I—SERIES V791



Mapped, edited, and published by the Geological Survey
in cooperation with the Army Map Service
Control by USGS and USC&GS
Topography by photogrammetric methods from aerial
photographs taken 1948 and planetable surveys 1951
Revised from aerial photographs taken 1964
Field checked 1965
Polyconic projection. 1927 North American datum
10,000-foot grid based on Washington coordinate system,
south zone
1000-meter Universal Transverse Mercator grid ticks,
zone 11, shown in blue



CONTOUR INTERVAL 20 FEET
NATIONAL GEODETIC VERTICAL DATUM OF 1929

THIS MAP COMPLIES WITH NATIONAL MAP ACCURACY STANDARDS
FOR SALE BY U.S. GEOLOGICAL SURVEY, DENVER, COLORADO 80225 OR RESTON, VIRGINIA 22092
A FOLDER DESCRIBING TOPOGRAPHIC MAPS AND SYMBOLS IS AVAILABLE ON REQUEST

ROAD CLASSIFICATION
Heavy duty ——— Light duty ———
Medium duty ——— Unimproved dirt - - - - -
○ State Route



HANFORD, WASH.
N4630-W11915/15

1965

AMS 2177 III—SERIES V791

UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

COYOTE RAPIDS QUADRANGLE
WASHINGTON
15 MINUTE SERIES (TOPOGRAPHIC)





60 m
200 E
200 W
200 N

Mapped, edited, and published by the Geological Survey in cooperation with the Army Map Service. Control by USGS and USC&GS.

Topography by photogrammetric methods from aerial photographs taken 1948 and planimetric surveys 1951. Revised from aerial photographs taken 1964. Field checked 1965.

Polyconic projection. 1927 North American datum. 10,000-foot grid based on Washington coordinate system, south zone. 1000-meter Universal Transverse Mercator grid ticks, zone 11, shown in blue. To place on the predicted North American Datum 1983 move the projection lines 19 meters north and 87 meters east.

There may be private inholdings within the boundaries of the National or State reservations shown on this map.



ROAD CLASSIFICATION

Heavy-duty	Light-duty
Medium-duty	Unimproved dirt
	State Route

COYOTE RAPIDS, WASH.
N4630-W11930/15

1965
DMA 2077 II-SERIES V791

THIS MAP COMPLIES WITH NATIONAL MAP ACCURACY STANDARDS FOR SALE BY U.S. GEOLOGICAL SURVEY DENVER, COLORADO 80225, OR RESTON, VIRGINIA 22092. A FOLDER DESCRIBING TOPOGRAPHIC MAPS AND SYMBOLS IS AVAILABLE ON REQUEST.

REFERENCE 9

Waste Management Operations, Volume 1, ERDA-1538

91118901060

TABLE II.1-5a (Continued)

CHARACTERIZATION OF 300 NORTH AND WYE BURIAL GROUNDS

- Field survey of waste disposal sites begun late FY-1975
- Drilling of current series of sampling wells completed late FY-1976
- Estimate of any migration of radionuclides based on available data completed early FY-1977
- Final report and disposal site maps completed and issued late FY-1977

PLUTONIUM RECOVERY

- Detailed schedules will be developed based on the results of the 216-Z-9 Crib plutonium recovery program.

FISSION PRODUCT RECOVERY

- No schedule for recovery has been established.

II.1.1.3 Plant Water Usage

Sanitary and process water for the Hanford Reservation operations are supplied from a number of different locations (Table II.1-6). The 100-B pumphouse, with the 100-D as backup, supplies water to 100-B-C, 100-D, 100-H, 100-F, and 200 Areas. (All of these areas are shown on Figure II.1-1.) A water plant at 100-N Area supplies sanitary and process water to both 100-N and the WPPSS generating station located adjacent to 100-N. The 300 Area pumphouse supplies all water needs for the 300 Area. The City of Richland water supply system is intertied to the 300 Area system as an emergency backup. Various wells in the 600 Areas on the site supply both sanitary and process water.

An average of 440 cfs (0.4% of annual average flow) of Columbia River water is continuously withdrawn, mainly for cooling purposes. Most of this water is discharged directly back to the river as nonradioactive disposal. In addition, approximately 34.7 cfs (less than 0.03% of the total annual average flow) of Columbia River water is withdrawn and discharged to the ground as sanitary and industrial wastes. Wells supply approximately 73,000 gal/day for sanitary and minor irrigation use which is almost entirely discharged to the ground.

II.1.1.4 Waste Inventories [X.18]

Multiple disposal sites for radioactive waste have been used at Hanford during its 30-yr history. The choice of disposal site is based on many factors including half-life and toxicity of the elements involved, quantity of material to be discarded, proximity to water table and/or the Columbia River, and optimum personnel radiation exposure and contamination control. The use of many of these sites has been terminated.

The waste inventory data reported subsequently and in greater detail in Appendix II.1-C for cribs, burial grounds, ponds, ditches, specific retention sites, and unplanned releases are subject to variations in sample collection and analysis.

High-level waste streams are analyzed for plutonium, uranium and neptunium. Fission product content is calculated from irradiation history. The total contents of all waste tanks are known with fair precision, although the contents of individual tanks and, to some extent, of individual tank farms are less certain because of transfers among tanks and farms.

Discharges to cribs, specific retention sites, ponds and ditches are based on line samples. Since the concentration of radionuclides in these streams is often low, accuracy is low and considerable uncertainty is involved. In some cases, particularly for discharges to ponds and ditches, the radioactivity is below limit of detection and the results are therefore reported as "less-than" figures.

Unplanned releases of liquids to grounds are based on estimates of volumes involved and of concentrations of the radionuclides in the stream. Gaseous releases are based on stack samples.

TABLE II.1-6
PLANT WATER USE

<u>Plant or Facility</u>	<u>Source of Supply</u>	<u>Primary Use</u>
100-B	Columbia River 100-B Pumphouse	Process water
100-D	Columbia River 100-D Pumphouse	Process water
100-K	From 100-K Pumphouse	Process water
100-N	Columbia River-100-N Pumphouse	Process, sanitary and cooling water
100-F	From 100-B Pumphouse	Process water
200 Areas	100-B Pumphouse Well 299-E26-6 Wells 299-E28-11 and 15	Process and cooling water Emergency cooling water supply Emergency process supply
300 Area	Columbia River 300 Area Pumphouse	Process and sanitary 331 Fish Ponds
FFTF	Wells 699-S0-7 699-S0-8 699-S1-7B	Construction and Sanitary
Atm. Physics	100-B Pumphouse	Sanitary
609 Fire Sta. (100 Area Central)	100-B Pumphouse	Sanitary
609 A Fire Sta. (200 Area Central)	100-B Area	Sanitary
6652C (Aeronomy)	Rattlesnake Springs	Sanitary
6652I (Ale Hq)	Well 699-26-89	Sanitary and Irrigation
Emergency Relocation Ctr.	Well 699-S18-51	Sanitary
BY Telephone Exchange	Well 699-50-28A	Sanitary
Hanford Road Maintenance	Well 699-Han-9	Industrial and Sanitary
Vernita Park	Wells 699-72-101A, B and C	Sanitary and Irrigation

Reported quantities of radioactivity in burial grounds or storage tunnels are estimates based on operating history and radiation levels of the equipment or waste buried. Generally, these data are of greater uncertainty than liquid and gaseous streams since there is no practical method by which a representative sample, particularly of large equipment pieces, can be obtained.

Overall, however, the accuracy of the inventory total is believed to be within a factor of two, probably within 50% or better. The reported inventories in this final environmental impact statement have been adjusted to a single significant figure to reflect the uncertainties in the data.

II.1.1.4.1 Contained Solids and Liquids

II.1.1.4.1.1 Disposal Sites

The approximate inventories of solid waste disposal in the 100 Areas through 1972 summarized below are shown in Appendix II.1-8, Part 3. These inventories are corrected for decay through the end of 1972.

REFERENCE 10

Inductively Coupled Plasma Method, Method 6010

91118901063

*Chromium Method
of Analysis.*

METHOD 6010

INDUCTIVELY COUPLED PLASMA METHOD

1.0 Scope and Application

1.1 Method 6010 is approved for determining the concentration of trace elements in mobility procedure extracts (exclusive of arsenic and selenium), wastes, and soils.

1.2 This method may be used for the determination of total elements in groundwater if the detection limit is less than primary interim drinking water standards or less than background levels.

1.3 All samples must be subjected to an appropriate dissolution step prior to analysis. (Refer to Section 2.1.)

1.4 Precaution must be taken to ensure that all spectral, chemical and physical interferences are detected and the appropriate corrective measures taken.

1.5 Use of this method is restricted to experienced spectroscopists.

1.6 Elements for which Method 6010 is an approved procedure are:

Aluminum	Cobalt	Potassium
Antimony	Copper	Silica
Barium	Iron	Silver
Beryllium	Lead	Sodium
Boron	Magnesium	Thallium
Cadmium	Manganese	Vanadium
Calcium	Molybdenum	Zinc
<u>Chromium</u>	Nickel	

2.0 Summary of Method

2.1 Prior to analysis by Method 6010, samples must be prepared for sample introduction. The method of sample preparation will vary according to the sample matrix. Aqueous samples are subjected to the acid digestion procedure described in this method. Sludge-type samples are prepared using the procedure described in Method 3050. For samples containing oils, greases or waxes, the procedures described in Methods 3030 and 3040 may be applicable.

2.2 Method 6010 describes a technique for the simultaneous or sequential multielement determination of trace elements in solution. The basis of the method is the measurement of atomic emission by an optical spectroscopic technique. Samples are nebulized and the aerosol that is produced is transported to the plasma torch where excitation occurs. Characteristic atomic-line emission spectra are produced by a radio-frequency inductively coupled plasma

2 / MULTIELEMENT INORGANIC ANALYTICAL TECHNIQUES

(ICP). The spectra are dispersed by a grating spectrometer and the intensities of the lines are monitored by photomultiplier tubes. The photocurrents from the photomultiplier tubes are processed and controlled by a computer system. A background correction technique is required to compensate for variable background contribution to the determination of trace elements. Background must be measured adjacent to analyte lines on samples during analysis. The position selected for the background intensity measurement, on either or both sides of the analytical line, will be determined by the complexity of the spectrum adjacent to the analyte line. The position used must be free of spectral interference and reflect the same change in background intensity as occurs at the analyte wavelength measured. Background correction is not required in cases of line broadening where a background correction measurement would actually degrade the analytical result. The possibility of additional interferences named in 3.1 (and tests for their presence as described in 3.2) should also be recognized and appropriate corrections made.

3.0 Interferences

3.1 Several types of interference effects may contribute to inaccuracies in the determination of trace elements. They can be summarized as follows:

3.1.1 Spectral interferences can be categorized as (1) overlap of a spectral line from another element; (2) unresolved overlap of molecular band spectra; (3) background contribution from continuous or recombination phenomena; and (4) background contribution from stray light from the line emission of high concentration elements. The first of these effects can be compensated by utilizing a computer correction of the raw data, requiring the monitoring and measurement of the interfering element. The second effect may require selection of an alternate wavelength. The third and fourth effects can usually be compensated by a background correction adjacent to the analyte line. In addition, users of simultaneous multielement instrumentation must assume the responsibility of verifying the absence of spectral interference for an element that could occur in a sample but for which there is no channel in the instrument array. Listed in Table 2 are some interference effects for the recommended wavelengths given in Table 1. The data in Table 2 are intended for use only as a rudimentary guide for indicating potential spectral interferences. For this purpose, linear relations between concentration and intensity for the analytes and the interferents can be assumed.

3.1.1.1 The interference information, which was collected at the Ames Laboratory,¹ is expressed as analyte concentration equivalents (i.e., false analyte concentrations) arising from 100 mg/liter of the interference element. The suggested use of this

¹Ames Laboratory, U.S. DOE, Iowa State University, Ames, Iowa 50011.

TABLE 1. RECOMMENDED WAVELENGTHS AND
ESTIMATED INSTRUMENTAL DETECTION LIMITS

Element	Wavelength ^a (nm)	Estimated detection limit ^b (µg/l)
Aluminum	308.215	45
Arsenic	193.696	53 ^c
Antimony	206.833	32
Barium	455.403	2
Beryllium	313.042	0.3
Boron	249.773	5
Cadmium	226.502	4
Calcium	317.933	10
Chromium	267.716	7
Cobalt	228.616	7
Copper	324.754	6
Iron	259.940	7
Lead	220.353	42
Magnesium	279.079	30
Manganese	257.610	2
Molybdenum	202.030	8
Nickel	231.604	15
Potassium	766.491	see ^d
Selenium	196.026	75 ^c
Silica (SiO ₂)	288.158	58
Silver	328.068	7
Sodium	588.995	29
Thallium	190.864	40
Vanadium	292.402	8
Zinc	213.856	2

^aThe wavelengths listed are recommended because of their sensitivity and overall acceptance. Other wavelengths may be substituted if they can provide the needed sensitivity and are treated with the same corrective techniques for spectral interference (see 3.1.1). In time, other elements may be added as more information becomes available and as required.

^bThe estimated instrumental detection limits as shown are taken from "Inductively Coupled Plasma-Atomic Emission Spectroscopy-Prominent Lines," EPA-600/4-79-017. They are given as a guide for an instrumental limit. The actual method detection limits are sample-dependent and may vary as the sample matrix varies.

^cMethod may not be applicable for groundwater analysis for these metals.

^dHighly dependent on operating conditions and plasma position.

TABLE 2. ANALYTE CONCENTRATION EQUIVALENTS
ARISING FROM INTERFERENTS AT THE 100 MG/L LEVEL

Analyte	Wavelength (nm)	Interferent (mg/l)									
		Al	Ca	Cr	Cu	Fe	Hg	Mn	Ni	Tl	V
Aluminum	308.215	--	--	--	--	--	--	0.21	--	--	1.4
Antimony	206.833	0.47	--	2.9	--	0.08	--	--	--	0.25	0.45
Arsenic	193.696	1.3	--	0.44	--	--	--	--	--	--	1.1
Barium	455.403	--	--	--	--	--	--	--	--	--	--
Beryllium	313.042	--	--	--	--	--	--	--	--	0.04	0.05
Boron	249.773	0.04	--	--	--	0.32	--	--	--	--	--
Cadmium	226.502	--	--	--	--	0.03	--	--	0.02	--	--
Calcium	317.933	--	--	0.08	--	0.01	0.01	0.04	--	0.03	0.03
Chromium	267.716	--	--	--	--	0.003	--	0.04	--	--	0.04
Cobalt	228.616	--	--	0.03	--	0.005	--	--	0.03	0.15	--
Copper	324.754	--	--	--	--	0.003	--	--	--	0.05	0.02
Iron	259.940	--	--	--	--	--	--	0.12	--	--	--
Lead	220.353	0.17	--	--	--	--	--	--	--	--	--
Magnesium	279.079	--	0.02	0.11	--	0.13	--	0.25	--	0.07	0.12
Manganese	257.610	0.005	--	0.01	--	0.002	0.002	--	--	--	--
Molybdenum	202.030	0.05	--	--	--	0.03	--	--	--	--	--
Nickel	231.604	--	--	--	--	--	--	--	--	--	--
Selenium	196.026	0.23	--	--	--	0.09	--	--	--	--	--
Silicon	288.158	--	--	0.07	--	--	--	--	--	--	0.01
Sodium	588.995	--	--	--	--	--	--	--	--	0.08	--
Thallium	190.864	0.30	--	--	--	--	--	--	--	--	--
Vanadium	292.402	--	--	0.05	--	0.005	--	--	--	0.02	--
Zinc	213.856	--	--	--	0.14	--	--	--	0.29	--	--

information is as follows: Assume that arsenic (at 193.696 nm) is to be determined in a sample containing approximately 10 mg/liter of aluminum. According to Table 2, 100 mg/liter of aluminum would yield a false signal for arsenic equivalent to approximately 1.3 mg/liter. Therefore, 10 mg/liter of aluminum would result in a false signal for arsenic equivalent to approximately 0.13 mg/liter. The reader is cautioned that other analytical systems may exhibit somewhat different levels of interference than those shown in Table 2, and that the interference effects must be evaluated for each individual system.

3.1.1.2 Only those interferents listed were investigated, and the blank spaces in Table 2 indicate that measurable interferences were not observed for the interferent concentrations listed in Table 3. Generally, interferences were discernible if they produced peaks or background shifts corresponding to 2-5% of the peaks generated by the analyte concentrations also listed in Table 3.

3.1.1.3 At present, information on the listed silver and potassium wavelengths is not available but it has been reported that second order energy from the magnesium 383.231-nm wavelength interferes with the listed potassium line at 766.491 nm.

3.1.2 Physical interferences are generally considered to be effects associated with the sample nebulization and transport processes. Such properties as change in viscosity and surface tension can cause significant inaccuracies especially in samples which may contain high dissolved solids or acid concentrations. The use of a peristaltic pump may lessen these interferences. If these types of interferences are operative, they must be reduced by diluting the sample and/or utilizing standard addition techniques. Another problem which can occur from high dissolved solids is salt buildup at the tip of the nebulizer. This affects aerosol flow rate, causing instrumental drift. Wetting the argon prior to nebulization, the use of a tip washer, or sample dilution have been used to control this problem. Also, the use of a high solid nebulizer can reduce salt build-up in the nebulizer and can also prevent drifting and loss of sensitivity in the instrument. In addition, it has been reported that better control of the argon flow rate improves instrument performance. This is accomplished with the use of mass flow controllers.

3.1.3 Chemical interferences are characterized by molecular compound formation, ionization effects and solute vaporization effects. Normally these effects are not pronounced with the ICP technique; however, if observed they can be minimized by careful selection of operating conditions (that is, incident power, observation position, and so forth), by buffering of the sample, by matrix matching, and by standard addition procedures. These types of interferences can be highly dependent on matrix type and the specific analyte element.

6 / MULTIELEMENT INORGANIC ANALYTICAL TECHNIQUES

TABLE 3. INTERFERENT AND ANALYTE ELEMENTAL CONCENTRATIONS
USED FOR INTERFERENCE MEASUREMENTS IN TABLE 2

Analytes	mg/liter	Interferents	mg/liter
Al	10	Al	1,000
As	10	Ca	1,000
B	10	Cr	200
Ba	1	Cu	200
Be	1	Fe	1,000
Ca	1	Mg	1,000
Cd	10	Mn	200
Co	1	Ni	200
Cr	1	Ti	200
Cu	1	V	200
Fe	1		
Mg	1		
Mn	1		
Mo	10		
Na	10		
Ni	10		
Pb	10		
Sb	10		
Se	10		
Si	1		
Ti	10		
V	1		
Zn	10		

3.2 It is recommended that whenever a new or unusual sample matrix is encountered, a series of tests be performed prior to reporting concentration data for analyte elements. These tests, as outlined in 3.2.1 through 3.2.4, will ensure the analyst that neither positive nor negative interferences will have on any of the analyte elements and distort the accuracy of the reported values.

3.2.1 Serial dilution: If the analyte concentration is sufficiently high (minimally a factor of 10 above the instrumental detection limit after dilution), an analysis of a dilution should agree within 5 percent of the original determination (or within some acceptable control limit that has been established for that matrix). If not, a chemical or physical interference effect should be suspected.

3.2.2 Spike addition: A spike addition added at a minimum level of 10x the instrumental detection limit (maximum 100x) to the original determination should be recovered to within 90 to 110 percent or within the established control limit for that matrix. If not, a matrix effect should be suspected. The use of a standard addition analysis procedure can usually compensate for this effect. CAUTION: The standard addition technique does not detect coincident spectral overlap. If suspected, use of computerized compensation, an alternate wavelength, or comparison with an alternate method is recommended (see 3.2.3).

3.2.3 Comparison with alternate method of analysis: When investigating a new sample matrix, comparison tests may be performed with other analytical techniques such as atomic absorption spectrometry or other approved methodology.

3.2.4 Wavelength scanning of analyte line region: If the appropriate equipment is available, wavelength scanning can be performed to detect potential spectral interferences.

4.0 Apparatus and Materials

4.1 Inductively coupled plasma-atomic emission spectrometer

4.1.1 Computer-controlled atomic emission spectrometer with background correction.

4.1.2 Radiofrequency generator.

4.1.3 Argon gas supply, welding grade or better.

8 / MULTIELEMENT INORGANIC ANALYTICAL TECHNIQUES

4.2 Operating conditions: Because of the differences between various makes and models of satisfactory instruments, no detailed operating instructions can be provided. Instead, the analyst should follow the instructions provided by the manufacturer of the particular instrument. Sensitivity, instrumental detection limit, precision, linear dynamic range, and interference effects must be investigated and established for each individual analyte line on that particular instrument. It is the responsibility of the analyst (1) to verify that the instrument configuration and operating conditions used satisfy the analytical requirements and (2) to maintain quality control data confirming instrument performance and analytical results.

5.0 Reagents

5.1 Acids used in the preparation of standards and for sample processing must be ultra-high purity grade or equivalent. Redistilled acids are acceptable.

5.1.1 Acetic acid, conc.

5.1.2 Hydrochloric acid, conc.

5.1.3 Hydrochloric acid (1:1): Add 500 ml conc. HCl to 400 ml Type II water and dilute to 1 liter.

5.1.4 Nitric acid, conc.

5.1.5 Nitric acid (1:1): Add 500 ml conc. HNO₃ to 400 ml Type II water and dilute to 1 liter.

5.2 Type II water: Prepare by passing distilled water through a mixed bed of cation and anion exchange resins. Use Type II water for the preparation of all reagents, calibration standards, and as dilution water. The purity of this water must be equivalent to or better than ASTM Type II reagent water of Specification D1193.

5.3 Standard stock solutions may be purchased or prepared from ultra-high purity grade chemicals or metals. All salts must be dried for 1 hr at 105° C unless otherwise specified. (CAUTION: Many metal salts are extremely toxic and may be fatal if swallowed. Wash hands thoroughly after handling.) Typical stock solution preparation procedures follow.

5.3.1 Aluminum solution, stock, 1 ml=100 µg Al: Dissolve 0.100 g of aluminum metal in an acid mixture of 4 ml of (1:1) HCl and 1 ml of conc. HNO₃ in a beaker. Warm gently to dissolve. When solution is complete, transfer quantitatively to a one liter flask and add an additional 10 ml of (1:1) HCl and dilute to 1,000 ml with Type II water.

5.3.2 Antimony solution, stock, 1 ml=100 µg Sb: Dissolve 0.2669 g $K(SbO)C_4H_4O_6$ in Type II water, add 10 ml (1:1) HCl and dilute to 1,000 ml with Type II water.

5.3.3 Arsenic solution, stock, 1 ml=100 µg As: Dissolve 0.1320 g of As_2O_3 in 100 ml of Type II water containing 0.4 g NaOH. Acidify the solution with 2 ml conc. HNO_3 and dilute to 1,000 ml with Type II water.

5.3.4 Barium solution, stock, 1 ml=100 µg Ba: Dissolve 0.1516 g $BaCl_2$ (dried at 250° C for 2 hr) in 10 ml Type II water with 1 ml (1:1) HCl. Add 10.0 ml (1:1) HCl and dilute to 1,000 ml with Type II water.

5.3.5 Beryllium solution, stock, 1 ml=100 µg Be: Do not dry. Dissolve 1.966 g $BeSO_4 \cdot 4H_2O$ in Type II water, add 10.0 ml conc. HNO_3 and dilute to 1,000 ml with Type II water.

5.3.6 Boron solution, stock, 1 ml=100 µg B: Do not dry. Dissolve 0.5716 g anhydrous H_3BO_3 in Type II water and dilute to 1,000 ml. Keep in a tightly stoppered bottle, and store in a desiccator to prevent the entrance of atmospheric moisture.

5.3.7 Cadmium solution, stock, 1 ml=100 µg Cd: Dissolve 0.1142 g CdO in a minimum amount of (1:1) HNO_3 . Heat to increase rate of dissolution. Add 10.0 ml conc. HNO_3 and dilute to 1,000 ml with Type II water.

5.3.8 Calcium solution, stock, 1 ml=100 µg Ca: Suspend 0.2498 g $CaCO_3$ dried at 180° C for 1 hr before weighing in Type II water and dissolve cautiously with a minimum amount of (1:1) HNO_3 . Add 10.0 ml conc. HNO_3 and dilute to 1,000 ml with Type II water.

5.3.9 Chromium solution, stock, 1 ml=100 µg Cr: Dissolve 0.1923 g CrO_3 in Type II water. When solution is complete, acidify with 10 ml conc. HNO_3 and dilute to 1,000 ml with Type II water.

5.3.10 Cobalt solution, stock, 1 ml=100 µg Co: Dissolve 0.1000 g of cobalt metal in a minimum amount of (1:1) HNO_3 . Add 10.0 ml (1:1) HCl and dilute to 1,000 ml with Type II water.

5.3.11 Copper solution, stock, 1 ml = 100 µg Cu: Dissolve 0.1252 g CuO in a minimum amount of (1+1) HNO_3 . Add 10.0 ml conc. HNO_3 and dilute to 1,000 ml with deionized, distilled water.

5.3.12 Iron solution, stock, 1 ml=100 µg Fe: Dissolve 0.1430 g Fe_2O_3 in a warm mixture of 20 ml (1:1) HCl and 2 ml of conc. HNO_3 . Cool, add an additional 5.0 ml of conc. HNO_3 and dilute to 1,000 ml with Type II water.

10 / MULTIELEMENT INORGANIC ANALYTICAL TECHNIQUES

5.3.13 Lead solution, stock, 1 ml=100 µg Pb: Dissolve 0.1599 g $\text{Pb}(\text{NO}_3)_2$ in a minimum amount of (1:1) HNO_3 . Add 10.0 ml (1:1) HNO_3 and dilute to 1,000 ml with Type II water.

5.3.14 Magnesium solution, stock, 1 ml=100 µg Mg: Dissolve 0.1658 g MgO in a minimum amount of (1:1) HNO_3 . Add 10.0 ml (1:1) conc. HNO_3 and dilute to 1,000 ml with Type II water.

5.3.15 Manganese solution, stock, 1 ml=100 µg Mn: Dissolve 0.1000 g of manganese metal in the acid mixture (10 ml conc. HCl and 1 ml conc. HNO_3) and dilute to 1,000 ml with Type II water.

5.3.16 Molybdenum solution, stock, 1 ml=100 µg Mo: Dissolve 0.2043 g $(\text{NH}_4)_2\text{MoO}_4$ in Type II water and dilute to 1,000 ml with Type II water.

5.3.17 Nickel solution, stock, 1 ml=100 µg Ni: Dissolve 0.1000 g of nickel metal in 10.0 ml hot conc. HNO_3 , cool, and dilute to 1,000 ml with Type II water.

5.3.18 Potassium solution, stock, 1 ml=100 µg K: Dissolve 0.1907 g KCl , dried at 110° C, in Type II water and dilute to 1,000 ml.

5.3.19 Selenium solution, stock, 1 ml=100 µg Se: Do not dry. Dissolve 0.1727 g H_2SeO_3 in Type II water and dilute to 1,000 ml.

5.3.20 Silica solution, stock, 1 ml=100 µg SiO_2 : Do not dry. Dissolve 0.4730 g $\text{Na}_2\text{SiO}_3 \cdot 9\text{H}_2\text{O}$ in Type II water. Add 10.0 ml conc. HNO_3 and dilute to 1,000 ml with Type II water.

5.3.21 Silver solution, stock, 1 ml=100 µg Ag: Dissolve 0.1575 g AgNO_3 in Type II water and 10 ml conc. HNO_3 . Dilute to 1,000 ml with Type II water.

5.3.22 Sodium solution, stock, 1 ml=100 µg Na: Dissolve 0.2542 g NaCl in Type II water. Add 10.0 ml conc. HNO_3 and dilute to 1,000 ml with Type II water.

5.3.23 Thallium solution, stock, 1 ml=100 µg Tl: Dissolve 0.1303 g TlNO_3 in Type II water. Add 10.0 ml conc. HNO_3 and dilute to 1,000 ml with Type II water.

5.3.24 Vanadium solution, stock, 1 ml=100 µg V: Dissolve 0.2297 g NH_4VO_3 in a minimum amount of conc. HNO_3 . Heat to increase rate of dissolution. Add 10.0 ml conc. HNO_3 and dilute to 1,000 ml with Type II water.

5.3.25 Zinc solution, stock, 1 ml=100 µg Zn: Dissolve 0.1245 g ZnO in a minimum amount of dilute HNO_3 . Add 10.0 ml conc. HNO_3 and dilute to 1,000 ml with Type II water.

5.4 Mixed calibration standard solutions: Prepare mixed calibration standard solutions by combining appropriate volumes of the stock solutions in volumetric flasks (see 5.4.1 through 5.4.5). Add 2 ml of (1:1) HNO_3 and 10 ml of (1:1) HCl and dilute to 100 ml with Type II water (see Note 1 in Section 5.4.5 and Note 6 in Section 7.3). Prior to preparing the mixed standards, each stock solution should be analyzed separately to determine possible spectral interference or the presence of impurities. Care should be taken when preparing the mixed standards that the elements be compatible and stable. Transfer the mixed standard solutions to a FEP fluorocarbon or unused polyethylene bottle for storage. Fresh mixed standards should be prepared as needed with the realization that concentration can change on aging. Calibration standards must be initially verified using a quality control sample and monitored weekly for stability (see 5.6.3). Although not specifically required, some typical calibration standard combinations follow when using those specific wavelengths listed in Table 1.

5.4.1 Mixed standard solution I: Manganese, beryllium, cadmium, lead, and zinc.

5.4.2 Mixed standard solution II: Barium, copper, iron, vanadium, and cobalt.

5.4.3 Mixed standard solution III: Molybdenum, silica, arsenic, and selenium.

5.4.4 Mixed standard solution IV: Calcium, sodium, potassium, aluminum, chromium, and nickel.

5.4.5 Mixed standard solution V: Antimony, boron, magnesium, silver, and thallium.

NOTE 1: If the addition of silver to the recommended acid combination results in an initial precipitation, add 15 ml of Type II water and warm the flask until the solution clears. Cool and dilute to 100 ml with Type II water. For this acid combination, the silver concentration should be limited to 2 mg/l. Silver under these conditions is stable in a tap water matrix for 30 days. Higher concentrations of silver require additional HCl .

5.5 Two types of blanks are required for the analysis. The calibration blank is used in establishing the analytical curve while the reagent blank is used to correct for possible contamination resulting from varying amounts of the acids used in the sample processing.

5.5.1 The calibration blank is prepared by diluting 2 ml of (1:1) HNO_3 and 10 ml of (1:1) HCl to 100 ml with Type II water (see note 6 in Section 7.3). Prepare a sufficient quantity to be used to flush the system between standards and samples.

12 / MULTIELEMENT INORGANIC ANALYTICAL TECHNIQUES

5.5.2 The reagent blank must contain all the reagents and in the same volumes as used in the processing of the samples. The reagent blank must be carried through the complete procedure and contain the same acid concentration in the final solution as the sample solution used for analysis.

5.6 Standards: In addition to the calibration standards, an instrument check standard, an interference check sample, and a quality control sample are also required for the analyses.

5.6.1 The instrument check standard is prepared by the analyst by combining compatible elements at a concentration equivalent to the midpoint of their respective calibration curves (see 8.5.1).

5.6.2 The interference check is prepared by the analyst in the following manner. Select a representative sample that contains minimal concentrations of the analytes of interest but known concentrations of interfering elements that will provide an adequate test of the correction factors. Spike the sample with the elements of interest at the approximate concentration of either 100 µg/liter or 5 times the estimated detection limits given in Table 1. (For effluent samples of expected high concentrations, spike at an appropriate level.) If the type of samples analyzed are varied, a synthetically prepared sample may be used if the above criteria and intent are met. A limited supply of a synthetic interference check sample will be available from the Quality Assurance Branch of EMSL-Cincinnati.

5.6.3 The quality control sample should be prepared in the same acid matrix as the calibration standards at a concentration near 1 mg/liter and in accordance with the instructions provided by the supplier.

6.0 Sample Collection, Handling, and Preservation

6.1 All samples must be collected using a sampling plan that addresses the considerations discussed in Section One of this manual.

6.2 For the determination of trace elements, contamination and loss are of prime concern. Dust in the laboratory environment, impurities in reagents, and impurities on laboratory apparatus which the sample contacts are all sources of potential contamination. Samples containers can introduce either positive or negative errors in the measurement of trace elements by (a) contributing contaminants through leaching or surface desorption and (b) by depleting concentrations through adsorption. Thus the collection and treatment of the sample prior to analysis requires particular attention. Laboratory glassware including the sample bottle (whether polyethylene, polypropylene, or FEP-fluorocarbon) should be thoroughly washed with detergent and tap water; and rinsed with (1:1) nitric acid, tap water, (1:1) hydrochloric acid, tap water, and finally Type II water, in that order (see Notes 2 and 3).

NOTE 2: Chromic acid may be useful to remove organic deposits from glassware; however, the analyst should be cautioned that the glassware must be thoroughly rinsed with water to remove the last traces of chromium. The use of chromic acid can cause a contamination problem for the determination of chromium if the glassware is not rinsed properly. A commercial product, NOCHROMIX, available from Godax Laboratories, 6 Varick Street, New York, NY 10013, may be used in place of chromic acid. Chromic acid should not be used with plastic bottles.

NOTE 3: If it can be documented through an active analytical quality control program using spiked samples and reagent blanks that certain steps in the cleaning procedure are not required for routine samples, those steps may be eliminated from the procedure.

6.3 Aqueous samples must be preserved and pretreated appropriately depending on whether data on dissolved, suspended, or total metals are desired.

6.4 Nonaqueous samples shall be refrigerated when possible, and analyzed as soon as possible.

7.0 Procedure

7.1 The following definitions are intended to clarify the ensuing discussion.

7.1.1 Instrumental detection limit: The concentration equivalent to a signal, due to the analyte, which is equal to three times the standard deviation of a series of twenty replicate measurements of a reagent blank signal at the same wavelength.

7.1.2 Sensitivity: The slope of the analytical curve, i.e., functional relationship between emission intensity and concentration.

7.1.3 Instrument check standard: A multielement standard of known concentration prepared by the analyst to monitor and verify instrument performance on a daily basis (see 5.6.1).

7.1.4 Interference check sample: A solution containing both interfering and analyte elements of known concentration that can be used to verify background and interelement correction factors (see 5.6.2).

7.1.5 Quality control sample: A solution obtained from an outside source having known concentration values to be used to verify the calibration standards (see 5.5.3).

14 / MULTIELEMENT INORGANIC ANALYTICAL TECHNIQUES

7.1.6 Calibration standards: A series of known standard solutions used by the analyst for calibration of the instrument (i.e., preparation of the analytical curve; see 5.4).

7.1.7 Linear dynamic range: The concentration range over which the analytical curve remains linear.

7.1.8 Reagent blank: A volume of Type II water containing the same acid matrix as the calibration standards carried through the entire analytical scheme (see 5.5.2).

7.1.9 Calibration blank: A volume of Type II water acidified with HNO_3 and HCl (see 5.5.1).

7.1.10 Method of standard addition: The standard addition technique involves the use of the unknown and unknown plus a known amount of standard (see 7.9.1).

7.2 Sample preparation: Aqueous samples should be prepared according to Section 7.3. Sludge-type samples should be prepared according to Method 3050 and samples containing oils, greases, or waxes may be prepared according to Methods 3030 and 3040. The applicability of a sample preparative technique to a new matrix type must be demonstrated by analyzing spiked sample or relevant standard reference materials.

7.3 Preparation of aqueous samples: For the determination of total elements, choose a measured volume of the well-mixed acid-preserved sample appropriate for the expected level of elements and transfer to a Griffin beaker (see note 5 below). Add 3 ml conc. HNO_3 . Place the beaker on a hot plate and evaporate to near dryness cautiously, making certain that the sample does not boil and that no area of the bottom of the beaker is allowed to go dry. Cool the beaker and add another 5-ml portion of conc. HNO_3 . Cover the beaker with a watch glass and return to the hot plate. Increase the temperature of the hot plate so that a gentle reflux action occurs. Continue heating, adding acid as necessary, until the digestion is complete (generally indicated when the digestate is light in color or does not change in appearance with continued refluxing). Again, evaporate to near dryness and cool the beaker. Add 10 ml of (1:1) HCl and 15 ml of Type II water per 100 ml final solution and warm the beaker gently for 15 min to dissolve any precipitate or residue resulting from evaporation. Allow to cool, wash down the beaker walls and watch glass with Type II water and filter the sample to remove insoluble material that could clog the nebulizer (see Note 4 below). Adjust the sample to a predetermined volume based on the expected concentrations of elements present. The sample is now ready for analysis (see Note 6 below). Concentrations so determined shall be reported as "total."

NOTE 4: In place of filtering, after diluting and mixing, the sample may be centrifuged or allowed to settle by gravity overnight to remove insoluble material.

NOTE 5: If low determinations of boron are critical, quartz or polypropylene glassware should be used. Polypropylene glassware is generally preferred since it is cheaper and just as accurate as quartz.

NOTE 6: If the sample analysis solution has a different acid concentration from that given in 7.3, but does not introduce a physical interference or affect the analytical result, the same calibration standards may be used.

7.4 Set up instrument with proper operating parameters established in Section 4.2. The instrument must be allowed to become thermally stable before beginning. This usually requires at least 30 min of operation prior to calibration.

7.5 Initiate appropriate operating configuration of computer.

7.6 Profile and calibrate instrument according to instrument manufacturer's recommended procedures, using the typical mixed calibration standard solutions described in Section 5.4. Flush the system with the calibration blank (5.5.1) between each standard (see Note 7 below). (The use of the average intensity of multiple exposures for both standardization and sample analysis has been found to reduce random error.)

NOTE 7: For boron concentrations greater than 500 $\mu\text{g/liter}$ extended flush times of 1 to 2 min may be required.

7.7 Before beginning the sample run, reanalyze the highest mixed calibration standard as if it were a sample. Concentration values obtained should not deviate from the actual values by more than +5% (or the established control limits, whichever is lower). If they do, follow the recommendations to the instrument manufacturer to correct for this condition.

7.8 Begin the sample run by flushing the system with the calibration blank solution (5.5.1) between each sample (see Note 7). Analyze the instrument check standard (5.6.1) and the calibration blank (5.5.1) after each 10 samples.

7.9 If it has been found that methods of standard addition are required, the following procedure is recommended.

7.9.1 The standard addition technique involves preparing new standards in the sample matrix by adding known amounts of standard to one or more aliquots of the processed sample solution. This technique compensates for a sample constituent that enhances or depresses the analyte signal, thus producing a different slope from that of the calibration standards. It will not correct for additive interference which causes a baseline shift. The simplest version of this technique is the single-addition method. The procedure is as follows. Two identical aliquots of the sample solution, each of volume V_x , are taken. To the first (labeled A) is added a small volume V_s of a standard analyte

16 / MULTIELEMENT INORGANIC ANALYTICAL TECHNIQUES

solution of concentration c_s . To the second (labeled B) is added the same volume V_s of the solvent. The analytical signals of A and B are measured and corrected for nonanalyte signals. The unknown sample concentration c_x is calculated:

$$c_x = \frac{S_B V_s c_s}{(S_A - S_B) V_x}$$

where S_A and S_B are the analytical signals (corrected for the blank) of solutions A and B, respectively. V_s and c_s should be chosen so that S_A is roughly twice S_B on the average. It is best if V_s is made much less than V_x , and thus c_s is much greater than c_x , to avoid excess dilution of the sample matrix. If a separation or concentration step is used, the additions are best made first and carried through the entire procedure. For the results from this technique to be valid, the following limitations must be taken into consideration.

1. The analytical response signals must be linear with respect to concentration.
2. The chemical form of the analyte added must respond the same as the analyte in the sample.
3. The interference effect must be constant over the working range of concern.
4. The signal must be corrected for any additive interference.

7.10 Calculations: Reagent blanks (5.5.2) should be subtracted from all samples. This is particularly important for digested samples requiring large quantities of acids to complete the digestion. If dilutions were performed, the appropriate factor must be applied to sample values. All results should be reported in mg/liter with up to three significant figures.

8.0 Quality Control

8.1 All quality control data should be maintained and available for easy reference or inspection.

8.2 Dilute samples if they are more concentrated than the highest standard or if they fall on the plateau of a calibration curve.

8.3 Employ a minimum of one procedural blank per sample batch to determine if contamination or any memory effects are occurring.

8.4 Run one duplicate sample for every 10 samples. A duplicate sample is a sample brought through the whole sample preparation process.

8.5 Check the instrument standardization by analyzing appropriate quality control check standards as follows:

8.5.1 Analyze an appropriate instrument check standard (5.6.1) containing the elements of interest at a frequency of 10%. This check standard is used to determine instrument drift. If agreement is not within $\pm 5\%$ of the expected values or within the established control limits, whichever is lower, the analysis is out of control. The analysis should be terminated, the problem corrected, and the instrument recalibrated.

8.5.2 Analyze the calibration blank (5.5.1) at a frequency of 10%. The result should be within the established control limits of 2 standard deviations of the mean value. If not, repeat the analysis two more times and average the three results. If the average is not within the control limit, terminate the analysis, correct the problem, and recalibrate the instrument.

8.5.3 To verify interelement and background correction factors, analyze the interference check sample (5.6.2) at the beginning, end, and at periodic intervals throughout the sample run. Results should fall within the established control limits of 1.5 times the standard deviation of the mean value. If not, terminate the analysis, correct the problem, and recalibrate the instrument.

8.5.4 A quality control sample (5.6.3) obtained from an outside source must first be used for the initial verification of the calibration standards. A fresh dilution of this sample shall be analyzed every week thereafter to monitor their stability. If the results are not within $\pm 5\%$ of the true value listed for the control sample, prepare a new calibration standard and recalibrate the instrument. If this does not correct the problem, prepare a new stock standard and a new calibration standard and repeat the calibration.

8.6 Spiked samples or standard reference materials shall be employed periodically to ensure that correct procedures are being followed and that all equipment is operating properly.

8.7 The method of standard additions shall be used for the analysis of all EP extracts and whenever a sample suffers from matrix interferences.

8.8 The method detection limit (MDL) is defined as the minimum concentration of a substance that can be measured and reported with 99% confidence that the value is above zero. The MDL concentrations listed in Table 1 were obtained using reagent water. Similar results were achieved using representative wastewaters. The MDL actually achieved in a given analysis will vary depending on instrument sensitivity and matrix effects.

8.9 In an EPA round-robin phase I study, seven laboratories applied the ICP technique to acid-distilled water matrices that had been dosed with various metal concentrates. Table 4 lists the true value, the mean reported value and the mean % relative standard deviation.

REFERENCE 11

Hanford Stream data base printout showing Priest Rapids Dam data
as background data

1
0
1
0
6
3
1
1
1
6

ENVIRONMENTAL SAMPLE ANALYSIS RESULT REPORT

 21 DEC 82 to 15 MAR 83

WATER RIVER WATER CUMULATIVE UNFILTERED
 Samp #: 1265 PRIEST RAPIDS-RIVER

121/SR 90

Use this sample for background

Date Off	Date On	Result +/- 2 Sigma (PCI/L)	Comments
-----	-----	-----	-----
21 DEC 82	23 NOV 82	3.37E-01 +/- 1.83E-01	
18 JAN 83	21 DEC 82	1.89E-01 +/- 2.44E-02	<i>BKG</i>
15 FEB 83	18 JAN 83	1.91E-01 +/- 3.18E-02	
15 MAR 83	15 FEB 83	2.57E-01 +/- 4.11E-02	

Fraction of Results >DL: 4/ 4 Mean: 2.44E-01
 Minimum: 1.89E-01 (18 JAN 83) Standard Error of Mean: 8.64E-02
 Maximum: 3.37E-01 (21 DEC 82) Standard Deviation: 1.73E-01
 Median: 1.91E-01

REFERENCE 12

Investigation of Ground-Water Seepage from Hanford Shoreline
of the Columbia River, PNL-5289, November 1984

91118901003

Investigation of Ground- Water Seepage from the Hanford Shoreline of the Columbia River

**W. D. McCormack
J. M. V. Carfile**

November 1984

**Prepared for the U.S. Department of Energy
under Contract DE-AC06-76RLO 1830**

**Pacific Northwest Laboratory
Operated for the U.S. Department of Energy
by Battelle Memorial Institute**






FIGURE 3. Locations and Analytical Results for Spring and River Samples from River Mile 3 through River Mile 12

in PNL-5289

TABLE B.1. Strontium-90 Analyses from Columbia River Samples

River Mile Location	Sample ID	Date Collected	Concentration, pCi/l $\pm 2\sigma$
3.0-5.0	B Comp RW ^(a)	01/22/83	0.55 \pm 0.23
5.5-7.5	K Comp RW	12/18/82	0.18 \pm 0.02
8.0-9.5	N Comp RW	12/18/82	28 \pm 0.47
10.0-12.0	D Comp RW	12/18/82	1.1 \pm 0.05
14.0-17.5	H Comp RW	01/22/83	0.50 \pm 0.14
18.0-22.0	F Comp RW	01/22/83	0.93 \pm 0.15
Upstream Columbia River Concentration (Average 1983)			0.18 \pm 0.22
DOE Concentration Guide (USDOE 1981)			300

DG
use this sample for down gradient
DG
because of close sampling dist. and order magnitude higher difference

(a) Comp-RW denotes composite river water sample comprised of aliquots from immediately preceding river sample locations.

TABLE B.2. Iodine-129 Analyses from Spring and Columbia River Samples

River Mile Location	Sample ID	Date Collected	Concentration, pCi/l $\pm 2\sigma$
27.0	27.0 RW ^(a)	01/22/83	$3.3 \times 10^{-6} \pm 1.4 \times 10^{-6}$
27.0	27-1 Sp ^(b)	09/11/83	$1.6 \times 10^{-4} \pm 2.1 \times 10^{-5}$
28.0	28-2 Sp	09/11/83	$6.2 \times 10^{-2} \pm 6.8 \times 10^{-3}$
29.0	29.0 RW	01/22/83	$6.3 \times 10^{-5} \pm 5.0 \times 10^{-6}$
31.75	31-5 Sp	09/11/83	$3.0 \times 10^{-5} \pm 4.0 \times 10^{-6}$
32.5	32-0 Sp	09/11/83	$4.4 \times 10^{-5} \pm 2.7 \times 10^{-5}$
Upstream Columbia River Concentration (1983 Average)			$2.4 \times 10^{-5} \pm 2.6 \times 10^{-5}$
DOE Concentration Guide (USDOE 1981)			60

(a) RW denotes composite river water sample.
 (b) Sp denotes river bank spring sample.

(in PNL-5289)

REFERENCE 13

Hanford Reservation Area Worker Census, BNWL-2298, July 1977

9 1 1 1 3 2 2 1 0 9 7

91118901088

Hanford Reservation
Area Worker Census

A Research Report for the
Rockwell Hanford Operation

Rockwell International Corporation
Hanford, California



TABLE 1. Concentration of Hanford Reservation Workers by Site

<u>Site</u>	<u>Number of Workers</u>	<u>Percent of Total</u>
100	760	5
200 E&W	2,355	16
WPPSS #1,2,&4	2,905	20
FFTF	2,420	16
300	3,110	21
Battelle, et al.	3,345	22
TOTAL	<u>14,895</u>	<u>100</u>

While the worker counts being reported by Reservation employers are usually shown concentrated around a designated site, in reality a substantial portion of these workers are likely to be distributed over the surrounding area. For convenience, however, they are credited to such particular sites as 200 East, 200 West, WPPSS 1, 2, & 4, etc.

Identification of shift workers posed some reporting difficulties since some firms run four shifts while most of the others conduct their operations in three. The 100 Area was a special problem since these workers operate over a wide area. Regardless, all workers have been accounted for in this census although some of the shift counts may be approximate.

DISTRIBUTION OF WORKERS BY RADII AND COMPASS DIRECTION

Figure 2 maps the distribution of Hanford Reservation workers by work shift over intervals of one-mile radii and 16 compass directions centered at the Purex Plant. These same worker distributions are repeated in Figure 3 without the mapped Reservation Area as a background. As a tabulating convenience, sector parcel counts have been rounded to units of 5 and 10, but were adjusted to the total count for the separate companies. (Because of confidentiality, worker counts for the separate companies are not being presented here.) For better readability, sector counts within the first two mile radii from the Purex Plant center are presented separately at the bottom of the figure. Table 2 presents work distribution in detail including a cumulative count of workers and percent of total as distance and direction from the Purex Plant center increases.

REFERENCE 14

Endangered and Threatened Wildlife and Plants, 50 CFR,
Part 17, Subpart B, October 86

91118901090

issued under Part 21 of this chapter, and

(2) Identified in the earliest applicable annual report required to be filed by a permittee under Part 21 of this chapter as in a permittee's possession on November 10, 1978, or as the progeny of such a raptor.

(b) This section does not apply to any raptor intentionally returned to the wild.

[48 FR 31607, July 8, 1983]

Subpart B—Lists

SOURCE: 48 FR 34182, July 27, 1983, unless otherwise noted.

§ 17.11 Endangered and threatened wildlife.

(a) The list in this section contains the names of all species of wildlife which have been determined by the Services to be Endangered or Threatened. It also contains the names of species of wildlife treated as Endangered or Threatened because they are sufficiently similar in appearance to Endangered or Threatened species (see § 17.50 *et seq.*).

(b) The columns entitled "Common Name," "Scientific Name," and "Vertebrate Population Where Endangered or Threatened" define the species of wildlife within the meaning of the Act. Thus, differently classified geographic populations of the same vertebrate subspecies or species shall be identified by their differing geographic boundaries, even though the other two columns are identical. The term "Entire" means that all populations throughout the present range of a vertebrate species are listed. Although common names are included, they cannot be relied upon for identification of any specimen, since they may vary greatly in local usage. The Services shall use the most recently accepted scientific name. In cases in which confusion might arise, a synonym(s) will be provided in parentheses. The Services shall rely to the extent practicable on the *International Code of Zoological Nomenclature*.

(c) In the "Status" column the following symbols are used: "E" for Endangered, "T" for Threatened, and "E

[or T] (S/A)" for similarity of appearance species.

(d) The other data in the list are nonregulatory in nature and are provided for the information of the reader. In the annual revision and compilation of this title, the following information may be amended without public notice: the spelling of species' names, historical range, footnotes, references to certain other applicable portions of this title, synonyms, and more current names. In any of these revised entries, neither the species, as defined in paragraph (b) of this section, nor its status may be changed without following the procedures of Part 424 of this title.

(e) The "historic range" indicates the known general distribution of the species or subspecies as reported in the current scientific literature. The present distribution may be greatly reduced from this historic range. This column does not imply any limitation on the application of the prohibitions in the Act or implementing rules. Such prohibitions apply to all individuals of the species, wherever found.

(f)(1) A footnote to the FEDERAL REGISTER publication(s) listing or reclassifying a species is indicated under the column "When listed." Footnote numbers to §§ 17.11 and 17.12 are in the same numerical sequence, since plants and animals may be listed in the same FEDERAL REGISTER document. That document, at least since 1973, includes a statement indicating the basis for the listing, as well as the effective date(s) of said listing.

(2) The "Special rules" and "Critical habitat" columns provide a cross reference to other sections in Parts 17, 222, 226, or 227. The "Special rules" column will also be used to cite the special rules that describe experimental populations and determine if they are essential or nonessential. Separate listing will be made for experimental populations, and the status column will include the following symbols: "XE" for an essential experimental population and "XN" for a nonessential experimental population. The term "NA" (not applicable) appearing in either of these two columns indicates that there are no special rules and/or critical habitat for that par-

ticular species appropriate through 2: that species other rules such wildlife requirements references column list two Service species or t Federal age erments.

(g) The list includes all example, t2 (bons) is list

ticular species. However, all other appropriate rules in Parts 17, 217 through 227, and 402 still apply to that species. In addition, there may be other rules in this title that relate to such wildlife, e.g., port-of-entry requirements. It is not intended that the references in the "Special rules" column list all the regulations of the two Services which might apply to the species or to the regulations of other Federal agencies or State or local governments.

(g) The listing of a particular taxon includes all lower taxonomic units. For example, the genus *Hylobates* (gibbons) is listed as Endangered through-

out its entire range (China, India, and SE Asia); consequently, all species, subspecies, and populations of that genus are considered listed as Endangered for the purposes of the Act. In 1978 (43 FR 6230-6233) the species *Haliaeetus leucocephalus* (bald eagle) was listed as Threatened in "USA (WA, OR, MN, WI, MI)" rather than its entire population; thus, all individuals of the bald eagle found in those five States are considered listed as Threatened for the purposes of the Act.

(h) The "List of Endangered and Threatened Wildlife" is provided below:

Species		Historic range	Vertebrate population where endangered or threatened	Status	When listed	Critical habitat	Special rules
Common name	Scientific name						
MAMMALS							
Anoa, lowland	<i>Bubalus depressicornis</i> (= <i>B. anoa depressicornis</i>)	Indonesia	Entire	E	3	NA	NA
Anoa, mountain	<i>Bubalus quarlesi</i> (= <i>B. anoa quarlesi</i>)	do	do	E	15	NA	NA
Antelope, giant sable	<i>Hippotragus niger varians</i>	Angola	do	E	15	NA	NA
Argali	<i>Ovis ammon hodgsoni</i>	China (Tibet, Himalayas)	do	E	15	NA	NA
Armadillo, giant	<i>Prodonates maximus</i> (= <i>giganteus</i>)	Venezuela and Guyana to Argentina	do	E	15	NA	NA
Armadillo, pink fairy	<i>Chlamyphorus truncatus</i>	Argentina	do	E	3	NA	NA
Ass, African wild	<i>Equus asinus</i> (= <i>africanus</i>)	Somalia, Sudan, Ethiopia	Somalia, Sudan, Ethiopia	E	3, 22	NA	NA
Ass, Asian wild (= kulan, onager)	<i>Equus hemionus</i>	Southwestern and Central Asia	Entire	E	3	NA	NA
Avahi	<i>Avahi</i> (= <i>Lichenotus) laniger</i> (= entire genus)	Madagascar Republic (= Madagascar)	do	E	3	NA	NA
Aye-aye	<i>Daubentonia madagascariensis</i>	Madagascar Republic (= Madagascar)	do	E	3	NA	NA
Babirusa	<i>Babirusa babirusa</i>	Indonesia	do	E	15	NA	NA
Beboon, gelada	<i>Theropithecus gelada</i>	Ethiopia	do	T	18	NA	17.40(c)
Bandicoot, barred	<i>Perameles bougainville</i>	Australia	do	E	4	NA	NA
Bandicoot, desert	<i>Perameles eromiana</i>	do	do	E	6	NA	NA
Bandicoot, lesser rabbit	<i>Macrotis leucura</i>	do	do	E	4	NA	NA
Bandicoot, pig-footed	<i>Cheeropus ecaudatus</i>	do	do	E	4	NA	NA
Bandicoot, rabbit	<i>Macrotis lagotis</i>	do	do	E	4	NA	NA
Banteng	<i>Bos javanicus</i> (= <i>banteng</i>)	Southeast Asia	do	E	3	NA	NA
Bat, Bulmer's fruit (flying fox)	<i>Aproteles bulmerae</i>	Papua New Guinea	do	E	139	NA	NA
Bat, bumblebee	<i>Craseonycteris thonglongyai</i>	Thailand	do	E	139	NA	NA
Bat, grey	<i>Myotis grisescens</i>	Central and Southeastern U.S.A.	do	E	13	NA	NA
Bat, Hawaiian hoary	<i>Lasiurus cinereus semotis</i>	U.S.A. (HI)	do	E	2	NA	NA
Bat, Indiana	<i>Myotis sodalis</i>	Eastern and Midwestern U.S.A.	do	E	1	17.95(a)	NA
Bat, little Mariana fruit	<i>Pteropus tokudae</i>	Western Pacific Ocean: U.S.A. (Guam)	do	E	156	NA	NA
Bat, Mariana fruit	<i>Pteropus mariannus mariannus</i>	Western Pacific Ocean: U.S.A. (Guam, Rota, Tinian, Saipan, Agiguan)	Guam	E	156	NA	NA
Bat, Ozark big-eared	<i>Plecotus townsendi ingens</i>	U.S.A. (MO, OK, AR)	Entire	E	85	NA	NA
Bat, Rodrigues fruit (flying fox)	<i>Pteropus rodricensis</i>	Indian Ocean: Rodrigues Island	do	E	139	NA	NA
Bat, Singapore roundleaf horseshoe	<i>Hipposideros ridleyi</i>	Malaysia	do	E	139	NA	NA
Bat, Virginia big-eared	<i>Plecotus townsendi virginianus</i>	U.S.A. (KY, NC, WV, VA)	do	E	85	17.95(a)	NA
Bear, Baluchistan	<i>Ursus thibetanus gedrosianus</i>	Iran, Pakistan	do	E	233	NA	NA
Bear, brown	<i>Ursus arctos prunensis</i>	China (Tibet)	Guam	E	15	NA	NA
Bear, brown	<i>Ursus arctos arctos</i>	China (Tibet)	Guam	E	15	NA	NA
Bear, brown or grizzly	<i>Ursus arctos</i> (= <i>U. a. harrisi</i>)	Alaska	11 B A 48	T	1, 2, 8	11 A	11 B
Bear, brown or grizzly	<i>Ursus arctos</i> (= <i>U. a. nelsoni</i>)	Holarctic	continental United States				
Beaver	<i>Castor fiber bairdii</i>	Mongolia	Mexico	E	3	NA	NA
Bison, wood	<i>Bison bison athabascae</i>	Canada, Northwestern U.S.A.	Entire	E	18	NA	NA
			Canada	E	3	NA	NA
Bobcat	<i>Felis rufus escuinapas</i>	Central Mexico	Entire	E	15	NA	NA
Bontebok (antelope)	<i>Damaliscus dorcas dorcas</i>	South Africa	do	E	15	NA	NA
Camel, Bactrian	<i>Camelus bactrianus</i> (= <i>ferus</i>)	Mongolia, China	do	E	15	NA	NA
Caribou, woodland	<i>Rangifer tarandus caribou</i>	Canada, U.S.A. (AK, ID, ME, MI, MN, MT, NH, VT, WA, WI)	Canada (that part of SE Brit. Col. bounded by the Can.-USA border.	E	126E, 136E, 143	NA	NA

75

Bobcat	<i>Felis rufus secunipes</i>	Central Mexico	Entire	E	16	NA	NA
Bontebok (antelope)	<i>Demaliscus dixcas dixcas</i>	South Africa	do	E	15	NA	NA
Camel, Bactrian	<i>Camelus bactrianus (= ferus)</i>	Mongolia, China	do	E	15	NA	NA
Caribou, woodland	<i>Rangifer tarandus caribou</i>	Canada, U.S.A. (AK, ID, ME, MI, MN, MT, NH, VT, WA, WI)	Canada (that part of S.E. Bnt Col. bounded by the Can.-USA border, Columbia R., Kootenay R., Kootenay L., and Kootenai R.), U.S.A. (ID, WA)	E	128E, 136E, 143	NA	NA
Cat, Andean	<i>Felis jacobita</i>	Chile, Peru, Bolivia, Argentina	Entire	E	15	NA	NA
Cat, black-footed	<i>Felis nigripes</i>	Southern Africa	do	E	15	NA	NA
Cat, flat-headed	<i>Felis planiceps</i>	Malaysia, Indonesia	do	E	15	NA	NA
Cat, Iriomote	<i>Felis (Mayailurus) iriomotensis</i>	Japan (Iriomote Island, Ryukyu Islands)	do	E	50	NA	NA
Cat, leopard	<i>Felis bengalensis bengalensis</i>	India, Southeast Asia	do	E	15	NA	NA
Cat, marbled	<i>Felis marmorata</i>	Nepal, Southeast Asia, Indonesia	do	E	15	NA	NA
Cat, Pakistan sand	<i>Felis margarita schaffell</i>	Pakistan	do	E	139	NA	NA
Cat, Temminck's (=golden cat)	<i>Felis temminckii</i>	Nepal, China, Southeast Asia, Indonesia (Sumatra)	do	E	15	NA	NA
Cat, tiger	<i>Felis tigrinus</i>	Costa Rica to northern Argentina	do	E	5	NA	NA
Clematis, Apennine	<i>Rupicapra rupicapra ornata</i>	Italy	do	E	15	NA	NA
Cheetah	<i>Acinonyx jubatus</i>	Africa to India	do	E	3, 5	NA	NA
Chimpanzee	<i>Pan troglodytes</i>	West and Central Africa	do	T	16	NA	17 40(c)
Chimpanzee, pygmy	<i>Pan paniscus</i>	Zaire	do	T	16	NA	17 40(c)
Chinchilla	<i>Chinchilla brevicaudata boliviana</i>	Bolivia	do	E	15	NA	NA
Civet, Malabar large-spotted	<i>Viverra zibetha</i>	India	do	E	50	NA	NA
Cochito (=Gulf of California harbor porpoise)	<i>Phocoena sinus</i>	Mexico (Gulf of California)	do	E	169	NA	NA
Colobus, Preuss's red	<i>Colobus bedfordi preussi</i>	Cameroon	do	E	139	NA	NA
Cougar, eastern	<i>Felis concolor cougar</i>	Eastern North America	do	E	8	NA	NA
Deer, Bactrian	<i>Cervus elaphus bactrianus</i>	U.S.S.R., Afghanistan	do	E	50	NA	NA
Deer, Bawean	<i>Axis (=Cervus) porcinus kuhli</i>	Indonesia	do	E	3	NA	NA
Deer, Barbary	<i>Cervus elaphus barbarus</i>	Morocco, Tunisia, Algeria	do	E	50	NA	NA
Deer, Cedros Island mule	<i>Odocoileus hemionus cedrosensis</i>	Mexico (Cedros Island)	do	E	10	NA	NA
Deer, Columbian white-tailed	<i>Odocoileus virginianus leucurus</i>	U.S.A. (WA, OR)	do	E	1	NA	NA
Deer, Corsican red	<i>Cervus elaphus corsicanus</i>	Corsica, Sardinia	do	E	50	NA	NA
Deer, Eld's brow-antlered	<i>Cervus eldi</i>	India to Southeast Asia	do	E	3	NA	NA
Deer, Formosan elk	<i>Cervus nippon taiouanus</i>	Taiwan	do	E	50	NA	NA
Deer, hog	<i>Axis (=Cervus) porcinus annamiticus</i>	Thailand, Indochina	do	E	15	NA	NA
Deer, key	<i>Odocoileus virginianus clavium</i>	U.S.A. (FL)	do	E	1	NA	NA
Deer, marsh	<i>Blastocercus dichotomus</i>	Argentina, Uruguay, Paraguay, Bolivia, Brazil	do	E	3	NA	NA
Deer, McNeill's	<i>Cervus elaphus macneilli</i>	China (Sichiang, Tibet)	do	E	3	NA	NA

U.S. Fish and Wildlife Serv., Interior

§ 17.11

Species		Historic range	Vertebrate population where endangered or threatened	Sta- tus	When listed	Critical habitat	Special rules
Common name	Scientific name						
Deer, musk	<i>Moschus</i> spp. (all species)	Central and East Asia	Afghanistan, Bhutan, Burma, China (Tibet, Yunnan), India, Nepal, Pakistan, Sikkim.	E	15	NA	NA
Deer, North China sika	<i>Cervus nippon mandchuricus</i>	China (Shantung and Chihli Provinces)	Entire	E	50	NA	NA
Deer, pampas	<i>Ozotoceros bezoarticus</i>	Brazil, Argentina, Uruguay, Bolivia, Para- guay.	do.	E	15	NA	NA
Deer, Persian fallow	<i>Dama dama mesopotamica</i>	Iraq, Iran	do.	E	3	NA	NA
Deer, Philippine	<i>Axis (= Cervus) porcinus calamianensis</i>	Philippines (Calamian Islands)	do.	E	15	NA	NA
Deer, Ryukyu sika	<i>Cervus nippon keramae</i>	Japan (Ryukyu Islands)	do.	E	50	NA	NA
Deer, Shanai sika	<i>Cervus nippon grassianus</i>	China (Shanai Province)	do.	E	50	NA	NA
Deer, South China sika	<i>Cervus nippon kopschi</i>	Southern China	do.	E	50	NA	NA
Deer, swamp (= barasingha)	<i>Cervus duvauceli</i>	India, Nepal	do.	E	3	NA	NA
Deer, Yarkand	<i>Cervus elaphus yarkandensis</i>	China (Sinkiang)	do.	E	50	NA	NA
Dhole (= Asiatic wild dog)	<i>Cuon alpinus</i>	U.S.S.R., Korea, China, India, Southeast Asia.	do.	E	3	NA	NA
Dibbler	<i>Antechinus apicalis</i>	Australia	do.	E	4	NA	NA
Dog, African wild	<i>Lycan pictus</i>	Sub-Saharan Africa	do.	E	139	NA	NA
Drill	<i>Papio leucophaeus</i>	Equatorial West Africa	do.	E	18	NA	NA
Dugong	<i>Dugong dugon</i>	East Africa to southern Japan, including U.S.A. (Trust Territories).	do.	E	4	NA	NA
Duiker, Jentink's	<i>Cephalophus jentinki</i>	Sierra Leone, Liberia, Ivory Coast	do.	E	50	NA	NA
Eland, Western giant	<i>Taurotragus oryxianus derbianus</i>	Senegal to Ivory Coast	do.	E	50	NA	NA
Elephant, African	<i>Loxodonta africana</i>	Africa	do.	T	40	NA	17.40(e)
Elephant, Asian	<i>Elephas maximus</i>	South-central and Southeast Asia	do.	E	15	NA	NA
Ferret, black-footed	<i>Mustela nigripes</i>	Western U.S.A., Western Canada	do.	E	1, 3	NA	NA
Fox, Northern swift	<i>Vulpes velox heses</i>	U.S.A. (northern plains), Canada	Canada	E	3	NA	NA
Fox, San Joaquin kit.	<i>Vulpes macrotis nutica</i>	U.S.A. (CA)	Entire	E	1	NA	NA
Fox, Sierran	<i>Canis (Simenia) simensis</i>	Ethiopia	do.	E	50	NA	NA
Gazelle, Clark's (= Dibatag)	<i>Ammodorcas clarkei</i>	Somalia, Ethiopia	do.	E	3	NA	NA
Gazelle, Cuvier's	<i>Gazella cuvieri</i>	Morocco, Algeria, Tunisia	do.	E	3	NA	NA
Gazelle, Mhor	<i>Gazella dama mhorr</i>	Morocco	do.	E	3	NA	NA
Gazelle, Moroccan (= Dorcas)	<i>Gazella dorcas massaesyala</i>	Morocco, Algeria, Tunisia	do.	E	3	NA	NA
Gazelle, Rio de Oro Dame	<i>Gazella dama kozanovi</i>	Western Sahara	do.	E	3	NA	NA
Gazelle, Arabian	<i>Gazella gazelle</i>	Arabian Peninsula, Palestine, Sinai	do.	E	50	NA	NA
Gazelle, sand	<i>Gazella subgutturosa marica</i>	Jordan, Arabian Peninsula	do.	E	50	NA	NA
Gazelle, Saudi Arabian	<i>Gazella dorcas saudya</i>	Israel, Iraq, Jordan, Syria, Arabian Penin- sula.	do.	E	50	NA	NA
Gazelle, Pelzeln's	<i>Gazella dorcas pelzelni</i>	Somalia	do.	E	50	NA	NA
Gazelle, slender-horned (= Rhim)	<i>Gazella leptoceros</i>	Sudan, Egypt, Algeria, Libya	do.	E	3	NA	NA
Gibbons	<i>Hylodactylus</i> spp. (including <i>Normoscus</i>)	China, India, Southeast Asia	do.	E	3, 18	NA	NA

Goat, wild (= Chilian markhor)	<i>Capra sibirica (= falconeri chilianensis)</i>	Southwestern Asia	Chilian Range of west-central Pakistan	E	15	NA	NA
Goral	<i>Nemorrhoea goral</i>						

Goat, wild (= Chilian markhor).....	<i>Capra zegrus (= isioneri chilianensis)</i>	Southwestern Asia.....	Chilian Range of west-central Pakistan.	E	16	NA	NA
Goral.....	<i>Nemorhaedus goral</i>	East Asia.....	Entire.....	E	15	NA	NA
Gorilla.....	<i>Gorilla gorilla</i>	Central and Western Africa.....	do.....	E	3	NA	NA
Guadalupe fur seal.....	<i>Arctoccephalus townsendi</i>	U.S.A. (Farallon Islands of CA) south to Mexico (Isles Revillagigedo).	do.....	T	212	NA	227.11
Hare, hispid.....	<i>Caprolagus hispidus</i>	India, Nepal, Bhutan.....	do.....	E	16	NA	NA
Hartebeest, Swayne's.....	<i>Alcelaphus buselaphus swaynei</i>	Ethiopia, Somalia.....	do.....	E	3, 60	NA	NA
Hartebeest, Tora.....	<i>Alcelaphus buselaphus tora</i>	Ethiopia, Sudan, Egypt.....	do.....	E	60	NA	NA
Hog, pygmy.....	<i>Sus salvanus</i>	India, Nepal, Bhutan, Sikkim.....	do.....	E	3	NA	NA
Horse, Przewalski's.....	<i>Equus przewalski</i>	Mongolia, China.....	do.....	E	15	NA	NA
Huemul, North Andean.....	<i>Hippocamelus antisensis</i>	Ecuador, Peru, Chile, Bolivia, Argentina.....	do.....	E	15	NA	NA
Huemul, South Andean.....	<i>Hippocamelus busculus</i>	Chile, Argentina.....	do.....	E	15	NA	NA
Hutia, Cabrera's.....	<i>Capromys argelsabrali</i>	Cuba.....	do.....	E	233	NA	NA
Hutia, dwarf.....	<i>Capromys nana</i>	do.....	do.....	E	233	NA	NA
Hutia, large-eared.....	<i>Capromys auritus</i>	do.....	do.....	E	233	NA	NA
Hutia, little eared.....	<i>Capromys sentelpensis</i>	do.....	do.....	E	233	NA	NA
Hyena, Barbary.....	<i>Hyena hyaena barbara</i>	Morocco, Algeria, Tunisia.....	do.....	E	3	NA	NA
Hyena, brown.....	<i>Hyena brunnea</i>	Southern Africa.....	do.....	E	3	NA	NA
Ibex, Pyrenean.....	<i>Capra pyrenaica pyrenaica</i>	Spain.....	do.....	E	3	NA	NA
Ibex, Wales.....	<i>Capra wale</i>	Ethiopia.....	do.....	E	3	NA	NA
Impati, black-faced.....	<i>Aepyproros melampus petersi</i>	Namibia, Angola.....	do.....	E	3	NA	NA
Indri.....	<i>Indri indri (= entle genus)</i>	Madagascar (= Madagascar).....	do.....	E	3	NA	NA
Jaguar.....	<i>Panthera onca</i>	U.S.A. (TX, NM, AZ), C. and S. America.....	Mexico southward.....	E	6	NA	NA
Jaguarundi.....	<i>Felis jagouaroundi cacomilli</i>	U.S.A. (TX), Mexico.....	Entire.....	E	15	NA	NA
Jaguarundi.....	<i>Felis jagouaroundi fossate</i>	Mexico, Nicaragua.....	do.....	E	15	NA	NA
Jaguarundi.....	<i>Felis jagouaroundi panamensis</i>	Nicaragua, Costa Rica, Panama.....	do.....	E	15	NA	NA
Jaguarundi.....	<i>Felis jagouaroundi tolfeca</i>	U.S.A. (AZ), Mexico.....	do.....	E	15	NA	NA
Kangaroo, eastern gray.....	<i>Macropus giganteus</i> (all subspecies except tasmanensis).	Australia.....	do.....	T	7	NA	17.40(a)
Kangaroo, red.....	<i>Macropus (Megalotis) rufus</i>	do.....	do.....	T	7	NA	17.40(a)
Kangaroo, Tasmanian forester.....	<i>Macropus giganteus tasmanensis</i>	Australia (Tasmania).....	do.....	E	8	NA	NA
Kangaroo, western gray.....	<i>Macropus fuliginosus</i>	Australia.....	do.....	T	7	NA	17.40(a)
Kouprey.....	<i>Bos sauveli</i>	Vietnam, Laos, Cambodia, Thailand.....	do.....	E	3	NA	NA
Langur, capped.....	<i>Presbytis pileata</i>	India, Burma, Bangladesh.....	do.....	E	15	NA	NA
Langur, entellus.....	<i>Presbytis entellus</i>	China (Tibet), India, Pakistan, Kashmir, Sri Lanka, Sikkim, Bangladesh.....	do.....	E	15	NA	NA
Langur, Douc.....	<i>Pygathrix nemaeus</i>	Cambodia, Laos, Vietnam.....	do.....	E	3	NA	NA
Langur, Francois.....	<i>Presbytis francoisi</i>	China (Kwangsi), Indochina.....	do.....	E	16	NA	NA
Langur, golden.....	<i>Presbytis goldi</i>	India (Assam), Bhutan.....	do.....	E	15	NA	NA
Langur, long-tailed.....	<i>Presbytis potenziani</i>	Indonesia.....	do.....	T	16	NA	17.40(c)
Langur, Pagi Island.....	<i>Nasalis (Simias) concolor</i>	do.....	do.....	E	3	NA	NA
Langur, purple-faced.....	<i>Presbytis senex</i>	Sri Lanka (= Ceylon).....	do.....	T	16	NA	17.40(c)
Langur, Tonkin snub-nosed.....	<i>Pygathrix (Rhinopithecus) avunculus</i>	Vietnam.....	do.....	T	16	NA	17.40(c)
Lechwe, red.....	<i>Kobus leche</i>	Southern Africa.....	do.....	Y	3, 15,	NA	NA
					100		

Species		Historic range	Vertebrate population where endangered or threatened	Status	When listed	Critical habitat	Special rules
Common name	Scientific name						
Lemurs.....	Lemuridae (incl. Cherogaleidae, Lepilemuridae); all members of genera <i>Lemur</i> , <i>Phaner</i> , <i>Haplemur</i> , <i>Lepilemur</i> , <i>Microcebus</i> , <i>Allocebus</i> , <i>Cheirogaleus</i> , <i>Varecia</i> .	Malagasy Republic (= Madagascar).....	do.....	E	3, 15	NA	NA
Leopard.....	<i>Panthera pardus</i>	Africa, Asia.....	Wherever found, except where it is listed as Threatened as set forth below.	E	3, 5, 114	NA	NA
Do.....	do.....	do.....	In Africa, in the wild, south of, and including, the following countries: Gabon, Congo, Zaire, Uganda, Kenya.	T	3, 5, 114	NA	17.40(f)
Leopard, clouded.....	<i>Neofelis nebulosa</i>	Southeast and south-central Asia, Taiwan.	Entire.....	E	3, 15	NA	NA
Leopard, snow.....	<i>Panthera uncia</i>	Central Asia.....	do.....	E	5	NA	NA
Linang, spotted.....	<i>Prionodon pardicolor</i>	Nepal, Assam, Vietnam, Cambodia, Laos, Burma.	do.....	E	15	NA	NA
Lion, Asiatic.....	<i>Panthera leo persica</i>	Turkey to India.....	do.....	E	3	NA	NA
Loris, lesser slow.....	<i>Nycticebus pygmaeus</i>	Indochina.....	do.....	T	18	NA	17.40(c)
Lynx, Spanish.....	<i>Felis (=Lynx) pardina</i>	Spain, Portugal.....	do.....	E	3	NA	NA
Macaque, Formosan rock.....	<i>Macaca cyclops</i>	Taiwan.....	do.....	T	16	NA	17.40(c)
Macaque, Japanese.....	<i>Macaca fuscata</i>	Japan (Shikoku, Kyushu and Honshu Islands).	do.....	T	16	NA	17.40(c)
Macaque, lion-tailed.....	<i>Macaca silenus</i>	India.....	do.....	E	3	NA	NA
Macaque, stump-tailed.....	<i>Macaca arctoides</i>	India (Assam) to southern China.....	do.....	T	16	NA	17.40(c)
Macaque, Tuque.....	<i>Macaca sinica</i>	Sri Lanka (=Ceylon).....	do.....	T	16	NA	17.40(c)
Manatee, Amazonian.....	<i>Trichechus inunguis</i>	South America (Amazon River Basin).....	do.....	E	3	NA	NA
Manatee, West African.....	<i>Trichechus senegalensis</i>	West Coast of Africa from Senegal River to Cuanza River.	do.....	T	52	NA	NA
Manatee, West Indian (Florida).....	<i>Trichechus manatus</i>	U.S.A. (southeastern), Caribbean Sea, South America.	do.....	E	1, 3	17.95(a)	NA
Mandrill.....	<i>Papio sphinx</i>	Equatorial West Africa.....	do.....	E	16	NA	NA
Mangabey, Tana River.....	<i>Cercocebus galantus</i>	Kenya.....	do.....	E	3	NA	NA
Mangabey, white-collared.....	<i>Cercocebus torquatus</i>	Senegal to Ghana; Nigeria to Gabon.....	do.....	E	16	NA	NA
Margay.....	<i>Felis wiedii</i>	U.S.A. (TX), C. and S. America.....	Mexico southward.....	E	6	NA	NA
Markhor, Kabal.....	<i>Capra falconeri megaceros</i>	Afghanistan, Pakistan.....	Entire.....	E	15	NA	NA
Markhor, straight-horned.....	<i>Capra falconeri jerdoni</i>	do.....	do.....	E	15	NA	NA
Marmoset, buff-headed.....	<i>Callithrix flaviceps</i>	Brazil.....	do.....	E	139	NA	NA

78

Marmoset, buff lifted-ear.....	<i>Callithrix jacchus auris</i>	do.....	do.....	E	233	NA	NA
Marmoset, cotton-top.....	<i>Saguinus oedipus</i>	Costa Rica to Colombia.....	do.....	E	16	NA	NA
Marmoset, Goeldi's.....	<i>Callimico goeldii</i>	Brazil, Colombia, Ecuador, Peru, Bolivia.....	do.....	E	3	NA	NA
Marmot, Vancouver Island.....	<i>Marmota vancouverensis</i>	Canada (Vancouver Island).....	do.....	E	176	NA	NA
Marsupial, eastern jerboa.....	<i>Antechinus leniger</i>	Australia.....	do.....	E	176	NA	NA

Marmoset, Kabali	<i>Callithrix jacchus</i>	Alghulistan, Pakistan	Entire	15	NA
Marmoset, straight-horned	<i>Callithrix jacchus</i>	do	do	15	NA
Marmoset, buff-headed	<i>Callithrix flaviceps</i>	Brazil	do	139	NA

(dition)

Marmoset, buffy tufted-ear	<i>Callithrix jacchus aurita</i>	do	do	233	NA	NA
Marmoset, cotton-top	<i>Saguinus oedipus</i>	Costa Rica to Colombia	do	16	NA	NA
Marmoset, Gould's	<i>Callimico goekii</i>	Brazil, Colombia, Ecuador, Peru, Bolivia	do	3	NA	NA
Marmot, Vancouver Island	<i>Marmota vancouverensis</i>	Canada (Vancouver Island)	do	139	NA	NA
Marsupial, eastern jerboa	<i>Antechinomys laniger</i>	Australia	do	4	NA	NA
Marsupial-mouse, large desert	<i>Sminthopsis psammophila</i>	do	do	4	NA	NA
Marsupial-mouse, long-tailed	<i>Sminthopsis longicaudata</i>	do	do	4	NA	NA
Marten, Formosan yellow-throated	<i>Martes flavigula chrysopsala</i>	Taiwan	do	3	NA	NA
Monkey, black colobus	<i>Colobus satanas</i>	Equatorial Guinea, People's Republic of Congo, Cameroon, Gabon	do	16	NA	NA
Monkey, black howler	<i>Alouatta pyra</i>	Mexico, Guatemala, Belize	do	10	NA	17.40(c)
Monkey Diana	<i>Cercopithecus diana</i>	Coastal West Africa	do	16	NA	NA
Monkey, howler	<i>Alouatta palliata (= villosa)</i>	Mexico to South America	do	15	NA	NA
Monkey, L'hoest's	<i>Cercopithecus lhoesti</i>	Upper Eastern Congo Basin, Cameroon	do	16	NA	NA
Monkey, Preuss' red colobus	<i>Colobus badius preussi</i>	Cameroon	do	139	NA	NA
Monkey, proboscis	<i>Nasalis larvatus</i>	Borneo	do	15	NA	NA
Monkey, red-backed squirrel	<i>Saimiri oerstedii</i>	Costa Rica, Panama	do	3	NA	NA
Monkey, red-bellied	<i>Cercopithecus erythrogaster</i>	Western Nigeria	do	16	NA	NA
Monkey, red-eared nose-spotted	<i>Cercopithecus erythrolis</i>	Nigeria, Cameroon, Fernando Po	do	16	NA	NA
Monkey, spider	<i>Ateles geoffroyi frontatus</i>	Costa Rica, Nicaragua	do	3	NA	NA
Monkey, spider	<i>Ateles geoffroyi panamensis</i>	Costa Rica, Panama	do	3	NA	NA
Monkey, Tana River red colobus	<i>Colobus rufomitratus (= badius) rufomitratus</i>	Kenya	do	3, 16	NA	NA
Monkey, woolly spider	<i>Brachyteles arachnoides</i>	Brazil	do	3	NA	NA
Monkey, yellow-tailed woolly	<i>Lagothrix flavicauda</i>	Andes of northern Peru	do	16	NA	NA
Monkey, Zanzibar red colobus	<i>Colobus tirkii</i>	Tanzania	do	3	NA	NA
Mouse, Alabama beach	<i>Peromyscus polionotus ammobates</i>	U.S.A. (AL)	do	183	17.95(a)	NA
Mouse, Australian native	<i>Zyromys (= Notomys) pedunculatus</i>	Australia	do	15	NA	NA
Mouse, Australian native	<i>Notomys aequi</i>	do	do	15	NA	NA
Mouse, Choclawhatchee beach	<i>Peromyscus polionotus allophrys</i>	U.S.A. (FL)	do	183	17.95(a)	NA
Mouse, Field's	<i>Pseudomys fieldi</i>	Australia	do	4	NA	NA
Mouse, Gould's	<i>Pseudomys gouldii</i>	do	do	6	NA	NA
Mouse, Key Largo cotton	<i>Peromyscus gossypinus allapaticola</i>	U.S.A. (FL)	do	131E	NA	NA
Mouse, New Holland	<i>Pseudomys novashollandiae</i>	Australia	do	160	NA	NA
Mouse, Perdido Key beach	<i>Peromyscus polionotus trissyllepsis</i>	U.S.A. (AL, FL)	do	4	NA	NA
Mouse, salt marsh harvest	<i>Rethrodontomys raviventris</i>	U.S.A. (CA)	do	183	17.95(a)	NA
Mouse, Shark Bay	<i>Pseudomys praeonis</i>	Australia	do	2	NA	NA
Mouse, Shortridge's	<i>Pseudomys shortridgei</i>	Australia	do	4	NA	NA
Mouse, Smoky	<i>Pseudomys tumeus</i>	do	do	4	NA	NA
Mouse, western	<i>Pseudomys occidentalis</i>	do	do	4	NA	NA
Muntjac, Fee's	<i>Muntiacus leae</i>	Northern Thailand, Burma	do	4	NA	NA
Native-cat, eastern	<i>Desyurus viverrinus</i>	Australia	do	50	NA	NA
Numbel	<i>Myrmecobius fasciatus</i>	do	do	6	NA	NA
Ocelot	<i>Felis pardalis</i>	U.S.A. (AZ, TX) to C. and S. America	do	4, 6	NA	NA
Orangutan	<i>Pongo pygmaeus</i>	Borneo, Sumatra	do	5, 119	NA	NA
Oryx, Arabian	<i>Oryx leucoryx</i>	Arabian Peninsula	do	3	NA	NA
Otter, Cameroon clawless	<i>Aonyx (Paronyx) congica microdon</i>	Cameroon, Nigeria	do	3	NA	NA

U.S. Fish and Wildlife Serv., Interior

§ 17.11

Species		Historic range	Vertebrate population where endangered or threatened	Sta-tus	When listed	Critical habitat	Special rules
Common name	Scientific name						
Otter, giant	<i>Pteronura brasiliensis</i>	South America	do	E	3	NA	NA
Otter, long-tailed	<i>Lutra longicauda</i> (incl. <i>platensis</i>)	do	do	E	3, 15	NA	NA
Otter, marine	<i>Lutra felina</i>	Peru south to Straits of Magellan	do	E	15	NA	NA
Otter, southern river	<i>Lutra provocax</i>	Chile, Argentina	do	E	15	NA	NA
Otter, southern sea	<i>Enhydra lutris nereis</i>	West coast U.S.A. (WA, OR, CA) south to Mexico (Baja California)	do	T	21	NA	NA
Panda, giant	<i>Ailuropoda melanoleuca</i>	People's Republic of China	do	E	139	NA	NA
Pangolin (= scaly anteater)	<i>Manis temmincki</i>	Africa	do	E	15	NA	NA
Panther, Florida	<i>Felis concolor coryi</i>	U.S.A. (LA and AR east to SC and FL)	do	E	1	NA	NA
Planigale, little	<i>Planigale ingrami subtilissima</i> (formerly <i>P. subtilissima</i>)	Australia	do	E	4	NA	NA
Planigale, southern	<i>Planigale tenuirostris</i>	do	do	E	4	NA	NA
Porcupine, thin-spined	<i>Chaetomys subspinosus</i>	Brazil	do	E	3	NA	NA
Possum, Leadbeater's	<i>Gymnobelideus leadbeateri</i>	do	do	E	233	NA	NA
Possum, mountain pygmy	<i>Burernys parvus</i>	Australia	do	E	4	NA	NA
Possum, scaly-tailed	<i>Wyulfa squamicaudata</i>	do	do	E	4	NA	NA
Prairie dog, Mexican	<i>Cynomys mexicanus</i>	Mexico	do	E	3	NA	NA
Prairie dog, Utah	<i>Cynomys pervidens</i>	U.S.A. (UT)	do	T	8, 149	NA	17 40(g)
Pronghorn, peninsular	<i>Antilocapra americana peninsularis</i>	Mexico (Baja California)	do	E	10	NA	NA
Pronghorn, Sonoran	<i>Antilocapra americana sonoriensis</i>	U.S.A. (AZ), Mexico	do	E	1, 3	NA	NA
Pudu	<i>Pudu pudu</i>	Southern South America	do	E	15	NA	NA
Puma, Costa Rican	<i>Felis concolor costaricensis</i>	Nicaragua, Panama, Costa Rica	do	E	15	NA	NA
Quokka	<i>Setonix brachyurus</i>	Australia	do	E	8	NA	NA
Rabbit, Ryukyu	<i>Pentalagus furnessi</i>	Japan (Ryukyu Islands)	do	E	50	NA	NA
Rabbit, volcano	<i>Romerolagus diazi</i>	Mexico	do	E	3	NA	NA
Rat, fales water	<i>Xeromya myodes</i>	Australia	do	E	4	NA	NA
Rat, Fresno kangaroo	<i>Dipodomys nitratoides exilis</i>	U.S.A. (CA)	do	E	170	17 95(a)	NA
Rat, Moro Bay kangaroo	<i>Dipodomys heermanni morroensis</i>	do	do	E	2	17 95(a)	NA
Rat, stick-nest	<i>Leponillus conditor</i>	Australia	do	E	8	NA	NA
Rat-kangaroo, brush-tailed	<i>Bettongia penicillata</i>	do	do	E	4	NA	NA
Rat-kangaroo, Gaimard's	<i>Bettongia gaimardi</i>	do	do	E	8	NA	NA
Rat-kangaroo, Lesueur's	<i>Bettongia lesueur</i>	do	do	E	4	NA	NA
Rat-kangaroo, plain	<i>Caloprymnus campestris</i>	do	do	E	4	NA	NA
Rat-kangaroo, Queensland	<i>Bettongia tropica</i>	do	do	E	4	NA	NA
Rhinoceros, black	<i>Diceros bicornis</i>	Sub-Saharan Africa	do	E	97	NA	NA
Rhinoceros, great Indian	<i>Rhinoceros unicornis</i>	India, Nepal	do	E	4	NA	NA
Rhinoceros, Javan	<i>Rhinoceros sondaicus</i>	Indonesia, Indochina, Burma, Thailand, Sikkim, Bangladesh, Malaysia	do	E	3	NA	NA
Rhinoceros, northern white	<i>Coelotherium simum cottoni</i>	Zaire, Sudan, Uganda, Central African Republic	do	E	3	NA	NA
Rhinoceros, Sumatran	<i>Dicerorhinus (= Dikermoceros) sumatrensis</i>	Bangladesh to Vietnam to Indonesia (Borneo)	do	E	3	NA	NA
Saiga, Mongolian (antelope)	<i>Saiga tatarica mongolica</i>	Mongolia	do	E	15	NA	NA

Saki, southern beared	<i>Chiropotes satanas satanas</i>	Brazil	do	E	233	NA	NA
Saki, white-nosed	<i>Chiropotes albinus</i>	do	do	E	3	NA	NA
Seal, Caribbean monk	<i>Monachus tropicalis</i>	Caribbean Sea, Gulf of Mexico	do	E	1, 2, 45	NA	NA
Seal, Hawaiian monk	<i>Monachus schauinslandi</i>	U.S.A. (HI)	do	E	18	NA	NA
Seal, Mediterranean monk	<i>Monachus monachus</i>	Mediterranean, Northwest African Coast and Black Sea	do	E	3	NA	NA
Seledang (= Gaur)	<i>Bos gaurus</i>	Bangladesh, Southeast Asia, India	do	E	3	NA	NA
Serow	<i>Capricornis sumatrensis</i>	East Asia, Sumatra	do	E	3	NA	NA
Serval, Barbary	<i>Felis serval constantina</i>	Africa	do	E	3	NA	NA

81

Seal, southern bearded.....	<i>Chiropterus eximius eximius</i>	Brazil	do	E	833	NA	NA
Seal, white-nosed.....	<i>Chiropterus albinus</i>	do	do	E	3	NA	NA
Seal, Caribbean monk.....	<i>Monachus tropicalis</i>	Caribbean Sea, Gulf of Mexico	do	E	1, 2, 46	NA	NA
Seal, Hawaiian monk.....	<i>Monachus schauinslandi</i>	U.S.A. (HI)	do	E	18	NA	NA
Seal, Mediterranean monk.....	<i>Monachus monachus</i>	Mediterranean, Northwest African Coast and Black Sea	do	E	3	NA	NA
Seledang (= Gaur).....	<i>Bos gaurus</i>	Bangladesh, Southeast Asia, India	do	E	3	NA	NA
Serow.....	<i>Capricornis sumatraensis</i>	East Asia, Sumatra	do	E	16	NA	NA
Serval, Barbary.....	<i>Felis serval constantina</i>	Algeria	do	E	3	NA	NA
Shapo.....	<i>Ovis vignei vignei</i>	Kashmir	do	E	16	NA	NA
Shou.....	<i>Cervus elaphus wallichii</i>	Tibet, Bhutan	do	E	3	NA	NA
Shrew, Diamal Swamp southeastern.....	<i>Sorex longirostris fisheri</i>	U.S.A. (VA, NC)	do	T	246	NA	NA
Siamang.....	<i>Symphalangus syndactylus</i>	Malaysia, Indonesia	do	E	15	NA	NA
Sifakas.....	<i>Propithecus</i> spp. (all species)	Malagasy Republic (= Madagascar)	do	E	4	NA	NA
Sloth, Brazilian three-toed.....	<i>Bradypus torquatus</i>	Brazil	do	E	3, 4	NA	NA
Solenodon, Cuban.....	<i>Solenodon (Atopogale) cubanus</i>	Cuba	do	E	3	NA	NA
Solenodon, Haitian.....	<i>Solenodon paraxius</i>	Dominican Republic, Haiti	do	E	3	NA	NA
Squirrel, Carolina northern flying.....	<i>Glaucomys sabrinus coloratus</i>	U.S.A. (NC, TN)	do	E	189	NA	NA
Squirrel, Delmarva Peninsula fox.....	<i>Sciurus nayar cinereus</i>	U.S.A. (Delmarva Peninsula to southeast PA)	do	E	1, 181, 168	NA	NA
Do.....	do	do	do	XN	161	NA	1784(a)
Squirrel, Virginia northern flying.....	<i>Glaucomys sabrinus tuciae</i>	U.S.A. (VA, WV)	do	E	189	NA	NA
Slag, Barbary.....	<i>Cervus elaphus barbarus</i>	Tunisia, Algeria	do	E	3	NA	NA
Slag, Kashmir.....	<i>Cervus elaphus hanglu</i>	Kashmir	do	E	3	NA	NA
Suni, Zanzibar.....	<i>Nesotragus (Nesotragus) moschatus moschatus</i>	Zanzibar (and nearby islands)	do	E	50	NA	NA
Tahr, Arabian.....	<i>Hemitragus jayakari</i>	Oman	do	E	50	NA	NA
Tamaraw.....	<i>Bubalus mindorensis</i>	Philippines	do	E	4	NA	NA
Tamarin, golden-rumped (= golden-headed Tamarin; = golden-blon Marmoset).	<i>Leontopithecus (= Leontideus) spp. (all species)</i>	Brazil	do	E	3	NA	NA
Tamarin, pied.....	<i>Saguinus bicolor</i>	do	do	E	16	NA	NA
Tamarin, white-footed.....	<i>Saguinus leucopus</i>	Colombia	do	T	16	NA	1740(c)
Tapir, Asian.....	<i>Tapirus taticus</i>	Burma, Laos, Cambodia, Vietnam, Malaysia, Indonesia, Thailand	do	E	15	NA	NA
Tapir, Brazilian.....	<i>Tapirus terrestris</i>	Colombia and Venezuela south to Paraguay and Argentina	do	E	3	NA	NA
Tapir, Central American.....	<i>Tapirus bairdi</i>	Southern Mexico to Colombia and Ecuador	do	E	3	NA	NA
Tapir, mountain.....	<i>Tapirus pinchaque</i>	Colombia, Ecuador and possibly Peru and Venezuela	do	E	3	NA	NA
Tarsier, Philippine.....	<i>Tarsius syrichta</i>	Philippines	do	T	16	NA	1740(c)
Tiger.....	<i>Panthera tigris</i>	Temperate and Tropical Asia	do	E	3, 5	NA	NA
Tiger, Tasmanian (= Thylacine)	<i>Thylacinus cynocephalus</i>	Australia	do	E	3	NA	NA
Uekari (all species).....	<i>Cacajao</i> spp (all species)	Peru, Brazil, Ecuador, Colombia, Venezuela	do	E	3	NA	NA
Urbel.....	<i>Ovis musimon (= orientalis) ophion</i>	Cyprus	do	E	15	NA	NA

Species		Historic range	Vertebrate population where endangered or threatened	Status	When listed	Critical habitat	Special rules
Common name	Scientific name						
Vicuna	<i>Vicugna vicugna</i>	South America (Andes)	do	E	3	NA	NA
Vole, Amargosa	<i>Microtus californicus scirpensis</i>	U.S.A. (CA)	do	E	166	17 95(a)	NA
Wallaby, banded hare	<i>Lagostrophus fasciatus</i>	Australia	do	E	4	NA	NA
Wallaby, bridled neck-tailed	<i>Onychogalea fraenata</i>	do	do	E	4	NA	NA
Wallaby, crescent neck-tailed	<i>Onychogalea lunata</i>	do	do	E	4	NA	NA
Wallaby, Parma	<i>Macropus parma</i>	do	do	E	4	NA	NA
Wallaby, Western hare	<i>Lagorchestes hirsutus</i>	do	do	E	4	NA	NA
Wallaby, yellow-footed rock	<i>Petrogale xanthopus</i>	do	do	E	6	NA	NA
Whale, blue	<i>Balaenoptera musculus</i>	Oceanic	do	E	3	NA	NA
Whale, bowhead	<i>Balaena mysticetus</i>	Oceanic (north latitudes only)	do	E	3	NA	NA
Whale, finback	<i>Balaenoptera physalus</i>	Oceanic	do	E	3	NA	NA
Whale, gray	<i>Eschrichtius robustus</i>	North Pacific Ocean: coastal and Bering Sea	do	E	3	NA	NA
Whale, humpback	<i>Megaptera novaeangliae</i>	Oceanic	do	E	3	NA	NA
Whale, right	<i>Balaena glacialis</i>	do	do	E	3	NA	NA
Whale, Sei	<i>Balaenoptera borealis</i>	do	do	E	3	NA	NA
Whale, sperm	<i>Physeter catodon</i>	do	do	E	3	NA	NA
Wolf, gray	<i>Canis lupus</i>	Holarctic	U.S.A. (48 conterminous States, except MN), Mexico	E	1, 8, 13, 15, 35	17.95(a)	NA
Do	do	do	U.S.A. (MN)	T	35	17.95(a)	17.40(d)
Wolf, maned	<i>Chrysocyon brachyurus</i>	Argentina, Bolivia, Brazil, Paraguay, Uruguay	Entire	E	4	NA	NA
Wolf, red	<i>Canis rufus</i>	U.S.A. (southeastern U.S.A. west to central TX)	do	E	1	NA	NA
Wombat, hairy-nosed (=Barnard's and Queensland hairy-nosed)	<i>Lasiornhinus krefftii</i> (formerly <i>L. barnardi</i> and <i>L. gillespiei</i>)	Australia	do	E	4, 6	NA	NA
Woodrat, Key Largo	<i>Neotoma floridana smalli</i>	U.S.A. (FL)	do	E	131E, 160	NA	NA
Yak, wild	<i>Bos grunniens</i>	China (Tibet), India	do	E	3	NA	NA
Zebra, Grevy's	<i>Equus grevyi</i>	Kenya, Ethiopia, Somalia	do	T	64	NA	NA
Zebra, Hartmann's mountain	<i>Equus zebra hartmannae</i>	Namibia, Angola	do	T	54, 111	NA	NA
Zebra, mountain	<i>Equus zebra zebra</i>	South Africa	do	E	15, 111	NA	NA
BIRDS							
Alepa, Hawaii (honeycreeper)	<i>Loxops coccyneus coccyneus</i>	U.S.A. (HI)	do	E	2	NA	NA
Alepa, Maui (honeycreeper)	<i>Loxops coccyneus ochraceus</i>	do	do	E	2	NA	NA
Akialoa, Kauai (honeycreeper)	<i>Hemignathus procerus</i>	do	do	E	1	NA	NA
Akialopau (honeycreeper)	<i>Hemignathus munroi</i> (= <i>wilsoni</i>)	do	do	E	1	NA	NA
Albatross, short-tailed	<i>Diomedea sibiricus</i>	North Pacific Ocean: Japan, U.S.S.R., U.S.A. (AK, CA, HI, OR, WA)	Entire, except U.S.A.	E	3	NA	NA
Blackbird, yellow-shouldered	<i>Agelaius xanthomus</i>	U.S.A. (PR)	Entire	E	17	17.95(b)	NA

Bobwhite, masked (quail)	<i>Colinus virginianus ridgwayi</i>	U.S.A. (AZ), Mexico (Sonora)	do	E	1, 3	NA	NA
Booby, Abbott's	<i>Sula abbotti</i>	Indian Ocean: Christmas Island	do	E	15	NA	NA
Bristlebird, western	<i>Dasyornis brachypterus longirostris</i>	Australia	do	E	3	NA	NA
Bristlebird, western rufous	<i>Dasyornis broadbentii littoralis</i>	do	do	E	3	NA	NA
Broadbill, Guam	<i>Aythya freycineti</i>	do	do	E	15	NA	NA
Bulbul, Mauritius oliveaceous	<i>Hypopetes barboticus oliveaceus</i>	Western Pacific Ocean: U.S.A. (Guam)	do	E	156	NA	NA
Bullfinch, Sao Miguel (finch)	<i>Pytilius pytilius murina</i>	Indian Ocean: Mauritius	do	E	3	NA	NA
Bushwren, New Zealand	<i>Vireo</i>	Eastern Atlantic Ocean: Azores	do	E	3	NA	NA

Bobwhite, masked (quail)	<i>Colinus virginianus ridgwayi</i>	U.S.A. (AZ), Mexico (Sonora)	do	E	1, 3	NA	NA
Booby, Abbott's	<i>Sula abbotti</i>	Indian Ocean: Christmas Island	do	E	15	NA	NA
Bristlebird, western	<i>Dasymotis brachypterus longirostris</i>	Australia	do	E	3	NA	NA
Bristlebird, western rufous	<i>Dasymotis broadbentii mitoralis</i>	do	do	E	15	NA	NA
Broadbill, Guam	<i>Myiagra breycani</i>	Western Pacific Ocean: U.S.A. (Guam)	do	E	15b	NA	NA
Bulbul, Mauritius oliveaceous	<i>Hypsipetes borbonicus oliveaceus</i>	Indian Ocean: Mauritius	do	E	3	NA	NA
Bullfinch, Sao Miguel (finch)	<i>Pyrrhula pyrrhula murina</i>	Eastern Atlantic Ocean: Azores	do	E	3	NA	NA
Bushwren, New Zealand	<i>Xenicus longipes</i>	New Zealand	do	E	3	NA	NA
Bustard, great Indian	<i>Chonotia nigricaps</i>	India, Pakistan	do	E	3	NA	NA
Cahow (=Bermuda Petrel)	<i>Pterodroma cahow</i>	North Atlantic Ocean: Bermuda	do	E	3	NA	NA
Condor, Andean	<i>Vultur gryphus</i>	Colombia to Chile and Argentina	do	E	3	NA	NA
Condor, California	<i>Gymnogyps californianus</i>	U.S.A. (OR, CA), Mexico (Baja California)	do	E	4	NA	NA
Coot, Hawaiian (=also keo keo)	<i>Fulica americana alai</i>	U.S.A. (HI)	do	E	1	17.95(b)	NA
Cotinga, bandud	<i>Cotinga maculata</i>	Brazil	do	E	2	NA	NA
Cotinga, white-winged	<i>Xipholena atropurpurea</i>	do	do	E	15	NA	NA
Crane, black-necked	<i>Grus nigricollis</i>	China (Tibet)	do	E	15	NA	NA
Crane, Cuba sandhill	<i>Grus canadensis nesiotis</i>	West Indies: Cuba	do	E	15	NA	NA
Crane, hooded	<i>Grus monacha</i>	Japan, U.S.S.R.	do	E	15	NA	NA
Crane, Japanese	<i>Grus japonensis</i>	China, Japan, Korea, U.S.S.R.	do	E	4	NA	NA
Crane, Mississippi sandhill	<i>Grus canadensis pulla</i>	U.S.A. (MS)	do	E	3	NA	NA
Crane, Siberian white	<i>Grus leucogeranus</i>	U.S.S.R. (Siberia) to India, including Iran and China	do	E	8	17.95(b)	NA
Crane, white-naped	<i>Grus vipio</i>	Mongolia	do	E	16	NA	NA
Crane, whooping	<i>Grus americana</i>	Canada, U.S.A. (Rocky Mountains east to Carolinas), Mexico	do	E	1, 3	17.95(b)	NA
Creeper, Hawaii	<i>Oreomystis (=Loxops) mana</i>	U.S.A. (HI)	do	E	10	NA	NA
Creeper, Molokai (=kekawahie)	<i>Paroreomystis (=Oreomystis, =Loxops) hemmae</i>	do	do	E	2	NA	NA
Creeper, Oahu (=alaawahio)	<i>Paroreomystis (=Oreomystis, =Loxops) maculata</i>	do	do	E	2	NA	NA
Crow, Hawaiian (=alala)	<i>Corvus hawaiiensis (=tropicus)</i>	do	do	E	1	NA	NA
Crow, Mariana	<i>Corvus kuberyi</i>	Western Pacific Ocean: U.S.A. (Guam, Rota)	do	E	15b	NA	NA
Cuckoo-shrike, Mauritius	<i>Coecus (=Coracina) typicus</i>	Indian Ocean: Mauritius	do	E	3	NA	NA
Cuckoo-shrike, Reunion	<i>Coecus (=Coracina) newtoni</i>	Indian Ocean: Reunion	do	E	3	NA	NA
Curassow, razor-billed	<i>Mitu (=Crax) mitu mitu</i>	Brazil (Eastern)	do	E	3	NA	NA
Curassow, red-billed	<i>Crax blumenbachii</i>	Brazil	do	E	15	NA	NA
Curassow, Trinidad white-headed	<i>Pipilo pipilo pipilo</i>	West Indies: Trinidad	do	E	4	NA	NA
Curlew, Eskimo	<i>Numenius borealis</i>	Alaska and northern Canada to Argentina	do	E	3	NA	NA
Dove, cloven-feathered	<i>Drepanoptila holosericea</i>	Southwest Pacific Ocean: New Caledonia	do	E	1, 3	NA	NA
Dove, Grenada gray-fronted	<i>Leptotila rufaxilla wellsii</i>	West Indies: Grenada	do	E	3	NA	NA
Duck, Hawaiian (=koloa)	<i>Anas wyvilliana</i>	U.S.A. (HI)	do	E	3	NA	NA
Duck, Laysan	<i>Anas laythanensis</i>	do	do	E	1	NA	NA
Duck, pink-headed	<i>Rhodonessa caryophyllacea</i>	India	do	E	1	NA	NA
Duck, white-winged wood	<i>Cairina scutulata</i>	India, Malaysia, Indonesia, Thailand	do	E	15	NA	NA

Species		Historic range	Vertebrate population where endangered or threatened	Sta- tus	When listed	Critical habitat	Special rules
Common name	Scientific name						
Eagle, bald	<i>Haliaeetus leucocephalus</i>	North America south to northern Mexico	U.S.A. (contiguous States, except WA, OR, MN, WI, MI).	E	1, 34	NA	NA
Do	do	do	U.S.A. (WA, OR, MN, WI, MI).	T	34	NA	17.41(a)
Eagle, Greenland white-tailed	<i>Haliaeetus albicilla groenlandicus</i>	Greenland and adjacent Atlantic islands	do	E	15	NA	NA
Eagle, harpy	<i>Harpia harpyja</i>	Mexico south to Argentina	do	E	16	NA	NA
Eagle, Philippine (= monkey-eating)	<i>Pithecopoga jafferyi</i>	Philippines	do	E	3	NA	NA
Eagle, Spanish imperial	<i>Aquila heliaca adalberti</i>	Spain, Morocco, Algeria	do	E	3	NA	NA
Egret, Chinese	<i>Egretta eulophotes</i>	China, Korea	do	E	3	NA	NA
Falcon, American peregrine	<i>Falco peregrinus anatum</i>	Nests from central Alaska across north-central Canada to central Mexico, winters south to South America.	do	E	2, 3, 145	17.95(b)	NA
Falcon, Arctic peregrine	<i>Falco peregrinus tundrius</i>	Nests from northern Alaska to Greenland; winters south to Central and South America.	do	T	2, 3, 145	NA	NA
Falcon, Eurasian peregrine	<i>Falco peregrinus peregrinus</i>	Europe, Eurasia south to Africa and Middle East	do	E	16	NA	NA
Falcon, northern aplomado	<i>Falco femoralis septentrionalis</i>	U.S.A. (AZ, NM, TX), Mexico, Guatemala	do	E	210	NA	NA
Falcon, peregrine	<i>Falco peregrinus</i>	Worldwide, except Antarctica and most Pacific Islands	Wherever found in wild in the contiguous 48 States.	E(S/A)	145	NA	NA
Finch, Laysan (honeycreeper)	<i>Telespiza (= Psittirostra) cantans</i>	U.S.A. (HI)	do	E	1	NA	NA
Finch, Nihoa (honeycreeper)	<i>Telespiza (= Psittirostra) ultima</i>	do	do	E	1	NA	NA
Flycatcher, Euler's	<i>Empidonax euleri johnstoni</i>	West Indies: Grenada	do	E	3	NA	NA
Flycatcher, Seychelles paradise	<i>Terpsiphone corvina</i>	Indian Ocean: Seychelles	do	E	3	NA	NA
Flycatcher, Tahiti	<i>Pomarea nigra</i>	South Pacific Ocean: Tahiti	do	E	3	NA	NA
Fody, Seychelles (weaver-finch)	<i>Foudia sechellarum</i>	Indian Ocean: Seychelles	do	E	3	NA	NA
Frigatebird, Andrew's	<i>Fregate andrewsi</i>	East Indian Ocean	do	E	3	NA	NA
Goose, Aleutian Canada	<i>Branta canadensis leucoparens</i>	do	do	E	15	NA	NA
Goose, Hawaiian (= nene)	<i>Nesochen (= Branta) sandvicensis</i>	U.S.A. (AK, CA, OR, WA), Japan	do	E	1, 3	NA	NA
Goshawk, Christmas Island	<i>Accipiter fasciatus natalis</i>	U.S.A. (HI)	do	E	1	NA	NA
Grackle, slender-billed	<i>Quiscalus (= Cassin's) palustris</i>	Indian Ocean: Christmas Island	do	E	3	NA	NA
Grasswren, Eyrean (flycatcher)	<i>Amytornis gouldi</i>	Mexico	do	E	3	NA	NA
Grebe, Atlanti	<i>Podilymbus gigas</i>	Australia	do	E	3	NA	NA
Greenshank, Nordmann's	<i>Tringa guttifer</i>	Guatemala	do	E	3	NA	NA
Gull, horned	<i>Oreophaps derbianus</i>	U.S.S.R., Japan, south to Malaya, Borneo	do	E	15	NA	NA
Gull, Audouin's	<i>Larus audouinii</i>	Guatemala, Mexico	do	E	3	NA	NA
Gull, relict	<i>Larus relictus</i>	Mediterranean Sea	do	E	3	NA	NA
Hawk, Anjouan Island sparrow	<i>Accipiter francesi pusillus</i>	India, China	do	E	16	NA	NA
		Indian Ocean: Comoro Islands	do	E	3	NA	NA

Hawk, Galapagos	<i>Buteo galapagoensis</i>	Ecuador (Galapagos Islands)	do	E	3	NA	NA
Hawk, Hawaiian (= lo)	<i>Buteo solitarius</i>	U.S.A. (HI)	do	E	1	NA	NA
Hermat, hook-billed (hummingbird)	<i>Glaucis (= Ramphodon) dolniri</i>	Brazil	do	E	16	NA	NA
Honeycreeper, crested (= akotekoke)	<i>Palmeria dolei</i>	U.S.A. (HI)	do	E	1	NA	NA
Hornbill, helmeted	<i>Rhinoplax vigil</i>	Thailand, Malaysia	do	E	16	NA	NA
Honeyeater, helmeted	<i>Meliphaga cassidix</i>	Australia	do	E	4	NA	NA
Ibis, Japanese crested	<i>Nipponia nippon</i>	China, Japan, U.S.S.R., Korea	do	E	3	NA	NA
Kagu	<i>Rhynchotus jubatus</i>	South Pacific Ocean: New Caledonia	do	E	3	NA	NA

Gull, relict *Larus* India, China do 15 NA NA
 Hawk, Anjouan island sparrow *Accipiter francesi pusillus* Indian Ocean, Comoro Islands do 3 NA NA

Hawk, Galapagos	<i>Buteo galapagoensis</i>	Ecuador (Galapagos Islands)	do	E	3	NA	NA
Hawk, Hawaiian (=lo)	<i>Buteo solitarius</i>	U.S.A. (HI)	do	E	1	NA	NA
Hermit, hook-billed (hummingbird)	<i>Glaucus (=Flamphodon) dohrni</i>	Brazil	do	E	15	NA	NA
Honeycreeper, crested (= 'akohekone)	<i>Palmene dohi</i>	U.S.A. (HI)	do	E	1	NA	NA
Hornbill, helmeted	<i>Rhinopias vigil</i>	Thailand, Malaysia	do	E	15	NA	NA
Honeyeater, helmeted	<i>Meliphaga cassidix</i>	Australia	do	E	4	NA	NA
Ibis, Japanese crested	<i>Nipponia nippon</i>	China; Japan, U.S.S.R., Korea	do	E	3	NA	NA
Kagu	<i>Rhynchotus jubatus</i>	South Pacific Ocean: New Caledonia	do	E	3	NA	NA
Kakapo (= owl-parrot)	<i>Strigops habroptilus</i>	New Zealand	do	E	3	NA	NA
Kestrel, Maurilius	<i>Falco punctatus</i>	Indian Ocean: Mauritius	do	E	3	NA	NA
Kestrel, Seychelles	<i>Falco areae</i>	Indian Ocean: Seychelles Islands	do	E	3	NA	NA
Kingfisher, Guam Micronesian	<i>Halcyon cinnamomina cinnamomina</i>	Western Pacific Ocean: U.S.A. (Guam)	do	E	158	NA	NA
Kita, Cuba hook-billed	<i>Chondrohierax uncinatus wilsoni</i>	West Indies: Cuba	do	E	3	NA	NA
Kita, Everglade snail	<i>Rostrhamus sociabilis plumbeus</i>	U.S.A. (FL), Cuba	Florida	E	1	17.95(b)	NA
Kita, Grenada hook-billed	<i>Chondrohierax uncinatus mirus</i>	West Indies: Grenada	Entire	E	3	NA	NA
Kokako (wattlebird)	<i>Callaeas cinerea</i>	New Zealand	do	E	3	NA	NA
Macaw, glaucous	<i>Anodorhynchus glaucus</i>	Paraguay, Uruguay, Brazil	do	E	15	NA	NA
Macaw, indigo	<i>Anodorhynchus leari</i>	Brazil	do	E	15	NA	NA
Macaw, little blue	<i>Cyanopsitta spixii</i>	do	do	E	15	NA	NA
Magpie-robin, Seychelles (thrush)	<i>Copsychus sechellianus</i>	Indian Ocean: Seychelles Islands	do	E	3	NA	NA
Malkoha, red-faced (cuckoo)	<i>Phaenicophaeus pyrrocephalus</i>	Sri Lanka (= Ceylon)	do	E	3	NA	NA
Mallard, Mariana	<i>Anas casteloti</i>	West Pacific Ocean: U.S.A. (Guam, Mariana Islands)	do	E	23	NA	NA
Megapode, Micronesian (= La Perouse's)	<i>Megapodius laepouse</i>	West Pacific Ocean: U.S.A. (Palau Island, Mariana Islands)	do	E	3	NA	NA
Megapode, Maleo	<i>Macrocephalon maleo</i>	Indonesia (Celebes)	do	E	3	NA	NA
Millerbird, Nihoa (old world warbler)	<i>Acrocephalus familiaris kingi</i>	U.S.A. (HI)	do	E	1	NA	NA
Monarch, Tinian (old world flycatcher)	<i>Monarcha takatsukasee</i>	Western Pacific Ocean: U.S.A. (Mariana Islands)	do	E	3	NA	NA
Moorhen (= gallinule), Hawaiian common	<i>Gallinula chloropus sandwicensis</i>	U.S.A. (HI)	do	E	1	NA	NA
Moorhen (= gallinule), Mariana common	<i>Gallinula chloropus guami</i>	Western Pacific Ocean: U.S.A. (Guam, Tinian, Saipan, Pagan)	do	E	158	NA	NA
Nightjar (= whip-poor-will), Puerto Rico	<i>Caprimulgus noctitherus</i>	U.S.A. (PR)	do	E	6	NA	NA
Nukupu (honeycreeper)	<i>Hemignathus lucidus</i>	U.S.A. (HI)	do	E	1, 2	NA	NA
'O'o, Kaula (= 'O'o 'A'a) (honeyeater)	<i>Moho braccatus</i>	do	do	E	1	NA	NA
Ostrich, Arabian	<i>Struthio camelus syriacus</i>	Jordan, Saudi Arabia	do	E	3	NA	NA
Ostrich, West African	<i>Struthio camelus spatzii</i>	Spanish Sahara	do	E	3	NA	NA
'O'u (honeycreeper)	<i>Psittirostra psittacea</i>	U.S.A. (HI)	do	E	1	NA	NA
Owl, Anjouan scops	<i>Otus rutilus capnodes</i>	Indian Ocean: Comoro Island	do	E	3	NA	NA
Owl, giant scops	<i>Otus gurneyi</i>	Philippines: Marinduque and Mindanao Island	do	E	15	NA	NA
Owl, Seychelles	<i>Otus insularis</i>	Indian Ocean: Seychelles Islands	do	E	3	NA	NA
Owlet, Morden's (= Sokoke)	<i>Otus kenae</i>	Kenya	do	E	3	NA	NA
Palia (honeycreeper)	<i>Loxoides (= Psittirostra) bailloni</i>	U.S.A. (HI)	do	E	1	17.95(b)	NA
Parakeet, Forbes'	<i>Cyanoramphus auriceps forbesi</i>	New Zealand	do	E	3, 15	NA	NA
Parakeet, golden	<i>Aratinga guarouba</i>	Brazil	do	E	4	NA	NA
Parakeet, golden-shouldered (= hooded)	<i>Psephotus chrysopterygius</i>	Australia	do	E	3	NA	NA
Parakeet, Mauritius	<i>Psittacula echo</i>	Indian Ocean: Mauritius	do	E	3	NA	NA

U.S. Fish and Wildlife Serv., Interior

§ 17.11

Species		Historic range	Vertebrate population where endangered or threatened	Sta- tus	When listed	Critical habitat	Special rules
Common name	Scientific name						
Parakeet, ochre-marked	<i>Pyrhura cruentata</i>	Brazil	do	E	3	NA	NA
Parakeet, orange-bellied	<i>Neophema chrysogaster</i>	Australia	do	E	4	NA	NA
Parakeet, paradise (= beautiful)	<i>Psephotus pulcherrimus</i>	do	do	E	4	NA	NA
Parakeet, scarlet-chested (= splendid)	<i>Neophema splendida</i>	do	do	E	4	NA	NA
Parakeet, turquoise	<i>Neophema pulchella</i>	do	do	E	3	NA	NA
Parrot, Australian	<i>Geopsittacus occidentalis</i>	do	do	E	3	NA	NA
Parrot, Bahaman or Cuban	<i>Amazona leucocephala</i>	West Indies: Cuba, Bahamas, Caymans	do	E	3, 15	NA	NA
Parrot, ground	<i>Pezoporus wallicus</i>	Australia	do	E	6	NA	NA
Parrot, imperial	<i>Amazona imperialis</i>	West Indies: Dominica	do	E	3	NA	NA
Parrot, Puerto Rican	<i>Amazona vittata</i>	U.S.A. (PR)	do	E	1	NA	NA
Parrot, red browed	<i>Amazona rhodocorytha</i>	Brazil	do	E	3	NA	NA
Parrot, red-capped	<i>Pionopsitta pileata</i>	do	do	E	15	NA	NA
Parrot, red-necked	<i>Amazona arausiaca</i>	West Indies: Dominica	do	E	60	NA	NA
Parrot, red-spectacled	<i>Amazona pretrei pretrei</i>	Brazil, Argentina	do	E	15	NA	NA
Parrot, St. Lucia	<i>Amazona versicolor</i>	West Indies: St. Lucia	do	E	3	NA	NA
Parrot, St. Vincent	<i>Amazona guildingii</i>	West Indies: St. Vincent	do	E	3	NA	NA
Parrot, thick-billed	<i>Rhynchopsitta pachyrhyncha</i>	Mexico, U.S.A. (AZ, NM)	Mexico	E	3	NA	NA
Parrot, vinaceous-breasted	<i>Amazona vinacea</i>	Brazil	Entire	E	15	NA	NA
Parrotbill, Maui (honeysreoper)	<i>Pseudonestor xanthophrys</i>	U.S.A. (HI)	do	E	1	NA	NA
Pelican, brown	<i>Pelecanus occidentalis</i>	U.S.A. (Carolinas to TX, CA), West Indies, C. and S. America: Coastal.	Entire, except U.S. Atlantic coast, FL, AL	E	2, 3, 171	NA	NA
Penguin, Galapagos	<i>Sphaniscus mendiculus</i>	Ecuador (Galapagos Islands)	Entire	E	3	NA	NA
Petrel, Hawaiian dark-rumped	<i>Pterodroma phaeopygia sandwichensis</i>	U.S.A. (HI)	do	E	2, 4, 1	NA	NA
Pheasant, bar-tailed	<i>Symaticus humale</i>	Burma, China	do	E	3	NA	NA
Pheasant, Blyth's tragopan	<i>Trigopon blythii</i>	Burma, China, India	do	E	3	NA	NA
Pheasant, brown eared	<i>Crossoptilon manichuricum</i>	China	do	E	3	NA	NA
Pheasant, Cabot's tragopan	<i>Trigopon caboti</i>	do	do	E	3	NA	NA
Pheasant, Chinese monal	<i>Lophophorus lhuysii</i>	do	do	E	3	NA	NA
Pheasant, Edward's	<i>Lophura edwardsi</i>	Vietnam	do	E	3	NA	NA
Pheasant, Elliot's	<i>Symaticus elioti</i>	China	do	E	15	NA	NA
Pheasant, imperial	<i>Lophura imperialis</i>	Vietnam	do	E	3	NA	NA
Pheasant, Mikado	<i>Symaticus mikado</i>	Taiwan	do	E	3	NA	NA
Pheasant, Palawan peacock	<i>Polyplectron emphanum</i>	Philippines	do	E	3	NA	NA
Pheasant, Slater's monal	<i>Lophophorus sclateri</i>	Burma, China, India	do	E	3	NA	NA
Pheasant, Swinhoe's	<i>Lophura swinhoi</i>	Taiwan	do	E	3	NA	NA
Pheasant, western tragopan	<i>Trigopon melanocephalus</i>	India, Pakistan	do	E	3	NA	NA
Pheasant, white eared	<i>Crossoptilon crossoptilon</i>	China (Tibet), India	do	E	4	NA	NA
Pigeon, Azores wood	<i>Columba palumbus azorica</i>	East Atlantic Ocean: Azores	do	E	3	NA	NA
Pigeon, Chatham Island	<i>Hemiphaea noveeseelandiae chathamensis</i>	New Zealand	do	E	3	NA	NA
Pigeon, Mindoro zone-tailed	<i>Ducula mindorensis</i>	Philippines	do	E	15	NA	NA
Pigeon, Puerto Rican plain	<i>Columba inornata wetmorei</i>	U.S.A. (PR)	do	E	2	NA	NA

Piping guan, black-fronted	<i>Fyxia leucotis</i>	Argentina	do	E	15	NA	NA
Pitta, Koch's	<i>Pitta kochi</i>	Philippines	do	E	15	NA	NA
Plover, New Zealand shore	<i>Thinornis noveeseelandiae</i>	New Zealand	do	E	15	NA	NA
Plover, piping	<i>Charadrius melodus</i>	U.S.A. (Great Lakes, northern Great Plains, Atlantic and Gulf of Mexico)	Great Lakes	E	3	NA	NA

Species		Historic range	Vertebrate population where endangered or threatened	Status	When listed	Critical habitat	Special rules
Common name	Scientific name						
Slit, Hawaiian (= Ae'o)	<i>Himantopus mexicanus</i> (= <i>himantopus</i>) <i>knudseni</i>	U.S.A. (Hawaii)	do	E	2	NA	NA
Stork, oriental white	<i>Ciconia ciconia boyciana</i>	China, Japan, Korea, U.S.S.R.	do	E	3	NA	NA
Stork, wood	<i>Mycteria americana</i>	U.S.A., (CA, AZ, TX, to Carolinas), Mexico, Central and South America	U.S.A. (AL, FL, GA, SC)	E	142	NA	NA
Swiftlet, Mariana gray (= Vanikoro)	<i>Aerodramus</i> (= <i>Collocalia</i>) <i>vanikorensis bartschi</i>	Western Pacific Ocean: U.S.A. (Guam, Rota, Tinian, Saipan, Agiguan)	Entire	E	156	NA	NA
Teal, Campbell Island flightless	<i>Anas aucklandica nesiotis</i>	New Zealand (Campbell Island)	do	E	15	NA	NA
Tern, California least	<i>Sterna antillarum</i> (= <i>albitrons</i>) <i>browni</i>	Mexico, U.S.A. (CA)	do	E	2, 3	NA	NA
Tern, least	<i>Sterna antillarum</i>	U.S.A. (Atlantic and Gulf coasts, Miss. R. Basin, CA), Gr. and Lesser Antilles, Bahamas, Mexico; winters C. America, northern S. America.	U.S.A. (AR, CO, IA, IL, IN, KS, KY, LA (Miss. R. and tribs. N of Baton Rouge), MS (Miss. R.), MO, MT, NE, NM, ND, OK, SD, TN, TX (Except within 50 miles of coast))	E	162	NA	NA
Thrasher, white-breasted	<i>Ramphocinclus brachyurus</i>	West Indies: St. Lucia, Martinique	Entire	E	3	NA	NA
Thrush, large Kauai	<i>Myadestes</i> (= <i>Phaeornis</i>) <i>myadestinus</i>	U.S.A. (HI)	do	E	2	NA	NA
Thrush, Molokai (= oloma'o)	<i>Myadestes</i> (= <i>Phaeornis</i>) <i>lanaiensis</i> (= <i>obscurus</i>) <i>rufa</i>	do	do	E	2	NA	NA
Thrush, New Zealand (wattlebird)	<i>Turnagra capensis</i>	New Zealand	do	E	3	NA	NA
Thrush, small Kauai (= puaihi)	<i>Myadestes</i> (= <i>Phaeornis</i>) <i>palmeri</i>	U.S.A. (HI)	do	E	1	NA	NA
Tinamou, solitary	<i>Tinamus solitarius</i>	Brazil, Paraguay, Argentina	do	E	15	NA	NA
Trembler, Martinique (thrasher)	<i>Cincloerthia ruficauda gutturalis</i>	West Indies: Martinique	do	E	3	NA	NA
Vireo, least Bell's	<i>Vireo bellii pusillus</i>	U.S.A. (CA), Mexico	do	E	228	NA	NA
Wanderer, plain (collared-hempnede)	<i>Pedionomus torquatus</i>	Australia	do	E	6	NA	NA
Warbler (wood), Bachman's	<i>Vermivora bachmani</i>	U.S.A. (Southeastern), Cuba	do	E	1, 3	NA	NA
Warbler (wood), Barbados yellow	<i>Dendroica petechia petechia</i>	West Indies: Barbados	do	E	3	NA	NA
Warbler (wood), Kirtland's	<i>Dendroica kirtlandii</i>	U.S.A. (principally MI), Canada, West Indies: Bahama Islands.	do	E	1, 3	NA	NA
Warbler (willow), nightingale reed	<i>Acrocephalus luscini</i>	Western Pacific Ocean	U.S.A. (Mariana Islands).	E	3, 4	NA	NA
Warbler (willow), Rodrigues	<i>Bebromis rodericanus</i>	Mauritius (Rodrigues Islands)	Entire	E	3	NA	NA
Warbler (wood), Semper's	<i>Leucopoea semperi</i>	West Indies: St. Lucia	do	E	3	NA	NA
Warbler (willow), Seychelles	<i>Bebromis sechellensis</i>	Indian Ocean: Seychelles Island	do	E	3	NA	NA
Whipbird, Western	<i>Psophodes nigrogularis</i>	Australia	do	E	3	NA	NA
White-eye, bridled	<i>Zosterops conspicillata conspicillata</i>	Western Pacific Ocean: U.S.A. (Guam)	do	E	156	NA	NA
White-eye, Norfolk Island	<i>Zosterops albogularis</i>	Indian Ocean: Norfolk Islands	do	E	15	NA	NA
White-eye, Ponape greater	<i>Rukia longirostra</i> (= <i>sanfordi</i>)	West Pacific Ocean: U.S.A. (Caroline Islands)	do	E	3	NA	NA
White-eye, Seychelles	<i>Zosterops modesta</i>	Indian Ocean: Seychelles	do	E	3	NA	NA
Woodpecker, imperial	<i>Campopophilus imperialis</i>	Mexico	do	E	3	NA	NA
Woodpecker, Ivory-billed	<i>Campopophilus principalis</i>	U.S.A. (southcentral and southeastern), Cuba	do	E	1, 3	NA	NA
Woodpecker, red-cockaded	<i>Picoides</i> (= <i>Dendrocopos</i>) <i>borealis</i>	U.S.A. (southcentral and southeastern)	do	E	2	NA	NA
Woodpecker, Titlam's	<i>Dryocopus javensis richardsi</i>	Korea	do	E	3	NA	NA
Wren, Guadeloupe house	<i>Troglodytes aedon guadeloupenis</i>	West Indies: Guadeloupe	do	E	3	NA	NA
Wren, St. Lucia house	<i>Troglodytes aedon mesoleucus</i>	West Indies: St. Lucia	do	E	3	NA	NA

RPT/FA

White-eye, Seychelles	<i>Zosterops modesta</i>	Indian Ocean: Seychelles	do	E	3	NA	NA
Woodpecker, imperial	<i>Campyphylus imperialis</i>	Mexico	do	E	3	NA	NA
Woodpecker, ivory-billed	<i>Campyphylus principalis</i>	U.S.A. (southcentral and southeastern), Cuba	do	E	1, 3	NA	NA
Woodpecker, red-cockaded	<i>Picoides</i> (= <i>Dendrocopos</i>) <i>borealis</i>	U.S.A. (southcentral and southeastern)	do	E	2	NA	NA
Woodpecker, Tristram's	<i>Dryocopus javensis richardsi</i>	Korea	do	E	3	NA	NA
Wren, Guadeloupe house	<i>Troglodytes aedon guadeloupensis</i>	West Indies: Guadeloupe	do	E	3	NA	NA
Wren, St. Lucia house	<i>Troglodytes aedon mesoleucus</i>	West Indies: St. Lucia	do	E	3	NA	NA
REPTILES							
Alligator, American	<i>Alligator mississippiensis</i>	Southeastern U.S.A.	Wherever found in wild except those areas where listed as threatened as set forth below	E	1, 11, 51, 60, 113, 134, 188	NA	NA
Do	do	do	U.S.A. (certain areas of GA and SC, as set forth in 17.42(a)(1), 17.42(b)(1))	T	20, 47, 51, 60, 134, 188	NA	17.42(a)
Do	do	do	U.S.A. (FL, LA and TX); in captivity wherever found	T(S/A)	11, 47, 51, 80, 113, 134, 188	NA	17.42(a)
Alligator, Chinese	<i>Alligator sinensis</i>	China	Entire	E	15	NA	NA
Anole, Culebra Island giant	<i>Anolis roosevelti</i>	U.S.A. (PR: Culebra Island)	do	E	25	17.95(c)	NA
Boa, Jamaican	<i>Epicratis subflavus</i>	Jamaica	do	E	3	NA	NA
Boa, Mona	<i>Epicratis monensis monensis</i>	U.S.A. (PR)	do	T	33	17.95(c)	NA
Boa, Puerto Rico	<i>Epicratis inornatus</i>	do	do	E	2	NA	NA
Boa, Round Island (no common name)	<i>Casarea dussumieri</i>	Indian Ocean: Mauritius	do	E	88	NA	NA
Boa, Round Island (no common name)	<i>Boyeria multocarinata</i>	do	do	E	88	NA	NA
Boa, Virgin Islands tree	<i>Epicratis monensis granti</i>	U.S. and British Virgin Islands	do	E	2, 86	NA	NA
Caiman, Apaporis River	<i>Caiman crocodilus apaporisensis</i>	Colombia	do	E	15	NA	NA
Caiman, black	<i>Melanosuchus nigus</i>	Amazon basin	do	E	15	NA	NA
Caiman, broad-snouted	<i>Caiman latirostris</i>	Brazil, Argentina, Paraguay, Uruguay	do	E	15	NA	NA
Caiman, Yacare	<i>Caiman crocodilus yacare</i>	Bolivia, Argentina, Peru, Brazil	do	E	3	NA	NA
Chuckwalla, San Esteban Island	<i>Sauromatus varus</i>	Mexico	do	E	88	NA	NA
Crocodile, African dwarf	<i>Osteoleaemus tetraspis tetraspis</i>	West Africa	do	E	15	NA	NA
Crocodile, African slender-snouted	<i>Crocodylus cataphractus</i>	Western and central Africa	do	E	5	NA	NA
Crocodile, American	<i>Crocodylus acutus</i>	U.S.A. (FL), Mexico, South America, Central America, Caribbean	do	E	10, 87	17.95(c)	NA
Crocodile, Ceylon mugger	<i>Crocodylus palustris limbata</i>	Sri Lanka	do	E	15	NA	NA
Crocodile, Congo dwarf	<i>Osteoleaemus tetraspis osborni</i>	Conjo River drainage	do	E	15	NA	NA
Crocodile, Cuban	<i>Crocodylus mombifer</i>	Cuba	do	E	3	NA	NA
Crocodile, Morelet's	<i>Crocodylus moreletii</i>	Mexico, Belize, Guatemala	do	E	3	NA	NA
Crocodile, mugger	<i>Crocodylus palustris palustris</i>	India, Pakistan, Iran, Bangladesh	do	E	15	NA	NA
Crocodile, Nile	<i>Crocodylus niloticus</i>	Africa	do	E	3	NA	NA

Species		Historic range	Vertebrate population where endangered or threatened	Status	When listed	Critical habitat	Special rules
Common name	Scientific name						
Crocodile, Orinoco	<i>Crocodylus intermedius</i>	South America: Orinoco River Basin	do	E	3	NA	NA
Crocodile, Philippine	<i>Crocodylus novaeguineae mindorensis</i>	Philippine Islands	do	E	18	NA	NA
Crocodile, saltwater (= estuarine)	<i>Crocodylus porosus</i>	Southeast Asia, Australia, Papua-New Guinea, Pacific Islands.	Entire, except Papua-New Guinea	E	87	NA	NA
Crocodile, Siamese	<i>Crocodylus siamensis</i>	Southeast Asia, Malay Peninsula	Entire	E	15	NA	NA
Gaviel (= gharial)	<i>Gavialis gangeticus</i>	Pakistan, Burma, Bangladesh, India, Nepal.	do	E	3, 15	NA	NA
Gecko, day	<i>Phelsuma edwardnewtoni</i>	Indian Ocean: Mauritius	do	E	3	NA	NA
Gecko, Monito	<i>Sphaerodactylus microplithecus</i>	U.S.A. (PR)	do	E	125	17.95(c)	NA
Gecko, Round Island day	<i>Phelsuma guentheri</i>	Indian Ocean: Mauritius	do	E	3	NA	NA
Gecko, Serpent Island	<i>Cyrtodactylus serpensinsula</i>	do	do	E	3	NA	NA
Iguana, Ackins ground	<i>Cyclura nileyi nuchels</i>	West Indies: Bahamas	do	T	129	NA	NA
Iguana, Allen's Cay	<i>Cyclura cychlura inornata</i>	do	do	T	129	NA	NA
Iguana, Andros Island ground	<i>Cyclura cychlura cychlura</i>	do	do	T	129	NA	NA
Iguana, Anegada ground	<i>Cyclura pinguis</i>	West Indies: British Virgin Islands (Anegada Island).	do	E	3	NA	NA
Iguana, Barrington land	<i>Conolophus pallidus</i>	Ecuador (Galapagos Islands)	do	E	3	NA	NA
Iguana, Cayman Brac ground	<i>Cyclura nubila caymanensis</i>	West Indies: Cayman Islands	do	T	129	NA	NA
Iguana, Cuban ground	<i>Cyclura nubila nubila</i>	Cuba	Entire (excluding population introduced in Puerto Rico).	T	129	NA	NA
Iguana, Exuma Island	<i>Cyclura cychlura figginsi</i>	West Indies: Bahamas	Entire	T	129	NA	NA
Iguana, Fiji banded	<i>Brachylophus fasciatus</i>	Pacific: Fiji, Tonga	do	E	88	NA	NA
Iguana, Fiji crested	<i>Brachylophus vitiensis</i>	Pacific: Fiji	do	E	88	NA	NA
Iguana, Grand Cayman ground	<i>Cyclura nubila lewisi</i>	West Indies: Cayman Islands	do	E	129	NA	NA
Iguana, Jamaican	<i>Cyclura colleyi</i>	West Indies: Jamaica	do	E	129	NA	NA
Iguana, Mayaguana	<i>Cyclura carinata bartschi</i>	West Indies: Bahamas	do	T	129	NA	NA
Iguana, Mona ground	<i>Cyclura stejnegeri</i>	U.S.A. (PR: Mona Island)	do	T	33	17.95(c)	NA
Iguana, Turks and Caicos	<i>Cyclura cannata cannata</i>	West Indies: Turks and Caicos Islands	do	T	129	NA	NA
Iguana, Wetling Island ground	<i>Cyclura nileyi nileyi</i>	West Indies: Bahamas	do	E	129	NA	NA
Iguana, White Cay ground	<i>Cyclura nileyi cristata</i>	do	do	T	129	NA	NA
Lizard, blunt-nosed leopard	<i>Gambelia (= Crotophytus) silus</i>	U.S.A. (CA)	do	E	1	NA	NA
Lizard, Coachella Valley fringe-toed	<i>Uma inornata</i>	do	do	T	105	17.95(c)	NA
Lizard, Hierro giant	<i>Gallotia simonyi simonyi</i>	Spain (Canary Islands)	do	E	144	NA	NA
Lizard, Ibiza wall	<i>Podarcis pityusensis</i>	Spain (Balearic Islands)	do	T	144	NA	NA
Lizard, Island night	<i>Xantusia (= Klausberina) riveralana</i>	U.S.A. (CA)	do	T	26	NA	NA
Lizard, St. Croix ground	<i>Ameiva polops</i>	U.S.A. (VI)	do	E	24	17.95(c)	NA
Monitor, Bengal	<i>Varanus bengalensis</i>	Iran, Iraq, India, Sri Lanka, Malaysia, Afghanistan, Burma, Vietnam, Thailand.	do	E	18	NA	NA

Monitor, desert	<i>Varanus griseus</i>	North Africa to Near East, Caspian Sea through U.S.S.R. to Pakistan, North	do	E	18	NA	NA
-----------------	------------------------	--	----	---	----	----	----

91

Monitor, desert.....	<i>Varanus griseus</i>	North Africa to Nearcast, Caspian Sea through USSR to Pakistan, North-west India.....	do.....	E	18	NA	NA
Monitor, Komodo Island.....	<i>Varanus komodoensis</i>	Indonesia (Komodo, Rinja, Pader, and western Flores Island).....	do.....	E	15	NA	NA
Monitor, yellow.....	<i>Varanus flavescens</i>	West Pakistan through India to Bangladesh.....	do.....	E	15	NA	NA
Python, Indian.....	<i>Python molurus molurus</i>	Sri Lanka and India.....	do.....	E	15	NA	NA
Rattlesnake, Aruba Island.....	<i>Crotalus unicolor</i>	Aruba Island (Netherland Antilles).....	do.....	T	129	NA	NA
Rattlesnake, New Mexican ridge-nosed.....	<i>Crotalus willardi obscurus</i>	U.S.A. (NM), Mexico.....	do.....	T	43	17 95(c)	NA
Skink, Round Island.....	<i>Leiolopisma telfairi</i>	Indian Ocean: Mauritius.....	do.....	T	129	NA	NA
Snake, Atlantic salt marsh.....	<i>Nerodia fasciata tenella</i>	U.S.A. (FL).....	do.....	T	30	NA	NA
Snake, Concho water.....	<i>Nerodia harteri paucimaculata</i>	U.S.A. (TX).....	do.....	T	241	NA	NA
Snake, eastern indigo.....	<i>Drymarchon corais couperi</i>	U.S.A. (AL, FL, GA, MS, SC).....	do.....	E	32	NA	NA
Snake, San Francisco garter.....	<i>Thamnophis sirtalis tetrataenia</i>	U.S.A. (CA).....	do.....	T	1	NA	NA
Tartaruga.....	<i>Podocnemis expansa</i>	South America: Orinoco and Amazon River basins.....	do.....	E	3	NA	NA
Terrapin, river (=Tuntong).....	<i>Bataguraska</i>	Malaysia, Bangladesh, Burma, India, Indonesia.....	do.....	E	3	NA	NA
Tomistoma.....	<i>Tomistoma schlegelii</i>	Malaysia, Indonesia.....	do.....	E	15	NA	NA
Tortoise, angulated.....	<i>Geochelone ymphora</i>	Malagasy Republic (= Madagascar).....	do.....	E	15	NA	NA
Tortoise, Bolson.....	<i>Gopherus flavomarginatus</i>	Mexico.....	do.....	E	46	NA	NA
Tortoise, desert.....	<i>Xerobates (= Scaptochelys = Gopherus) agassizi</i>	U.S.A. (UT, AZ, CA, NV); Mexico.....	Beaver Dam Slope, Utah.....	T	103	17 95(c)	NA
Tortoise, Galapagos.....	<i>Geochelone elephantopus</i>	Ecuador (Galapagos Islands).....	Entire.....	E	3	NA	NA
Tortoise, redated.....	<i>Geochelone (= Testudo) radiata</i>	Malagasy Republic (= Madagascar).....	do.....	E	3	NA	NA
Traceja.....	<i>Podocnemis unifilis</i>	South America: Orinoco and Amazon River basins.....	do.....	E	3	NA	NA
Tuatara.....	<i>Sphenodon punctatus</i>	New Zealand.....	do.....	E	6	NA	NA
Turtle, aquatic box.....	<i>Terrapene carolina</i>	Mexico.....	do.....	E	15	NA	NA
Turtle, black softshell.....	<i>Trionyx nigricans</i>	Bangladesh.....	do.....	E	15	NA	NA
Turtle, Burmese peacock.....	<i>Morenia ocellata</i>	Burma.....	do.....	E	129	NA	NA
Turtle, Central American river.....	<i>Dermatemys mawii</i>	Mexico, Belize, Guatemala.....	do.....	E	15	NA	NA
Turtle, Cuatro Ciénegas softshell.....	<i>Trionyx ater</i>	Mexico.....	do.....	E	15	NA	NA
Turtle, geometric.....	<i>Psammobates geometricus (= Geochelone geometrica)</i>	South Africa.....	do.....	E	15	NA	NA
Turtle, green sea.....	<i>Chelonia mydas</i>	Circumglobal in tropical and temperate seas and oceans.....	Wherever found except where listed as endangered below.....	T	2, 42	NA	17 42(b) and Parts 220 and 227.
Do.....	do.....	do.....	Breeding colony populations in FL and on Pacific coast of Mexico.....	E	2, 42	NA	NA
Turtle, hawkbill sea (= carey).....	<i>Eretmochelys imbricata</i>	Tropical seas.....	Entire.....	E	3	17.95(c)	NA
Turtle, Indian sawback.....	<i>Kachuga tecta tecta</i>	India.....	do.....	E	15	NA	NA
Turtle, Indian softshell.....	<i>Trionyx gangeticus</i>	Pakistan, India.....	do.....	E	15	NA	NA

U.S. Fish and Wildlife Serv., Interior

§ 17.11

Species		Historic range	Vertebrate population where endangered or threatened	Status	When listed	Critical habitat	Special rules
Common name	Scientific name						
Turtle, Kemp's (= Atlantic) Ridley sea	<i>Lepidochelys kempi</i>	Tropical and temperate seas in Atlantic Basin.	do	E	4	NA	NA
Turtle, leatherback sea	<i>Dermochelys coriacea</i>	Tropical, temperate, and subpolar seas	do	E	3	17.95(c), 226.71	NA
Turtle, loggerhead sea	<i>Caretta caretta</i>	Circumglobal in tropical and temperate seas and oceans.	do	T	42	NA	17.42(b) and Parts 220 and 227.
Turtle, Olive (Pacific) Ridley sea	<i>Lepidochelys olivacea</i>	Tropical and temperate seas in Pacific Basin.	Wherever found except where listed as endangered below.	T	42	NA	17.42(b) and Parts 220 and 227.
Do	do	do	Breeding colony populations on Pacific coast of Mexico.	E	42	NA	NA
Turtle, peacock softshell	<i>Trionyx hurum</i>	India, Bangladesh	Entire	E	15	NA	NA
Turtle, Plymouth red-bellied	<i>Pseudemys (= Chrysemys) rubriventris bangsi</i>	U.S.A. (MA)	do	E	90	17.95(c)	NA
Turtle, short-necked or western swamp	<i>Pseudemys urbinii</i>	Australia	do	E	3	NA	NA
Turtle, spotted pond	<i>Geoclemys (= Demyssia) hamiltoni</i>	North India, Pakistan	do	E	15	NA	NA
Turtle, three keeled Asian	<i>Melanochelys (= Geoemyde, Nicoria) tri-carinata</i>	Central India to Bangladesh and Burma	do	E	15	NA	NA
Viper, Lar Valley	<i>Vipera latif</i>	Iran	do	E	129	NA	NA
AMPHIBIANS							
Coqui, golden	<i>Eleutherodactylus jasperi</i>	U.S.A. (PR)	do	T	29	17.95(d)	NA
Frog, Israel painted	<i>Discoglossus nigriventris</i>	Israel	do	E	3	NA	NA
Frog, Panamanian golden	<i>Atelopus varius zeteki</i>	Panama	do	E	15	NA	NA
Frog, Stephen Island	<i>Leiopelma hamiltoni</i>	New Zealand	do	E	3	NA	NA
Salamander, Chinese giant	<i>Andrias davidianus davidianus</i>	Western China	do	E	15	NA	NA
Salamander, desert slender	<i>Batrachoseps andrus</i>	U.S.A. (CA)	do	E	6	NA	NA
Salamander, Japanese giant	<i>Andrias davidianus japonicus</i>	Japan	do	E	15	NA	NA
Salamander, Red Hills	<i>Pleurodactylus hubrichti</i>	U.S.A. (AL)	do	T	19	NA	NA
Salamander, San Marcos	<i>Eurycea nana</i>	U.S.A. (TX)	do	T	98	17.95(d)	17.43(a)
Salamander, Santa Cruz long-toed	<i>Ambystoma macrodactylum croceum</i>	U.S.A. (CA)	do	E	1	NA	NA
Salamander, Texas blind	<i>Typhlomolge rathbuni</i>	U.S.A. (TX)	do	E	1	NA	NA
Toad, African viviparous	<i>Nectophrynoides spp.</i>	Tanzania, Guinea, Ivory Coast, Cameroon, Liberia, Ethiopia.	do	E	15	NA	NA
Toad, Cameroon	<i>Bufo superciliosus</i>	Equatorial Africa	do	E	15	NA	NA
Toad, Houston	<i>Bufo houstonensis</i>	U.S.A. (TX)	do	E	2	17.95(d)	NA
Toad, Monte Verde	<i>Bufo periglenes</i>	Costa Rica	do	E	15	NA	NA

Toad, Wyoming	<i>Bufo hemophrys baxteri</i>	U.S.A. (WY)	do	E	138	NA	NA
FISHES							
Ale Balk (trout)	<i>Salmo platycephalus</i>	Turkey	Entire	E	3	NA	NA
Ayudodoki (loach)	<i>Hymenophysa (= Botia) curta</i>	Japan	do	E	3	NA	NA
Blindcat, Mexican (catfish)	<i>Pristella phreatophila</i>	Mexico	do	E			
Bonytongue, Asian	<i>Schizothorax ferox</i>						

Toad, Wyoming	<i>Bufo hemophrys baxteri</i>	U.S.A. (WY)	do	E	138	NA	NA
FISHES							
Ala Balk (trout)	<i>Salmo platycephalus</i>	Turkey	Enwe	E	3	NA	NA
Ayumodoki (loach)	<i>Hymanophysa (= Botia) curta</i>	Japan	do	E	3	NA	NA
Blindcat, Mexican (catfish)	<i>Priestella phreatophila</i>	Mexico	do	E	3	NA	NA
Bonytongue, Asian	<i>Scleropages formosus</i>	Thailand, Indonesia, Malaysia	do	E	15	NA	NA
Catfish [no common name]	<i>Pangasius sanitwongsei</i>	Thailand	do	E	3	NA	NA
Catfish, giant	<i>Pangasianodon gigas</i>	do	do	E	3	NA	NA
Catfish, Yaqui	<i>Ictalurus pinnai</i>	U.S.A. (AZ), Mexico	do	T	167	17.95(e)	17.44(g)
Cavefish, Alabama	<i>Speoplatythinus poulsoni</i>	U.S.A. (AL)	do	T	28	17.95(e)	NA
Cavefish, Ozark	<i>Amblyopsis rosae</i>	U.S.A. (AR, MO, OK)	do	T	184	NA	NA
Chub, bonytail	<i>Gila elegans</i>	U.S.A. (AZ, CA, CO, NV, UT, WY)	do	E	92	NA	NA
Chub, Borax Lake	<i>Gila boraxobius</i>	U.S.A. (OR)	do	E	124	17.95(u)	NA
Chub, Chihuahua	<i>Gila nigrescens</i>	U.S.A. (NM), Mexico (Chihuahua)	do	T	132	NA	17.44(g)
Chub, humpback	<i>Gila cypha</i>	U.S.A. (AZ, CO, UT, WY)	do	E	1	NA	NA
Chub, Huron tul	<i>Gila bicolor sep</i>	U.S.A. (OR)	do	T	174	NA	17.44(j)
Chub, Mohave tul	<i>Gila bicolor mohavensis</i>	U.S.A. (CA)	do	E	2	NA	NA
Chub, Owens tul	<i>Gila bicolor snyderi</i>	do	do	E	185	17.95(e)	NA
Chub, Pahranagat roundtail	<i>Gila robusta jordani</i>	U.S.A. (NV)	do	E	2	NA	NA
Chub, slender	<i>Hybopsis cahnii</i>	U.S.A. (TN, VA)	do	T	28	17.95(e)	17.44(c)
Chub, Sonora	<i>Gila ditaenia</i>	U.S.A. (AZ), Mexico	do	T	227	17.95(e)	17.44(o)
Chub, spotfin	<i>Hybopsis monache</i>	U.S.A. (AL, GA, NC, TN, VA)	do	T	28	17.95(e)	17.44(c)
Chub, Yaqui	<i>Gila purpurea</i>	U.S.A. (AZ), Mexico	do	E	167	17.95(e)	NA
Cicok (minnow)	<i>Acanthorhynchus hendrichsi</i>	Turkey	do	E	3	NA	NA
Cul-ul	<i>Chasmistes cyjus</i>	U.S.A. (NV)	do	E	1	NA	NA
Dace, Ash Meadows speckled	<i>Rhinichthys osculus nevadensis</i>	do	do	E	117E, 127E, 130	17.95(e)	NA
Dace, desert	<i>Eremichthys acroa</i>	do	do	T	210	17.95(e)	17.44(m)
Dace, Fosken speckled	<i>Rhinichthys osculus sep</i>	U.S.A. (OR)	do	T	174	NA	17.44(j)
Dace, Kendall Warm Springs	<i>Rhinichthys osculus thermata</i>	U.S.A. (WY)	do	E	2	NA	NA
Dace, Moapa	<i>Moapa coriacea</i>	U.S.A. (NV)	do	E	1	NA	NA
Darter, amber	<i>Percina antesella</i>	U.S.A. (GA, TN)	do	E	108	17.95(e)	NA
Darter, bayou	<i>Etheostoma rubrum</i>	U.S.A. (MS)	do	T	10	NA	17.44(b)
Darter, fountain	<i>Etheostoma lenticola</i>	U.S.A. (TX)	do	E	2	17.95(e)	NA
Darter, leopard	<i>Percina pantherina</i>	U.S.A. (AR, OK)	do	T	31	17.95(e)	17.44(d)
Darter, Maryland	<i>Etheostoma sellera</i>	U.S.A. (MD)	do	E	1	NA	NA
Darter, Niangua	<i>Etheostoma nianguae</i>	U.S.A. (MO)	do	T	185	17.95(e)	17.44(k)
Darter, Okaloosa	<i>Etheostoma okaloosae</i>	U.S.A. (FL)	do	E	6	NA	NA
Darter, slackwater	<i>Etheostoma boechungi</i>	U.S.A. (AL, TN)	do	T	28	17.95(e)	17.44(c)
Darter, snail	<i>Percina tenasi</i>	U.S.A. (AL, GA, TN)	do	T	12, 150	NA	NA
Darter, watercress	<i>Etheostoma nuchale</i>	U.S.A. (AL)	do	E	2	NA	NA
Gambusia, Big Bend	<i>Gambusia gaigei</i>	U.S.A. (TX)	do	E	1	NA	NA
Gambusia, Clear Creek	<i>Gambusia heterochir</i>	do	do	E	1	NA	NA
Gambusia, Amated	do	do	do	E	93	NA	NA
Gambusia, Pecos	<i>Gambusia nobilis</i>	U.S.A. (NM, TX)	do	E	2	NA	NA
Gambusia, San Marcos	<i>Gambusia georgei</i>	U.S.A. (TX)	do	E	98	17.95(e)	NA

Species		Historic range	Vertebrate population where endangered or threatened	Status	When listed	Critical habitat	Special rules
Common name	Scientific name						
Killifish, Pahump	<i>Epiplatys lateralis</i>	U.S.A. (NV)	do	E	1	NA	NA
Logperch, Conasauga	<i>Percina jenkinsi</i>	U.S.A. (GA, TN)	do	E	196	17 95(e)	NA
Madtom, Scioto	<i>Noturus trautmani</i>	U.S.A. (OH)	do	E	10	NA	NA
Madtom, Smoky	<i>Noturus baileyi</i>	U.S.A. (TN)	do	E	163	17 95(e)	NA
Madtom, yellowfin	<i>Noturus flavipinnis</i>	U.S.A. (GA, TN, VA)	do	T	26	17 95(e)	17 44(c)
Nakogigi (catfish)	<i>Licobagrus ichikawai</i>	Japan	do	E	3	NA	NA
Pupfish, Ash Meadows Amargosa	<i>Cyprinodon nevadensis mionectes</i>	U.S.A. (NV)	do	E	117E, 127E, 130	17 95(e)	NA
Pupfish, Comanche Springs	<i>Cyprinodon elegans</i>	U.S.A. (TX)	do	E	1	NA	NA
Pupfish, desert	<i>Cyprinodon macularius</i>	U.S.A. (AZ, CA) Mexico	do	E	222	17 95(e)	NA
Pupfish, Devils Hole	<i>Cyprinodon diabolis</i>	U.S.A. (NV)	do	E	1	NA	NA
Pupfish, Leon Springs	<i>Cyprinodon bovinus</i>	U.S.A. (TX)	do	E	102	17 95(e)	NA
Pupfish, Owens	<i>Cyprinodon radiosus</i>	U.S.A. (CA)	do	E	1	NA	NA
Pupfish, Warm Springs	<i>Cyprinodon nevadensis pectoralis</i>	U.S.A. (NV)	do	E	2	NA	NA
Shiner, beautiful	<i>Notropis formosus</i>	U.S.A. (AZ, NM), Mexico	do	T	157	17 95(e)	17 44(g)
Spinedace	<i>Meda fulgida</i>	do	do	T	236	NA	17 44(p)
Spinedace, Big Spring	<i>Lepidomeda mollispinis pratensis</i>	U.S.A. (NV)	do	T	173	17 95(e)	17 44(l)
Spinedace, White River	<i>Lepidomeda albivialis</i>	do	do	E	203	17 95(e)	NA
Springfish, Hiko White River	<i>Crenichthys baileyi grandis</i>	do	do	E	206	17 95(e)	NA
Springfish, Railroad Valley	<i>Crenichthys nevadae</i>	do	do	T	224	17 95(e)	17 44(n)
Springfish, White River	<i>Crenichthys baileyi baileyi</i>	do	do	E	206	17 95(e)	NA
Squawfish, Colorado	<i>Ptychocheilus lucius</i>	U.S.A. (AZ, CA, CO, NM, NV, UT, WY), Mexico	Entire, except Salt and Verde R. drainages, AZ.	E	1, 193	NA	NA
do	do	do	Salt and Verde R. drainages, AZ.	XN	193	NA	17 84(b)
Stickleback, unarmored threespine	<i>Gasterosteus aculeatus williamsoni</i>	U.S.A. (CA)	Entire	E	2	NA	NA
Sturgeon, shortnose	<i>Acipenser brevirostrum</i>	U.S.A. and Canada (Atlantic Coast)	do	E	1	NA	NA
Sucker, June	<i>Chasmistes florus</i>	U.S.A. (UT)	do	E	223	17 95(e)	NA
Sucker, Modoc	<i>Catostomus microps</i>	U.S.A. (CA)	do	E	184	17 95(e)	NA
Sucker, Warner	<i>Catostomus warnerensis</i>	U.S.A. (OR)	do	T	205	17 95(e)	17 44(f)
Tango, Miyako (Tokyo bitterling)	<i>Tanaka tenago</i>	Japan	do	E	3	NA	NA
Temoiak, Ikan (minnow)	<i>Probarbus jullieni</i>	Thailand, Cambodia, Vietnam, Malaysia, Laos	do	T	15	NA	NA
Topminnow, Gila	<i>Poeciliopsis occidentalis</i>	U.S.A. (AZ, NM), Mexico	do	E	1	NA	NA
Totoaba (searout or weakfish)	<i>Omyoscion macdonaldi</i>	Mexico (Gulf of California)	do	E	45	NA	NA
Trout, Apache	<i>Salmo apache</i>	U.S.A. (AZ)	do	T	1, 8	NA	17 44(a)
Trout, Gila	<i>Salmo gilas</i>	U.S.A. (AZ, NM)	do	E	1	NA	NA
Trout, greenback cutthroat	<i>Salmo clarki stomias</i>	U.S.A. (CO)	do	T	1, 38	NA	17 44(f)
Trout, Lahontan cutthroat	<i>Salmo clarki henshawi</i>	U.S.A. (CA, NV)	do	T	2, 8	NA	17 44(a)
Trout, Little Kern golden	<i>Salmo gairdneri whitei</i>	U.S.A. (CA)	do	T	37	17 95(e)	17 44(e)
Trout, Paiute cutthroat	<i>Salmo clarki selenitis</i>	do	do	T	1, 8	NA	17 44(a)

Woundlin	<i>Plagopterus argentissimus</i>	U.S.A. (AZ, NV, UT)	Entire, except Gila R. drainage, AZ, NM.	E	2, 193	NA	NA
do	do	do	Gila R. drainage AZ, NM.	XN	193	NA	17 84(b)
SNAILS							
Snail, Chittenango oval amber	<i>Succinea chittenangoensis</i>	U.S.A. (NY)	NA	T	41	NA	NA
Snail, flat-spined three-toothed	<i>Tridopala platysaides</i>	U.S.A. (WV)	NA	T	41	NA	NA
Snail, Iowa Pleistocene	<i>Physa macclintocki</i>						

Woundfin.....	<i>Pogonius argenteus</i>	U.S.A. (AZ, NV, UT).....	Entire, except Gila R drainage, AZ, NM.	E	2, 193	NA	NA
Do.....	do.....	do.....	Gila R. drainage AZ, NM.	XN	193	NA	17.84(b)
SNAILS							
Snail, Chittanooga ovate amber.....	<i>Succinea chittanoogaensis</i>	U.S.A. (NY).....	NA	T	41	NA	NA
Snail, flat-spined three-toothed.....	<i>Triodopsis platyspyoides</i>	U.S.A. (WV).....	NA	T	41	NA	NA
Snail, Iowa Pleistocene.....	<i>Discus macclintocki</i>	U.S.A. (IA).....	NA	E	41	NA	NA
Snail, Manus Island tree.....	<i>Papustyla pulcherrima</i>	Pacific Ocean: Admiralty Is. (Manus Is.).....	NA	E	3	NA	NA
Snail, noonday.....	<i>Mesodon clarki nantahala</i>	U.S.A. (NC).....	NA	T	41	NA	NA
Snail, Oahu tree.....	<i>Achatineta</i> spp. (all species).....	U.S.A. (HI).....	NA	E	108, 112	NA	NA
Snail, painted snake coiled forest.....	<i>Anguspira picta</i>	U.S.A. (TN).....	NA	T	41	NA	NA
Snail, Stock Island.....	<i>Orthalicus reses</i> (not incl. <i>nasodryas</i>).....	U.S.A. (FL).....	NA	T	41	NA	NA
Snail, Virginia fringed mountain.....	<i>Polygyris virginiensis</i>	U.S.A. (VA).....	NA	E	41	NA	NA
CLAMS							
Pearly mussel, Alabama lamp.....	<i>Lampsis virescens</i>	U.S.A. (AL, TN).....	NA	E	15	NA	NA
Pearly mussel, Appalachian monkeyface.....	<i>Quadrula sparsa</i>	U.S.A. (TN, VA).....	NA	E	15	NA	NA
Pearly mussel, birdwing.....	<i>Conradilla caelata</i>	do.....	NA	E	15	NA	NA
Pearly mussel, Cumberland bean.....	<i>Villosa</i> (= <i>Micromya</i>) <i>trabalis</i>	U.S.A. (KY, TN).....	NA	E	15	NA	NA
Pearly mussel, Cumberland monkeyface.....	<i>Quadrula intermedia</i>	U.S.A. (AL, TN, VA).....	NA	E	15	NA	NA
Pearly mussel, Curtis'.....	<i>Epioblasma</i> (= <i>Dynomia</i>) <i>florentina curti</i>	U.S.A. (MO).....	NA	E	15	NA	NA
Pearly mussel, dromedary.....	<i>Dromus dromes</i>	U.S.A. (TN, VA).....	NA	E	15	NA	NA
Pearly mussel, green-blossom.....	<i>Epioblasma</i> (= <i>Dynomia</i>) <i>torulosa gubernaculum</i>	do.....	NA	E	15	NA	NA
Pearly mussel, Higgins' eye.....	<i>Lampsis higginsii</i>	U.S.A. (IL, IA, MN, MO, NE, WI).....	NA	E	16	NA	NA
Pearly mussel, Nicklin's.....	<i>Megalomias nickliniana</i>	Mexico.....	NA	E	15	NA	NA
Pearly mussel, orange-footed.....	<i>Plethobasus cooperianus</i>	U.S.A. (AL, IN, IA, KY, OH, PA, TN).....	NA	E	15	NA	NA
Pearly mussel, pale liliput.....	<i>Toxolasma</i> (= <i>Cerunculina</i>) <i>cyllindrellus</i>	U.S.A. (AL, TN).....	NA	E	15	NA	NA
Pearly mussel, pink mucket.....	<i>Lampsis orbiculata</i>	U.S.A. (AL, IL, IN, KY, MO, OH, PA, TN, WV).....	NA	E	15	NA	NA
Pearly mussel, Tampico.....	<i>Cyrtomias tampicoensis tocomatensis</i>	Mexico.....	NA	E	15	NA	NA
Pearly mussel, tubercled-blossom.....	<i>Epioblasma</i> (= <i>Dynomia</i>) <i>torulosa torulosa</i>	U.S.A. (IL, IN, KY, TN, WV).....	NA	E	15	NA	NA
Pearly mussel, turgid-blossom.....	<i>Epioblasma</i> (= <i>Dynomia</i>) <i>turgida</i>	U.S.A. (AL, TN).....	NA	E	15	NA	NA
Pearly mussel, white cat's paw.....	<i>Epioblasma</i> (= <i>Dynomia</i>) <i>sulcata delcata</i>	U.S.A. (IN, MI, OH).....	NA	E	15	NA	NA
Pearly mussel, white wartyback.....	<i>Plethobasus cicatricosus</i>	U.S.A. (AL, IN, TN).....	NA	E	15	NA	NA
Pearly mussel, yellow-blossom.....	<i>Epioblasma</i> (= <i>Dynomia</i>) <i>florentina florentina</i>	U.S.A. (AL, TN).....	NA	E	15	NA	NA
Pigtoe, fine-rayed.....	<i>Fusconia cuneata</i>	U.S.A. (AL, TN, VA).....	NA	E	15	NA	NA
Pigtoe, rough.....	<i>Pleurobema planum</i>	U.S.A. (IN, KY, TN, VA).....	NA	E	15	NA	NA
Pigtoe, shiny.....	<i>Fusconia edgariana</i>	U.S.A. (AL, TN, VA).....	NA	E	15	NA	NA
Pocketbook, fat.....	<i>Potamius</i> (= <i>Proptera</i>) <i>capax</i>	U.S.A. (AR, IN, MO, OH).....	NA	E	15	NA	NA
Riffle shell, tan.....	<i>Epioblasma walkeri</i>	U.S.A. (KY, TN, VA).....	NA	E	27	NA	NA

Species		Historic range	Vertebrate population where endangered or threatened	Sta- tus	When listed	Critical habitat	Special rules
Common name	Scientific name						
Spiry mussel, Ter River	<i>Elipido (Cerithyia) stevensiana</i>	U.S.A. (NC)	NA	E	180	NA	NA
CRUSTACEANS							
Amphipod, Hey's Spring	<i>Stygobromus hayi</i>	U.S.A. (DC)	NA	E	118	NA	NA
Crayfish, Nashville	<i>Oreochelone sike</i>	U.S.A. (TN)	NA	E	242	NA	NA
Isopod, Madison Cave	<i>Ambloplites</i>	U.S.A. (VA)	NA	T	123	NA	17 40(a)
Isopod, Socorro	<i>Thermophilus (-Ectophasma)</i>	U.S.A. (NM)	NA	E	36	NA	NA
Shrimp, Kentucky cave	<i>Palaemonetes ganteri</i>	U.S.A. (KY)	NA	E	135	17 95(i)	NA
INSECTS							
Beetle, delta green ground	<i>Elaphus vridis</i>	U.S.A. (CA)	NA	T	100	17 95(i)	NA
Beetle, valley elderberry longhorn	<i>Desmocerus californicus dimorphus</i>	do	NA	T	98	17 95(i)	NA
Butterfly, El Segundo blue	<i>Euphydryas (-Styriaroides) bellatoides silybi</i>	do	NA	E	14	NA	NA
Butterfly, Large's metalmark	<i>Apodemia mormo larigi</i>	do	NA	E	14	NA	NA
Butterfly, lola blue	<i>Lycasides argyrognomon lola</i>	do	NA	E	14	NA	NA
Butterfly, mission blue	<i>Lycasides karibides missionensis</i>	do	NA	E	14	NA	NA
Butterfly, Oregon silverspot	<i>Speyeria zerene hippolyta</i>	U.S.A. (OR, WA)	NA	T	95	17 95(i)	NA
Butterfly, Palos Verdes blue	<i>Glaucopsyche lydianus palosverdesensis</i>	U.S.A. (CA)	NA	E	96	17 95(i)	NA
Butterfly, San Bruno elfin	<i>Callophrys mossii bayensis</i>	do	NA	E	14	NA	NA
Butterfly, Schaus swallowtail	<i>Heliconius (-Papilio) aristodemus pot- carius</i>	U.S.A. (FL)	NA	E	13,158	NA	NA
Butterfly, Smith's blue	<i>Euphydryas (-Styriaroides) anoplos smithi</i>	U.S.A. (CA)	NA	E	14	NA	NA
Moth, Kern pinrose sphinx	<i>Euproserpinus euterpe</i>	do	NA	T	91	NA	NA
Nauccord, Ash Meadows	<i>Amblyscus emargatus</i>	U.S.A. (NV)	NA	T	181	17 95(i)	NA

EDITORIAL NOTE: For "When listed" citations, see list following for symbols in "When listed" see below:
 #—Indicates FR where species was deleted; relisting of the species is indicated by subsequent number(s).
 E—Indicates Emergency rule publication (see FR document for effective dates); subsequent number(s) indicate FR final rule, if applicable under "When listed".

- 1—32 FR 4001; Me
- 2—35 FR 16047; O
- 3—35 FR 8495; Ju
- 4—35 FR 18320; D
- 5—37 FR 6476; Me
- 6—38 FR 14678; Ju
- 7—39 FR 44991; D
- 8—40 FR 29864; Ju
- 9—40 FR 31738; Ju
- 10—40 FR 44151; S
- 11—40 FR 44418; S
- 12—40 FR 47508; C
- 13—41 FR 17740; J
- 14—41 FR 22044; J
- 15—41 FR 24064; J
- 16—41 FR 45993; C
- 17—41 FR 51021; M
- 18—41 FR 51612; M
- 19—41 FR 53034; I
- 20—42 FR 2078; Ja
- 21—42 FR 2968; Ja
- 22—42 FR 15971; M
- 23—42 FR 28137; J
- 24—42 FR 28545; J
- 25—42 FR 37373; J
- 26—42 FR 40885; A
- 27—42 FR 42353; A
- 28—42 FR 45528; S
- 29—42 FR 58755; N
- 30—42 FR 60745; N
- 31—43 FR 3715; Ja
- 32—43 FR 4028; Ja
- 33—43 FR 4621; Fe
- 34—43 FR 6233; Fe
- 35—43 FR 9612; M
- 36—43 FR 12891; M
- 37—43 FR 15428; A
- 38—43 FR 16345; A
- 40—43 FR 20504; M
- 41—43 FR 28932; J
- 42—43 FR 32808; J
- 43—43 FR 34479; A
- 45—44 FR 21289; A
- 46—44 FR 23064; A
- 48—44 FR 29480; M
- 50—44 FR 37126; J
- 51—44 FR 37132; J
- 52—44 FR 42911; J
- 54—44 FR 49220; A
- 55—44 FR 54007; S
- 60—44 FR 59084; O
- 85—44 FR 69208; N
- 86—44 FR 70677; D
- 87—44 FR 75076; D
- 88—45 FR 18010; M
- 90—45 FR 21833; A
- 91—45 FR 24090; A
- 92—45 FR 27713; A
- 93—45 FR 28722; A
- 94—45 FR 35821; M
- 95—45 FR 44935; J
- 96—45 FR 44939; J
- 97—45 FR 47352; J
- 98—45 FR 47355; J
- 99—45 FR 52803; A
- 100—45 FR 52807; J
- 102—45 FR 54678; J

- 1-32 FR 4001: March 11, 1967.
 2-35 FR 16047: October 13, 1970.
 3-35 FR 8495: June 2, 1970.
 4-35 FR 18320: December 2, 1970.
 5-37 FR 8478: March 30, 1972.
 6-38 FR 14678: June 4, 1973.
 7-39 FR 44991: December 30, 1974.
 8-40 FR 29864: July 16, 1975.
 9-40 FR 31736: July 28, 1975.
 10-40 FR 44151: September 25, 1975.
 11-40 FR 44418: September 26, 1975.
 12-40 FR 47506: October 9, 1975.
 13-41 FR 17740: April 28, 1976.
 14-41 FR 22044: June 1, 1976.
 15-41 FR 24064: June 14, 1976.
 16-41 FR 45993: October 19, 1976.
 17-41 FR 51021: November 19, 1976.
 18-41 FR 51612: November 23, 1976.
 19-41 FR 53034: December 3, 1976.
 20-42 FR 2076: January 10, 1977.
 21-42 FR 2968: January 14, 1977.
 22-42 FR 15971, March 24, 1977.
 23-42 FR 28137: June 2, 1977.
 24-42 FR 28545: June 3, 1977.
 25-42 FR 37373: July 21, 1977.
 26-42 FR 40685: August 11, 1977.
 27-42 FR 42353: August 23, 1977.
 28-42 FR 45528: September 9, 1977.
 29-42 FR 58755: November 11, 1977.
 30-42 FR 60745: November 29, 1977.
 31-43 FR 3715: January 27, 1978.
 32-43 FR 4028: January 31, 1978.
 33-43 FR 4621: February 3, 1978.
 34-43 FR 6233: February 14, 1978.
 35-43 FR 9612: March 9, 1978.
 36-43 FR 12691: March 27, 1978.
 37-43 FR 15429: April 13, 1978.
 38-43 FR 16345: April 18, 1978.
 40-43 FR 20504: May 12, 1978.
 41-43 FR 28932: July 3, 1978.
 42-43 FR 32808: July 28, 1978.
 43-43 FR 34479: August 4, 1978.
 45-44 FR 21289: April 10, 1979.
 46-44 FR 23064: April 17, 1979.
 48-44 FR 29480: May 21, 1979.
 50-44 FR 37126: June 25, 1979.
 51-44 FR 37132: June 25, 1979.
 52-44 FR 42911: July 20, 1979.
 54-44 FR 49220: August 21, 1979.
 55-44 FR 54007: September 17, 1979.
 60-44 FR 59084: October 12, 1979.
 85-44 FR 69208: November 30, 1979.
 86-44 FR 70677: December 7, 1979.
 87-44 FR 75076: December 18, 1979.
 88-45 FR 18010: March 20, 1980.
 90-45 FR 21833: April 2, 1980.
 91-45 FR 24090: April 8, 1980.
 92-45 FR 27713: April 23, 1980.
 93-45 FR 28722: April 30, 1980.
 94-45 FR 35821: May 28, 1980.
 95-45 FR 44935: July 2, 1980.
 96-45 FR 44939: July 2, 1980.
 97-45 FR 47352: July 14, 1980.
 98-45 FR 47355: July 14, 1980.
 99-45 FR 52803: August 8, 1980.
 100-45 FR 52807: August 8, 1980.
 102-45 FR 54678: August 15, 1980.
 103-45 FR 55654: August 20, 1980.
 105-45 FR 63812: September 25, 1980.
 108-45 FR 65132: October 1, 1980.
 108-46 FR 3178: January 13, 1981.
 111-46 FR 11665: February 10, 1981.
 112-46 FR 40025: August 6, 1981.
 113-46 FR 40664: August 10, 1981.
 114-47 FR 4204: January 28, 1982.
 115-47 FR 5425: February 5, 1982.
 117-47 FR 19995: May 10, 1982.
 119-47 FR 31670: July 21, 1982.
 123-47 FR 43701: October 4, 1982.
 124-47 FR 43962: October 5, 1982.
 125-47 FR 46093: October 15, 1982.
 127-48 FR 612: January 5, 1983.
 128-48 FR 1726: January 14, 1983.
 129-48 FR 28464: June 22, 1983.
 130-48 FR 40184: September 2, 1983.
 131-48 FR 43043: September 21, 1983.
 132-48 FR 46057: October 11, 1983.
 134-48 FR 46336: October 12, 1983.
 135-48 FR 46341: October 12, 1983.
 136-48 FR 49249: October 25, 1983.
 137-49 FR 1058: January 9, 1984.
 138-49 FR 1994: January 17, 1984.
 139-49 FR 2783: January 23, 1984.
 142-49 FR 7335: February 28, 1984.
 143-49 FR 7394: February 29, 1984.
 144-49 FR 7398: February 29, 1984.
 145-49 FR 10526: March 20, 1984.
 146-49 FR 14356: April 11, 1984.
 149-49 FR 22334: May 29, 1984.
 150-49 FR 27514: July 5, 1984.
 156-49 FR 33885: August 27, 1984.
 157-49 FR 34494: August 31, 1984.
 159-49 FR 34504: August 31, 1984.
 160-49 FR 34510: August 31, 1984.
 161-49 FR 35954: September 13, 1984.
 163-49 FR 43069: October 26, 1984.
 164-49 FR 43969: November 1, 1984.
 166-49 FR 45163: November 15, 1984.
 168-49 FR 49639: December 21, 1984.
 169-50 FR 1056: January 9, 1985.
 170-50 FR 4226: January 30, 1985.
 171-50 FR 4945: February 4, 1985.
 173-50 FR 12302: March 28, 1985.
 174-50 FR 12305: March 28, 1985.
 181-50 FR 20785: May 20, 1985.
 182-50 FR 21792: May 28, 1985.
 183-50 FR 23884: June 6, 1985.
 184-50 FR 24530: June 11, 1985.
 185-50 FR 24653: June 12, 1985.
 186-50 FR 25678: June 20, 1985.
 188-50 FR 26575: June 27, 1985.
 189-50 FR 27002: July 1, 1985.
 193-50 FR 30194: July 24, 1985.
 195-50 FR 31596: August 5, 1985.
 196-50 FR 31603: August 5, 1985.
 203-50 FR 37198: September 12, 1985.
 205-50 FR 39117: September 27, 1985.
 206-50 FR 39123: September 27, 1985.
 210-50 FR 50308: December 10, 1985.
 211-50 FR 50733: December 11, 1985.
 212-50 FR 51252: December 16, 1985.
 216-51 FR 6690: February 25, 1986.
 222-51 FR 10850: March 31, 1986.

223—51 FR 10857; March 31, 1986.
 224—51 FR 10864; March 31, 1986.
 227—51 FR 16047; April 30, 1986.
 228—51 FR 16482; May 2, 1986.
 233—51 FR 17980; May 16, 1986.
 236—51 FR 23781; July 1, 1986.
 239—51 FR 27495; July 31, 1986.
 241—51 FR 31422; September 3, 1986.
 242—51 FR 34412; September 26, 1986.
 246—51 FR 34425; September 26, 1986.

[48 FR 34182, July 27, 1983; 48 FR 34961, Aug. 2, 1983, as amended at 48 FR 39943, Sept. 2, 1983; 48 FR 46337, Oct. 12, 1983; 48 FR 52743, Nov. 22, 1983; 49 FR 1058, Jan. 9, 1984; 49 FR 33892, Aug. 27, 1984]

EDITORIAL NOTE For additional FEDERAL REGISTER citations affecting the table in § 17.11(h), see the listing which follows the table.

EFFECTIVE DATE NOTE At 51 FR 34412, 34425, Sept. 26, 1986, the table in § 17.11(h) was amended by adding "Shrew, Dismal Swamp southeastern" alphabetically under "Mammals" and adding "Crayfish, Nashville" alphabetically under "Crustaceans", effective October 27, 1986.

§ 17.12 Endangered and threatened plants.

(a) The list in this section contains the names of all species of plants which have been determined by the Services to be Endangered or Threatened. It also contains the names of species of plants treated as Endangered or Threatened because they are sufficiently similar in appearance to Endangered or Threatened species (see § 17.50 *et seq.*).

(b) The columns entitled "Scientific name" and "Common name" define the species of plant within the meaning of the Act. Although common names are included, they cannot be relied upon for identification of any specimen, since they may vary greatly in local usage. The Services shall use the most recently accepted scientific name. In cases in which confusion might arise, a synonym(s) will be provided in parentheses. The Services shall rely to the extent practicable on the *International Code of Botanical Nomenclature*.

(c) In the "Status" column the following symbols are used: "E" for Endangered, "T" for Threatened, and "E [or T] (S/A)" for similarity of appearance species.

(d) The other data in the list are nonregulatory in nature and are provided for the information of the

reader. In the annual revision and compilation of this title, the following information may be amended without public notice: the spelling of species' names, historical range, footnotes, references to certain other applicable portions of this title, synonyms, and more current names. In any of these revised entries, neither the species, as defined in paragraph (b) of this section, nor its status may be changed without following the procedures of Part 424 of this title.

(e) The "Historic range" indicates the known general distribution of the species or subspecies as reported in the current scientific literature. The present distribution may be greatly reduced from this historic range. This column does not imply any limitation on the application of the prohibitions in the Act or implementing rules. Such prohibitions apply to all individuals of the plant species, wherever found.

(f)(1) A footnote to the FEDERAL REGISTER publication(s) listing or reclassifying a species is indicated under the column "When listed." Footnote numbers to §§ 17.11 and 17.12 are in the same numerical sequence, since plants and animals may be listed in the same FEDERAL REGISTER document. That document, at least since 1973, includes a statement indicating the basis for the listing, as well as the effective date(s) of said listing.

(2) The "Special rules" and "Critical habitat" columns provide a cross reference to other sections in Parts 17, 222, 226, or 227. The "Special rules" column will also be used to cite the special rules which describe experimental populations and determine if they are essential or nonessential. Separate listings will be made for experimental populations, and the status column will include the following symbols: "XE" for an essential experimental population and "XN" for a nonessential experimental population. The term "NA" (not applicable) appearing in either of these two columns indicates that there are no special rules and/or critical habitat for that particular species. However, all other appropriate rules in Parts 17, 217 through 227, and 402 still apply to that species. In addition, there may be other rules in this title that relate to

such plants. I
 erences
 list all th
 ices whi
 or to th

Species		Historic range	Sta- tus	When listed	Critical habitat	Special rules	
Scientific name	Common name						
Agavaceae—Agave family: <i>Agave arizonica</i>		Arizona agave.....	U.S.A. (AZ).....	E	147	NA	NA
Alismaceae—Water-plantain family: <i>Sagittaria fasciculata</i>		Bunched arrowhead.....	U.S.A. (NC, SC).....	E	63	NA	NA
Amaranthaceae—Amaranth family: <i>Achyranthes rotundata</i>		None.....	U.S.A. (HI).....	E	220	NA	NA
Annonaceae—Custard-apple family: <i>Asimina tetramera</i>		Four-petal pawpaw.....	U.S.A. (FL).....	E	244	NA	NA
		<i>Deeringothamnus pulchellus</i>do.....	E	244	NA	NA
		<i>Deeringothamnus rugeli</i>do.....	E	244	NA	NA
Apiaceae—Parsley family: <i>Orypolis canbyi</i>		Canby's dropwort.....	U.S.A. (DE, GA, MD, NC, SC).....	E	217	NA	NA
Apocynaceae—Dogbane family: <i>Cycladenia humilis</i> var. <i>jonesii</i>		Jones cycladenia.....	U.S.A. (AZ, UT).....	T	229	NA	NA
Asteraceae—Aster family: <i>Argyroxiphum sandwicense</i> ssp. <i>sandwicense</i>		'Ahihina (Mauna Kea silversword).....	U.S.A. (HI).....	E	219	NA	NA
		<i>Bidens cuneata</i>do.....	E	141	NA	NA
		<i>Chrysopsis floridana</i> (= <i>Heterotheca floridana</i>).....	Florida golden aster.....	E	232	NA	NA
		<i>Dysodia tephroleuca</i>	U.S.A. (TX).....	E	152	NA	NA
		<i>Echinacea tennesseensis</i>	Tennessee purple coneflower.....	E	49	NA	NA
		<i>Encelopsis nudaucalis</i> var. <i>conjugata</i>	Ash Meadows sunray.....	T	181	17.96(a)	NA
		<i>Erigeron maguirei</i> var. <i>maguirei</i>	U.S.A. (NV).....	E	202	NA	NA
		<i>Erigeron rhizomatus</i>	Maguire daisy.....	T	177	NA	NA
		<i>Grindelia travinoprasensis</i>	Rhizome fleabane.....	E	181	17.96(a)	NA
		<i>Hymenoxys texana</i>	Ash Meadows gumplant.....	T	218	NA	NA
		<i>Lipochaeta venosa</i>	U.S.A. (TX).....	E	73	NA	NA
		<i>Pityopsis ruthii</i> (= <i>Heterotheca ruthii</i> , = <i>Chrysopsis ruthii</i>).....	U.S.A. (HI).....	E	191	NA	NA
		<i>Senecio franciscanus</i>	Ruth's golden aster.....	T	137	17.96(a)	NA
		<i>Solidago shortii</i>	San Francisco Peaks groundsel.....	E	201	NA	NA
		<i>Solidago spilihamae</i>	U.S.A. (KY).....	T	175	NA	NA
		<i>Stephanomeria malheurensis</i>	Short's goldenrod.....	E	128	17.96(a)	NA
		<i>Townsendia aprica</i>	Blue Ridge goldenrod.....	T	200	NA	NA
			Malheur wire-lettuce.....	E	128	17.96(a)	NA
			Last Chance townsendia.....	T	200	NA	NA
Berberidaceae—Barberry family: <i>Mahonia sonnei</i> (= <i>Berberis</i> s.).....		Truckee barberry.....	U.S.A. (CA).....	E	76	NA	NA
Betulaceae—Birch family: <i>Betula ulmifera</i>		Virginia round-leaf birch.....	U.S.A. (VA).....	E	39	NA	NA

Boraginaceae—Borage family:

Amalanchia grandiflora.....

Large-flowered liddeneck.....

U.S.A. (CA).....

E

179

17.96(a)

NA

Brassicaceae—Mustard family:

Arabis mcdonaldiana.....

McDonald's rock-creep.....

U.S.A. (CA).....

E

44

NA

NA

Erysimum capitatum var. *angustatum*.....

Contra Costa wallflower.....

.....do.....

E

39

17.96(a)

NA

Thelypodium stenopetalum.....

Slender-petaled mustard.....

.....do.....

E

158

NA

NA

Buxaceae—Boxwood family:

Buxus vahlia.....

Vahl's boxwood.....

U.S.A. (PR).....

E

197

NA

NA

Celastraceae—Cedar family:

Barberraceae—Barberry family:
Mahonia sonnei (= Berberis s.)

9 1 1 1 8 1 1 1 1 1 9

juva round leaf birch

U S A. (VA)

E

39

NA

NA

Betuleaceae—Birch family:
Betula ulmi

Boraginaceae—Borage family:
Amarantus grandiflorus

Brassicaceae—Mustard family:
Arabis macdonaldiana
Erysimum capitatum var. angustatum
Thelypodium stenopetalum

Buxaceae—Boxwood family:
Buxus vahli

Cactaceae—Cactus family:

Ancistrocactus tobuschii (= Echinocactus t., Mammillaria t.)
Cereus enoplorus var. flagrans
Cereus robbini
Coryphantha minima (= C. neliae, Escobaria n., Mammillaria n.)
Coryphantha ramulosa

Coryphantha robbinsorum (= Cochisea r., Escobaria r.)

Coryphantha sneedi var. laei (= Escobaria l., Mammillaria l.)
Coryphantha sneedi var. sneedi (= Escobaria s., Mammillaria s.)
Echinocactus horizontalis var. nicholii
Echinocereus engelmannii var. purpureus
Echinocereus kuenzleri var. kuenzleri (= E. kuenzleri, E. hempeli of authors, not Fobes)

Echinocereus lloydii (= E. roetteri var. l.)
Echinocereus reichenbachii var. stertii (= E. melanocentrus)
Echinocereus triglochidatus var. arizonicus (= E. arizonicus)
Echinocereus triglochidatus var. inermis (= E. coccineus var. l., E. phoeniceus var. l.)
Echinocereus vridiflorus var. davisi (= E. davisi)
Neolloydia mariposensis (= Echinocactus m., Echinomastus m.)

Pediocactus bradyi (= Tourneya b.)
Pediocactus knowltoni (= P. bradyi var. k. Tourneya k.)
Pediocactus peeblesianus var. peeblesianus (= Echinocactus p., Navajo p., Tourneya p., Utahia p.)

Pediocactus stierli (= Echinocactus s., Utahia s.)
Sclerocactus glaucus (= Echinocactus g., E. subglaucus, E. whipplei var. g.,
Pediocactus g., S. franklini, S. whipplei var. g.)
Sclerocactus mesa-verdeae (= Colorado m., Echinocactus m., Peducactus m.)
Sclerocactus wrightii (= Pediocactus w.)

Caryophyllaceae—Pink family:
Schiedea adamanis

Large-flowered fiddleneck	U.S.A. (CA)	E	179	17.96(a)	NA
McDonald's rock-cree	U.S.A. (CA)	E	44	NA	NA
Contra Costa wallflower	do	E	39	17.96(a)	NA
Slender-petaled mustard	do	E	158	NA	NA
Vahl's boxwood	U.S.A. (PR)	E	197	NA	NA
Tobusch fleehook cactus	U.S.A. (TX)	E	80	NA	NA
Fragrant prickly-apple	U.S.A. (FL)	E	208	NA	NA
Key tree-cactus	U.S.A. (FL), Cuba	E	153	NA	NA
Nellie cory cactus	U.S.A. (TX)	E	81	NA	NA
Bunched cory cactus	U.S.A. (TX), Mexico (Cochula)	T	77	NA	NA
Cochise pincushion cactus	U.S.A. (AZ), Mexico (Sonora)	T	214	NA	NA
Lee pincushion cactus	U.S.A. (NM)	T	81	NA	NA
Sneed pincushion cactus	U.S.A. (TX, NM)	E	82	NA	NA
Nichol's Turk's head cactus	U.S.A. (AZ)	E	71	NA	NA
Purple-spined hedgehog cactus	U.S.A. (UT)	E	58	NA	NA
Kuenzler hedgehog cactus	U.S.A. (NM)	E	70	NA	NA
Lloyd's hedgehog cactus	U.S.A. (TX)	E	67	NA	NA
Black lace cactus	do	E	68	NA	NA
Arizona hedgehog cactus	U.S.A. (AZ)	E	82	NA	NA
Spineless hedgehog cactus	U.S.A. (CO, UT)	E	83	NA	NA
Davis' green piteya	U.S.A. (TX)	E	81	NA	NA
Lloyd's Mariposa cactus	U.S.A. (TX), Mexico (Cochula)	T	77	NA	NA
Brady pincushion cactus	U.S.A. (AZ)	E	63	NA	NA
Knowlton cactus	U.S.A. (NM, CO)	E	72	NA	NA
Peebles Navajo cactus	U.S.A. (AZ)	E	69	NA	NA
Siler pincushion cactus	U.S.A. (AZ, UT)	E	64	NA	NA
Uinta Basin hookless cactus	U.S.A. (CO, UT)	T	69	NA	NA
Mesa Verde cactus	U.S.A. (CO, NM)	T	75	NA	NA
Wright fishhook cactus	U.S.A. (UT)	E	58	NA	NA
Diamond Head schledes	U.S.A. (HI)	E	141	NA	NA

<i>Nitrophila mohaveensis</i>	Amargosa waterwort.....	U.S.A. (CA)				
Cistaceae—Rockrose family: <i>Hudsonia montana</i>	Mountain golden heather.....	U.S.A. (NC)	Q	107	17.00(a)	NA
Crassulaceae—Stonecrop family: <i>Dudleya traskiae</i>	Santa Barbara Island liveforever.....	U.S.A. (CA)	E	39	NA	NA
Cucurbitaceae—Gourd family: <i>Tumamoca maddockii</i>	Tumamoc globe-berry.....	U.S.A. (AZ), Mexico (Sonora)	E	228	NA	NA
Cupressaceae—Cypress family: <i>Fitzroya cupressoides</i>	Chilean false larch (= alerce).....	Chile, Argentina	T	79	NA	NA
Cyperaceae—Sedge family: <i>Carex specuicola</i>	None.....	U.S.A. (AZ)	T	178	7.00(a)	NA
Ericaceae—Heath family: <i>Arctostaphylos purpurea</i> var. <i>ravenii</i> (= <i>A. hookeri</i> ssp. <i>ravenii</i>) <i>Rhododendron chapmanii</i>	Presidio (= Raven's) manzanilla Chapman rhododendron.....	U.S.A. (CA) U.S.A. (FL)	E E	65 47	NA NA	NA NA
Euphorbiaceae—Spurge family: <i>Euphorbia</i> (= <i>Chamaesyce</i>) <i>deltoides</i> ssp. <i>deltoides</i> <i>Euphorbia</i> (= <i>Chamaesyce</i>) <i>garberi</i> <i>Euphorbia skottsbergii</i> var. <i>kalaheleana</i> <i>Jatropha costaricensis</i>	Spurge..... do..... Ewa Plains 'akoko Costa Rican <i>Jatropha</i>	U.S.A. (FL) do U.S.A. (HI) Costa Rica	E T E E	192 192 120 154	NA NA NA NA	NA NA NA NA
Fabaceae—Pea family: <i>Amorpha crenulata</i> <i>Astragalus humillimus</i> <i>Astragalus perianus</i> <i>Astragalus phoenix</i> <i>Baptisia arachnifera</i> <i>Galactia smallii</i> <i>Hoffmannseggia tenella</i> <i>Lotus dendroideus</i> ssp. <i>traskiae</i> (= <i>L. scopertus</i> ssp. <i>?</i>) <i>Mazoneuron kavaiense</i> <i>Vicia menziesii</i>	Crenulate lead-plant..... Manco's milk-vetch..... Rydberg milk-vetch..... Ash Meadows milk-vetch..... Hairy rattletweed..... Small's milkpea..... Slender rush-pea..... San Clemente Island broom..... Uhiuhi..... Hawaiian vetch.....	U.S.A. (FL) U.S.A. (CO, NM) U.S.A. (UT) U.S.A. (NV) U.S.A. (GA) U.S.A. (FL) U.S.A. (TX) U.S.A. (CA) U.S.A. (HI) do	E E T T E E E E E E	192 167 39 181 39 192 209 26 238 39	NA NA NA 17.00(a) NA NA NA NA NA NA	NA NA NA NA NA NA NA NA NA NA
Frankeniaceae—Frankenia family: <i>Frankenia johnstonii</i>	Johnston's frankenia.....	U.S.A. (TX), Mexico (Nuevo Leon)	E	155	NA	NA

Gentianaceae—Gentian family: <i>Centaurium namphyllum</i>	Spring-loving centaury.....	U.S.A. (CA, NV)	T	181	17.00(a)	NA
Goodeniaceae—Goodenia family: <i>Scaevola coriacea</i>	Dwarf naupaka.....	U.S.A. (HI)	E	231	NA	NA
Hydrophyllaceae—Waterleaf family: <i>Phacelia argillacea</i> <i>Phacelia formosula</i>	Clay phacelia North Park phacelia.....	U.S.A. (UT) U.S.A. (CO)	E E	44 121	NA NA	NA NA

Frankeniaceae—Frankenia family:
Frankenia johnstonii

Johnston

erie

U.S.A. (TX), Mexico
(Bluevale Loc.)

E

188

NA

NA

(ition)

Gentianaceae—Gentian family: <i>Gentiana nana</i>	Spring-loving gentian	U.S.A. (CA, NV)	T	181	17.98(a)	NA
Goodeniaceae—Goodenia family: <i>Scaevola concolor</i>	Dwarf neupaka	U.S.A. (HI)	E	231	NA	NA
Hydrophyllaceae—Waterleaf family: <i>Phacelia argillacea</i>	Clay phacelia	U.S.A. (UT)	E	44	NA	NA
<i>Phacelia formosula</i>	North Park phacelia	U.S.A. (CO)	E	121	NA	NA
Lamiaceae—Mint family: <i>Acanthomintha obovata</i> ssp. <i>duttonii</i>	San Mateo thornmint	U.S.A. (CA)	E	204	NA	NA
<i>Dicerandra cornutissima</i>	Longjapurred mint	U.S.A. (FL)	E	207	NA	NA
<i>Dicerandra lutescens</i>	Scrub mint	do	E	207	NA	NA
<i>Dicerandra immaculata</i>	Lakula's mint	do	E	180	NA	NA
<i>Haplostachya haplostachya</i> var. <i>angustifolia</i>	None	U.S.A. (HI)	E	73	NA	NA
<i>Hedeoma apiculatum</i>	McKintick pennyroyal	U.S.A. (TX, NM)	T	118	17.98(a)	NA
<i>Hedeoma tomentosum</i>	Todeen's pennyroyal	U.S.A. (NM)	E	110, 112	17.98(a)	NA
<i>Pogogyne abramis</i>	San Diego mesa mint	U.S.A. (CA)	E	44	NA	NA
<i>Scutellaria montana</i>	Large-flowered skullcap	U.S.A. (GA, TN)	E	234	NA	NA
<i>Stenogyne angustifolia</i> var. <i>angustifolia</i>	None	U.S.A. (HI)	E	73	NA	NA
Lauraceae—Laurel family: <i>Lindera melastrobata</i>	Ponuberry	U.S.A. (AL, AR, FL, GA, LA, MO, MS, NC, SC)	E	240	NA	NA
Liliaceae—Lily family: <i>Erythronium propullans</i>	Minnesota trout lily	U.S.A. (MN)	E	221	NA	NA
<i>Harperocalyx flava</i>	Harper's beauty	U.S.A. (FL)	E	67	NA	NA
<i>Trillium persistens</i>	Persistent trillium	U.S.A. (GA, SC)	E	39	NA	NA
Loasaceae—Loasa family: <i>Mentzelia leucophylla</i>	Ash Meadows blazing star	U.S.A. (NV)	T	181	17.98(a)	NA
Malvaceae—Mallow family: <i>Abutilon menziesii</i>	Ko'olalo'ula	U.S.A. (HI)	E	243	NA	NA
<i>Callirhoe scaberrimula</i>	Texas poppy-mallow	U.S.A. (TX)	E	109, 112	NA	NA
<i>Hibiscadelphus distans</i>	Kaula hau kushiki	U.S.A. (HI)	E	225	NA	NA
<i>Ilama corei</i>	Peter's Mountain mallow	U.S.A. (VA)	E	230	NA	NA
<i>Kotia cookii</i>	Cooke's kokio	U.S.A. (HI)	E	74	NA	NA
<i>Kotia drynarioides</i>	Koki'o (=hau-hale'ula or Hawaii tree cotton)	do	E	187	17.98(a)	NA
<i>Malacothamnus clementinus</i>	San Clemente Island bush-mallow	U.S.A. (CA)	E	28	NA	NA
<i>Sidaicea pedata</i>	Pedate checker mallow	do	E	158	NA	NA
Nyctaginaceae—Four-o'clock family: <i>Marattia macfarlandii</i>	MacFarlane's four-o'clock	U.S.A. (ID, OR)	E	68	NA	NA

U.S. Fish and Wildlife Serv., Interior

§ 17.12

Onagraceae—Evening-primrose family:							
<i>Camissonia benthamii</i>	San Benito evening-primrose.....	U.S.A. (CA).....	I	172	NA	NA	
<i>Oenothera avila</i> ssp. <i>eurekaensis</i>	Eureka Valley evening-primrose.....	do.....	E	39	NA	NA	
<i>Oenothera deltoides</i> ssp. <i>howellii</i>	Antioch Dunes evening primrose.....	do.....	E	39	17.96(a)	NA	
Orchidaceae—Orchid family:							
<i>Isoetes medeoloides</i>	Small whorled pogonia.....	U.S.A. (CT, IL, MA, MD, ME, MI, MO, NC, NH, NJ, NY, PA, RI, SC, VA, VT), Canada (Ont.).....	E	122	NA	NA	
<i>Spiranthes parviflora</i>	Navasota ladies'-tresses.....	U.S.A. (TX).....	E	116	NA	NA	
Papaveraceae—Poppy family:							
<i>Arctomecon humilis</i>	Dwarf bear-poppy.....	U.S.A. (UT).....	E	78	NA	NA	
Pinaceae—Pine family:							
<i>Abies guatemalensis</i>	Guatemalan fir (=phabele).....	Mexico, Guatemala, Honduras, El Salvador.....	T	84	NA	NA	
Poaceae—Grass family:							
<i>Tuctoria mucronata</i> (= <i>Orcuttia m.</i>).....	Solano grass.....	U.S.A. (CA).....	E	44	NA	NA	
<i>Panicum carteri</i>	Carter's panicgrass.....	U.S.A. (HI).....	E	133	17.96(a)	NA	
<i>Swallenia alexandriae</i>	Eureka Dune grass.....	U.S.A. (CA).....	E	39	NA	NA	
<i>Zizania texana</i>	Texas wild-rice.....	U.S.A. (TX).....	E	39	17.96(a)	NA	
Polygalaceae—Milkwort family:							
<i>Polypala arnoldii</i>	Tiny polygala.....	U.S.A. (FL).....	E	192	NA	NA	
Polygonaceae—Buckwheat family:							
<i>Eriogonum gypsophilum</i>	Gypsum wild buckwheat.....	U.S.A. (NM).....	T	110, 112	17.96(a)	NA	
<i>Eriogonum ovalifolium</i> var. <i>williamsiae</i>	Steamboat buckwheat.....	U.S.A. (NV).....	E	237	NA	NA	
<i>Eriogonum peltophilum</i>	Clay-loving wild-buckwheat.....	U.S.A. (CO).....	E	161	17.96(a)	NA	
Primulaceae—Primrose family:							
<i>Primula maguirei</i>	Maguire primrose.....	U.S.A. (UT).....	T	199	NA	NA	
Ranunculaceae—Buttercup family:							
<i>Aconitum noveboracense</i>	Northern wild monkshood.....	U.S.A. (IA, NY, OH, WI).....	T	39	NA	NA	
<i>Clematis socialis</i>	Alabama leather flower.....	U.S.A. (AL).....	E	245	NA	NA	
<i>Delphinium kinkense</i>	San Clemente Island larkspur.....	U.S.A. (CA).....	E	26	NA	NA	
Rhamnaceae—Buckthorn family:							
<i>Gouania hillebrandii</i>	None.....	U.S.A. (HI).....	E	165	17.96(a)	NA	

Rosaceae—Rose family:							
<i>Cowania subintegra</i>	Arizona cliffrose.....	U.S.A. (AZ).....	E	148	NA	NA	
<i>Ivesia eremica</i>	Ash Meadows Ivesia.....	U.S.A. (NV).....	T	181	17.96(a)	NA	
<i>Potentilla robbinsiana</i>	Robbins' cinquefoil.....	U.S.A. (NH, VT).....	E	104	17.96(a)	NA	
Rubiaceae—Coffee family:							
<i>Gardenia brighamii</i>	Na'u (Hawaiian gardenia).....	U.S.A. (HI).....	E	198	NA	NA	
Rutaceae—Citrus family:							
	St. Thomas prickly-ash.....	U.S.A. (PR, VI).....	E	213	NA	NA	

REFERENCE 15

Hanford Environmental Health Foundation

Drinking Water Report, HEHF-45

91118901123

HANFORD SANITARY WATER SYSTEMS

During CY-1984, sanitary water was supplied on the Hanford Site by 17 individual drinking water systems, each consisting of a raw water supply, treatment facilities, and distribution piping. All of the systems are operated by Department of Energy contractors with the exception of the City of Richland municipal system which provides water to the 700, 1100, and 3000 Areas. Twelve of the systems utilize Columbia River water as a raw water source, four systems utilize groundwater, and one system (Richland municipal) a combination of the two. The systems range in size from those providing plant-scale treatment and serving extensive areas to nine systems which supply water to single or small complexes of facilities only.

The systems, along with their sources of supply, service areas, and additional pertinent information, are listed below. A Hanford Site map depicting locations of the systems (with the exception of the Richland municipal system) is given in Figure 1.

<u>System</u>	<u>Source of Supply</u>	<u>Notes</u>
100-B	Columbia River via 181-B Pumphouse	Filtered and chlorinated. Sanitary water supplied to 182-B Bldg. only. System is Rockwell (RHO) operated.
100-D	Columbia River via 181-D Pumphouse	Filtered and chlorinated at 183-D. Treated water also supplied to 100-F and 100-H Areas as needed. System is RHO operated.
100-K	Columbia River via 181-K Pumphouse	Filtered and chlorinated. System is United Nuclear Corporation (UNC) operated.
100-N	Columbia River via 181-N Pumphouse	Filtered and chlorinated. System is UNC operated.
200-E	Columbia River via 100-B raw water export	Filtered and chlorinated at 283-E. System is RHO operated.

REFERENCE 16

Draft Phase I Installation Assessment of Inactive
Waste-Disposal Sites at Hanford, Volume 3

91118971125

SITE ID NO.: Pickling Acid Crib

ALIAS: None

FACILITY: Crib

STATUS: Inactive

ELEVATION:
WATERTABLE:

DIMENSIONS:

Length: 50 ft
Width: 30 ft
Depth: 10 ft
Diameter: 0

LOCATION: 600 Area
COORDINATES: To be determined

SITE DESCRIPTION:

A crib type structure 50 feet by 30 feet and 10 feet deep located 500 yards from the old White Bluffs Ice Plant and 300 yards east of Federal Avenue. Vent pipes extrude every 18 inches at the surface.

SERVICE DATES: 1943 - 1945

SERVICE HISTORY:

The site was used to pickle galvanized piping for use in the reactor buildings during construction. The process used several thousand gallons of nitric and hydrofluoric acid. The site was retired in 1945 and the surface was covered with large cobbles.

REFERENCES:

Documents:

Photographs: 122440-464-CN, 122440-465-CN

Drawings:

SITE ID NO.: Pickling Acid Crib

CHEMICALS DISPOSED

No chemical inventory is available.

RADIONUCLIDE INVENTORY
(in curies)

H-3:	0.00000	CE-144:	0.00000
C-14:	0.00000	PR-144:	0.00000
MN-54:	0.00000	PM-147:	0.00000
CO-60:	0.00000	EU-152:	0.00000
NI-63:	0.00000	EU-154:	0.00000
KR-85:	0.00000	EU-155:	0.00000
SR-90:	0.00000	NP-237:	0.00000
Y-91:	0.00000	PU-238:	0.00000
NB-95:	0.00000	PU-239:	0.00000
ZR-95:	0.00000	PU-240:	0.00000
TC-99:	0.00000	PU-241:	0.00000
RU-103:	0.00000	AM-241:	0.00000
RU-106:	0.00000	U-233:	0.00000
SN-113:	0.00000	U-235:	0.00000
SB-125:	0.00000	U-238:	0.00000
I-129:	0.00000	TH-232:	0.00000
CS-134:	0.00000	BETA:	0.00000
CS-137:	0.00000	GAMMA:	0.00000
CE-141:	0.00000	ALPHA:	0.00000

This site was not used to deliberately dispose of radioactive waste.

These values are decayed through April 1, 1986.

DIMENSIONS:

ELEVATION: 440 feet

WATERTABLE: 40 feet

Length: 450 feet
Width: 230 feet
Depth: 24 feet
Diameter:

FACILITY: Retention Basin

LOCATION: 100-B/C

COORDINATES: N71660/W80560, N71660/W80090, N71430/W80560, N71430/W80090

DESCRIPTION OF FACILITY

Concrete lined basin, 230 ft. x 467 ft., with a vertical baffle down the middle (lengthwise). The floor of the basin consists of concrete slabs, their joints originally closed with neoprene water seals. To a height of almost 10 ft. above the floor, the walls slope and are about 4 inches thick. The upper sections of the walls (about 10 ft) area vertical and range in thickness from about 5'8" at the bottom to one foot at the top.

The basin has been backfilled with soil to a depth of almost four feet.

DATE OF OPERATIONS: 1944-1968

DESCRIPTION OF WASTE

Retained cooling water effluent from the 105 reactor for radioactive decay and thermal cooling prior to release to the Columbia River. Total radionuclide inventories in the vicinity of the basins ranged from 5 to over 400 curies. 80% of the total radionuclide inventory is contained within the soil adjacent to the basins. Approximately 10 curies have leached into the concrete floor and walls.

UNPLANNED RELEASE

The following is a summary of unplanned releases in the basin area:

- In early 1952, gross leakage at the inlet for the 105-B effluent line was detected and steadily increased in volume.
- In late 1952, two known leaks from the effluent line occurred: (1) near the #2 diversion box for the 30" line and (2) near the 8" riser for the temporary by-pass line northeast of the 105-B Building.
- In February, 1954, a break occurred in the 107-B Basin.

EXTENT OF CONTAMINATION

The extent of the contamination from these releases is well within the zone encompassed by the retention basin and is within the AC-5-40 permanent posting.

*** MEMO FIELD EMPTY ***

REFERENCES:

Documents: UNI-946

Photographs:

Drawings:

9111890129

SITE ID NO.: 107-C

DIMENSIONS:

ELEVATION: 440 feet

WATERTABLE: 40 feet

Length:
Width:
Depth: 16 feet
Diameter: 330 feet

FACILITY: Retention Basins (2)

LOCATION: 100-B/C
COORDINATES: N-71045/W-79970, N-71045/W-80320

DESCRIPTION OF FACILITY

Two carbon steel tanks, each 330 feet in diameter and 16 feet deep. Each has a series of steel baffle plates inside to prevent water from channeling across the tank into the discharge line. Both tanks have been backfilled with soil to a depth of about four feet.

DATE OF OPERATIONS: 1942-1969

DESCRIPTION OF WASTE

Retained cooling water effluent from the 105 reactor for radioactive decay and thermal cooling prior to release to the Columbia River. Total radionuclide inventories in the vicinity of the basins ranged from 5 to over 400 curies. 80% of the total radionuclide inventory is contained within the soil adjacent to the basins. Approximately 10 curies have leached into the concrete floor and walls.

UNPLANNED RELEASE

The basin and its approximate 5 foot diameter effluent line has developed leaks during its operating life. It has been said the leaks could have been as high as 5,000 - 10,000 gallons per minute.

EXTENT OF CONTAMINATION

The extent of the contamination from these releases is well within the zone encompassed by the retention basin and is within the AC-5-40 permanent posting.

*** MEMO FIELD EMPTY ***

REFERENCES:

Documents: UNI-946

Photographs:

Ref. 16.5

Drawings: P-5242

911399131

SITE ID NO.: 107-D

DIMENSIONS:

ELEVATION: 420 feet

WATERTABLE: 35 feet

Length: 450 feet
Width: 230 feet
Depth: 24 feet
Diameter:

FACILITY: Retention Basin

LOCATION: 100 D-DR

COORDINATES: N-94455/W-52730 to N-94685/W-53250

DESCRIPTION OF FACILITY

A concrete-lined basin, 230 ft. x 467 ft., with 20 foot walls. The design is about the same as 107-B. the basin has been backfilled to a depth of about 2 ft. The walls appear to be coated with asphalt.

DATE OF OPERATIONS: 1944-1967

DESCRIPTION OF WASTE

Retained reactor cooling water effluent water from the 105 reactor for radioactive decay and thermal cooling prior to release to the Columbia River. Total radionuclide inventories in the vicinity of the basins ranged from 5 to over 400 curies. 70% of the total radionuclide inventory is contained within the soil adjacent to the basins. Approximately 10 curies have leached into the concrete floor and walls.

UNPLANNED RELEASE

The following is a summary of unplanned releases in the basin area:

- In early 1950, leakage occurred on the north side between the basin and the river. Effluent water had drained under the road to the section between the road and the perimeter fence.
- In the fall of 1951, excess leakage was detected above ground. As a result, two excavations were made. The holes were then covered, but effluent water continued to seep to the surface.

EXTENT OF CONTAMINATION

The extent of the contamination from these releases is well within the zone encompassed by the retention basin and is within the AC-5-40 permanent posting.

*** MEMO FIELD EMPTY ***

REFERENCES:

Documents: UNI-946

Photographs:

Drawings: M-1901-D Sh.8

91118901133

SITE ID NO.: 107-DR

DIMENSIONS:

ELEVATION: 420 feet

WATERTABLE: 35 feet

Length: 600 feet
Width: 230 feet
Depth:
Diameter:

FACILITY: Retention Basin

LOCATION: 100-D/DR

COORDINATES: N-94058/W52516, N94680/W52219

DESCRIPTION OF FACILITY

A concrete-lined basin, 230 ft. x 600 ft., with 20 foot walls. It has been backfilled to a depth of one to 3 feet.

DATE OF OPERATIONS: 1950-1964

DESCRIPTION OF WASTE

Retained cooling water effluent from the 105 reactor for radioactive decay and thermal cooling prior to release to the Columbia River. Total radionuclide inventories in the vicinity of the basins ranged from 5 to over 400 curies. 70% of the total radionuclide inventory is contained within the soil adjacent to the basins. Approximately 10 curies have leached into the concrete floor and walls.

UNPLANNED RELEASE

The following is a summary of unplanned releases in the basin area:

- In late 1951, extensive leakage of effluent water at the inlet end of the 107-DR retention basin was caused by the pipes pulling loose from the basin wall.

EXTENT OF CONTAMINATION

The extent of the contamination from these releases is well within the zone encompassed by the retention basin and is within the AC-5-40 permanent posting.

*** MEMO FIELD EMPTY ***

REFERENCES:

Documents: UNI-946

Photographs:

Drawings: M-1901-D Sh. 8

91118901135

SITE ID NO.: 107-F

DIMENSIONS:

ELEVATION: 400 feet

WATERTABLE: 30 feet

Length: 450 feet
Width: 230 feet
Depth: 24 feet
Diameter:

FACILITY: Retention Basin

LOCATION: 100-F

COORDINATES: N79967/W24130, N79967/W28900, N79500/W28900, N79500/W29130

DESCRIPTION OF FACILITY

A concrete-lined basin, 230 ft. x 467 ft., with 20 foot walls, similar in design to 100-B and 100-D. The basin has back-filled to a depth of about 5 feet, with soil piled to cover the walls.

DATE OF OPERATIONS: 1945-1965

DESCRIPTION OF WASTE

Retained cooling water effluent from the 105 reactor for radioactive decay and thermal cooling prior to release to the Columbia River. 70% of the total radionuclide inventory is contained within the soil adjacent to the basins. Approximately 10 curies have leached into the concrete floor and walls.

UNPLANNED RELEASE

EXTENT OF CONTAMINATION

The extent of the contamination from these releases is well within the zone encompassed by the retention basin and is within the AC-5-40 permanent posting.

*** MEMO FIELD EMPTY ***

REFERENCES:

Documents: UNI-946

Photographs:

Drawings: M-1600-F Sh.5

SITE ID NO.: 107-H

DIMENSIONS:

ELEVATION: 420 feet

WATERTABLE: 10 feet

Length: 600 feet
Width: 273 feet
Depth: 20 feet
Diameter:

FACILITY: Retention Basin

LOCATION: 100-H

COORDINATES: N96000/W38740, N96000/W38466, N95368/W38466, N95368/W38740

DESCRIPTION OF FACILITY

Concrete-lined rectangular basin, 273 ft. x 600 ft.; 20 feet deep. The basin has been back-filled to a depth of about 4 feet above the floor and slopes to the top of the walls.

DATE OF OPERATIONS: 1949-1965

DESCRIPTION OF WASTE

Retained cooling water effluent from the 105 reactor for radioactive decay and thermal cooling prior to release to the Columbia River. 70% of the total radionuclide inventory is contained within the soil adjacent to the basins. Approximately 10 curies have leached into the concrete floor and walls.

UNPLANNED RELEASE

The basin and its approximate 5 foot diameter effluent line has developed leaks during its operating life. It has been said the leaks could have been as high as 5,000 - 10,000 gallons per minute.

EXTENT OF CONTAMINATION

The extent of contamination from these releases is well within the zone encompassed by the retention basin and is within the AC-5-40 permanent posting.

*** MEMO FIELD EMPTY ***

REFERENCES:

Documents: UNI-946

Photographs:

Drawings: M-1600-H Sh.4

Ref. 16.12

SITE ID NO.: 107-KE

DIMENSIONS:

ELEVATION: 480 feet

WATERTABLE: 70 feet

Length:

Width:

Depth: 29 feet

Diameter: 250 feet

FACILITY: Retention Basins (3)

LOCATION: 100-K

COORDINATES: N(K)565699/W426802, N(K)532850/W453651, N(K)53000/W480500

DESCRIPTION OF FACILITY

Three carbon steel tanks, 250 ft. in diameter and 29 ft. deep. They have been backfilled with soil to a depth of about 4 feet.

DATE OF OPERATIONS: 1955-1971

DESCRIPTION OF WASTE

Retained cooling water effluent from the 105 reactor for radioactive decay and thermal cooling prior to release to the Columbia River. 80% of the total radionuclide inventory is contained within the soil adjacent to the basins. Approximately 10 curies have leached into the concrete floor and walls.

UNPLANNED RELEASE

The basin and its approximate 5 foot diameter effluent line has developed leaks during its operating life. The leak rate from the butterfly valves (that went to an adjacent trench) could have been as high as 5,000-10,000 gallons per minute. Most of the basin leakage was diverted to an open canal and disposed to the river.

EXTENT OF CONTAMINATION

The extent of the contamination from these releases is well within the zone encompassed by the retention basin and is within the AC-5-40 permanent posting.

*** MEMO FIELD EMPTY ***

REFERENCES:

Documents: UNI-946

Photographs:

Drawings: H-1-25529

Ref. 16.13

SITE ID NO.: 107-KW

DIMENSIONS:

ELEVATION: 480 feet

WATERTABLE: 70 feet

Length:

Width:

Depth: 29 feet

Diameter: 250 feet

FACILITY: Retention Basins (3)

LOCATION: 100-K

COORDINATES: N(K)5245/W(K)6190, N(K)5245/W(K)6460, N(K)5245/W(K)6730

DESCRIPTION OF FACILITY

Three carbon steel tanks, 250 ft. in diameter and 29 ft. deep. They have been backfilled with soil to a depth of about 4 feet.

DATE OF OPERATIONS: 1944-1970

DESCRIPTION OF WASTE

Retained cooling water effluent from the 105 reactor for radioactive decay and thermal cooling prior to release to the Columbia River. 80% of the total radionuclide inventory is contained within the soil adjacent to the basins. Approximately 10 curies have leached into the concrete floor and walls.

UNPLANNED RELEASE

The basin and its approximate 5 foot diameter effluent line has developed leaks during its operating life. The leak rate from the butterfly valves (that went to an adjacent trench) could have been as high as 5,000-10,000 gallons per minute. Most of the basin leakage was diverted to an open canal and disposed to the river.

EXTENT OF CONTAMINATION

The extent of the contamination from these releases is well within the zone encompassed by the retention basin and is within the AC-5-40 permanent posting.

*** MEMO FIELD EMPTY ***

REFERENCES:

Documents: UNI-946

Photographs:

Ref. 16.14

Drawings: H-1-24429

91118901140

Drawings: P-5242

91118901141

REFERENCE 17

File Memo regarding recreational use of the
Columbia River

9110001116

KH Cramer
RD Stenner
File/LB

Date August 26, 1987
To IWSS Files
From DR Sherwood *DRS*
Subject Recreational Use of Hanford Reach

The Hanford Reach of the Columbia River has many recreational uses. Sportsman's access for fishing and waterfowl, big game, and upland bird hunting is provided at several locations. Access points to the Columbia River along the Hanford Reach are located at the old White Bluffs Ferry landing, the Ringold Hatchery, and Leslie Grove Park in Richland. I have fished for steelhead and salmon along the Hanford Reach since 1980.

DRS/mgs

9 1 1 0 6 0 1 1 1 6

REFERENCE 18

Letter from RD Stenner to DM Bennett regarding 100 Area
Ground Water Contaminant Plume, October 14, 1987

91118901144



Battelle

Pacific Northwest Laboratories
P.O. Box 999
Richland, Washington U.S.A. 99352
Telephone (509)

Telex 15-2874

October 14, 1987

Mr. D. M. Bennett
U.S. Environmental Protection Agency
Region X
Superfund Program
1200 6th Avenue
Seattle, WA 98101

Dear Dave:

Enclosed are the three descriptions of the 100 Area, 200 Area and 300 Area ground water contaminant plumes we discussed on the telephone yesterday. I have included some attached figures and maps to help show the independency of the detected ground water contamination in each of the three aggregate areas.

If there are any questions regarding the descriptions, please contact me at 509-375-2916.

Sincerely,

Bab

R.D. Stenner, Sr. Research Engr.
Earth and Environmental Sciences Center
GEOSCIENCES DEPARTMENT

RDS:th

100 Area Ground Water Contamination

The ground water chromium and strontium-90 concentrations in the 100 Area are attributable only to the activities in the 100 Area because of the isolation of these areas and the fact that several years of ground water monitoring in the area show that any concentrations in ground water are found contained in close proximity to each of the individual reactor sites. Upstream samples of the surface water show that the contamination detected downstream are attributable to activity in the 100 Area. The furthest downstream location of contaminants from the 100 Area entering the river is at river mile 22. The furthest upstream entry point of the 100 Area contaminants is just above river mile 2, which is near the 100 B/C Area.

9119331145

15




FIGURE 3. Locations and Analytical Results for Spring and River Samples from River Mile 3 through River Mile 12

16




FIGURE 4. Locations and Analytical Results for Spring and River Samples from River Mile 14 through River Mile 22

REFERENCE 19

Letter from RD Stenner to DM Bennett regarding Liquid
Waste Sites and Burning Pits, October 26, 1987

91118901149



Pacific Northwest Laboratories
P.O. Box 999
Richland, Washington U.S.A. 99352
Telephone (509)

Telex 15-2874

October 26, 1987

D.M. Bennett
EPA/NPL Coordinator
EPA Superfund Program
U.S. Environmental Protection Agency Region X
1200 Sixth Ave.
Seattle, Washington 98101

Dear Dave:

Per your telephone request, enclosed are the statements on liquid waste sites and burning pits. It is our understanding that you will incorporate them as references in the appropriate sections of the NPL packages and provide us with a finalized copy of the packages following completion of the MITRE Corp. review.

If there are any questions regarding these statements, please don't hesitate to contact me at 509-375-2916.

Sincerely,

Bob

R.D. Stenner, Senior Research Engineer
Environmental Pathways and Assessment Section
Geosciences Department
Earth and Environmental Sciences Center

RDS:dar

ENCLOSURE

bcc: KH Cramer
MS Hanson
DA Lamar
RM Mangin - DOE/RL
TJ McLaughlin
RG Schreckhise
WB Schulze - DOE/RL
DR Sherwood

100, 200, 300 Area Statement Regarding Liquid Wastes

The general operating procedures for liquid waste sites in the 100, 200 and 300 Areas were such that the waste constituents listed for each site generally entered the process lines and were mixed with each other prior to being disposed of at the site. This process mixing of these waste constituents occurred over the period of site operation.

1
5
1
6
6
1
1
6

Burning Pit Statement

Due to the time period for which the burning pits operated, the nonhazardous combustible waste materials (i.e., paper products, cans, etc.) would have been mixed (i.e., mixed together in the garbage truck or waste container) with the hazardous waste materials (i.e., paints, solvents, etc.) prior to the waste mixture being disposed of at the site.

91110201172

Reference 20

Memo from Kathleen Galloway, MITRE, to Sandy Crystall, EPA
December 29, 1987

91118901153

